List of Tables

Table 2.1 Name of the fungi and accession number of ITS sequences
submitted to genebank, used in the present study. Similarity with sequences in
NCBI and Fusarium MLST database
Table 2.2 Similarity of TEF sequences of isolates used in present study with
existing sequences in NCBI and Fusarium MLST database71
Table 3.1: The genes investigated in current study, their known function and
primers used for qRT-PCR93
Table 3.2: Specific activity of extracted proteases from Fusarium spp. at
different pH using Azocasein assay99
Table 3.3: % residual activity of proteases in presence of inhibitors (EDTA,
PMSF and Pepstatin A) by using azocasein assay
Table 3.4: Protease identified from FSSC isolate Cs1 by HRLC-MS/MS
(orbitrap) in exoproteome and after column purification
Table 4.1: The genes investigated in the current study, their known functions,
and the primers used for quantitative real-time polymerase chain reaction 144
Table 4.2: Initial fungal dry weight and yield of crude β-glucan extracted from
Fusarium isolates
Table 4.3: Quantification of β -1,3-glucan and total β -glucan in <i>Fusarium</i>
isolates using ELISA and Congo red assay
Table 4.4: Functional group and their corresponding absorption frequencies in
FTIR spectra of β-glucan
Table 5.1: Summarised table for presence and absence of ZEA and T2 toxin in
Fusarium isolates based on TLC
Table 5.2: Concentration of ZEA toxin in crude toxin extracted from
pathogenic isolates
Table 6.1: Different solvent systems were used to separate and to characterize
the naphthoguinone pigment 219