

List of Figures

Figure 1 A quorum sensing model based on acyl homoserine lactone signalling systems	7
Figure 2 Structure of some representative AHLs	7
Figure 3 Soft rot caused due to quorum sensing in <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>	12
Figure 4 LasI/LasR and RhII/RhIR quorum sensing in <i>Pseudomonas aeruginosa</i>	14
Figure 5 Diversity of quorum quenching enzymes and inhibitors in different species of bacteria, Archaea and Eukaryote	18
Figure 6 Distribution of quorum quenching enzymes in various bacterial genera	32
Figure 7 Representative culture of <i>Bacillus</i> sp. isolated from root sample	41
Figure 8 Screening of isolates for C6-HSL degrading ability	41
Figure 9 Genotypic characterization of selected isolates	43
Figure 10 Dendrogram based on ARDRA analysis showing three major clusters of the twenty AHL degrading isolates	44
Figure 11 <i>B. subtilis</i> Pls8 inhibiting the growth of <i>PccBR1</i>	56
Figure 12 <i>B. aerius</i> Pls17 showing maceration on potato host tissue	57
Figure 13 Influence of <i>Bacillus</i> isolates on 3-oxo-C6HSL accumulation and growth of <i>PccBR1</i> in co-culture assays	58
Figure 14 Effect of AHL degrading <i>Bacillus</i> isolates on virulence enzymes production by <i>PccBR1</i>	59
Figure 15 <i>In vitro</i> soft rot attenuation assay on different host of <i>PccBR1</i>	61
Figure 16 <i>In vitro</i> curative biocontrol of potato soft rot caused by <i>PccBR1</i>	62
Figure 17 <i>In vitro</i> preventive biocontrol assay	63
Figure 18 <i>In planta</i> assay for biocontrol of spoilage of bean sprouts caused by <i>PccBR1</i> on susceptible variety of mung bean (<i>Vigna radiata</i>)	65
Figure 19 Strategy for cloning of adeH	78
Figure 20 AHL degrading enzyme in <i>Lysinibacillus</i> sp. Gs50 cells	83
Figure 21 Characterisation of AHL degrading enzyme of <i>Lysinibacillus</i> sp. Gs50	84
Figure 22 Cloning of gene responsible for AHL degradation in <i>Lysinibacillus</i> sp. Gs50	86

Figure 23 Expression of recombinant adeH in <i>E.coli</i> BL21(DE3)	87
Figure 24 Degradation of different chain length AHLs by <i>Lysinibacillus</i> sp. Gs50 and <i>E.coli</i> BL21(DE3) pET22b(+)/adeH	88
Figure 25 Soft rot attenuation assay of <i>E.coli</i> BL21 (DE3) pET22b(+)/adeH caused by <i>PccBR1</i>	88
Figure 26 Purification of active AdeH	89
Figure 27 AdeH is an AHL lactonase	90
Figure 28 Multiple sequence alignment of AdeH and other representative AHL lactonases	92
Figure 29 Neighbor-joining tree of AHL lactonases belonging to the metallo- β -lactamase, phosphotriesterase and α/β hydrolase-fold family based on amino acid sequences	93
Figure 30 Quantification of C6-HSL using Agar diffusion bioassay	94
Figure 31 Biochemical characterization of AdeH	96
Figure 32 Influence of temperature on quorum quenching activity of <i>Lysinibacillus</i> sp. Gs50	108
Figure 33 Influence of pH on quorum quenching activity of <i>Lysinibacillus</i> sp. Gs50	108
Figure 34 Influence of substrate concentration on quorum quenching activity of <i>Lysinibacillus</i> sp. Gs50	109
Figure 35 Comparison of <i>P. aeruginosa</i> PAO1 wild type and <i>P. aeruginosa</i> PAO1 (<i>lasI- rhII-</i>) treated mung bean plant growth parameters	112
Figure 36 CSLM images of validation of <i>P. aeruginosa</i> PAO1 strains	113
Figure 37 CSLM images of different plant root samples for visualizing active QS on mung bean roots	116
Figure 38 <i>In vitro</i> degradation of C4-HSL and 3-oxo-C12HSL by <i>Lysinibacillus</i> sp. Gs50	117
Figure 39 <i>In planta</i> virulence attenuation of <i>P. aeruginosa</i> PAO1 by quorum quenching <i>Lysinibacillus</i> sp. Gs50	119
Figure 40 CSLM micrograph of <i>P. aeruginosa</i> PAO1-gfp and <i>Lysinibacillus</i> sp. Gs50 co-inoculated plant root	120