

LIST OF TABLES

Table 1. 1 Distribution of Anopheles mosquito in India.....	10
Table 1. 2 Mouse and Human monocyte subsets (Shi et al., 2011).....	24
Table 1. 3 CC, C, CXC, and CX3C chemokine/receptor families (Borish et al., 2003)	26
Table 2. 1 Media composition	42
Table 2. 2 Antibiotics used for the study	42
Table 2. 3 Primer sequences of AMA1, MSP7 and WARP used for cloning.....	45
Table 2. 4 PCR system used for amplification of AMA1, MSP7 and WARP.....	45
Table 2. 5 PCR conditions used for amplification of (i) AMA1, (ii) MSP7 and (iii) WARP gene.....	46
Table 2. 6 Reagents and composition for competent cell preparation.....	47
Table 2. 7 Ligation system of AMA1 (PCR product) in pJET 1.2.....	48
Table 2. 8 Ligation system of AMA1 in pET30a (+)	48
Table 2. 9 Ligation system of MSP7 PCR product with pJET1.2 vector.	49
Table 2. 10 Ligation system of MSP7 in pET30a (+).....	49
Table 2. 11 Ligation system of WARP PCR product with pJET1.2.	50
Table 2. 12 Ligation system of WARP in pET30a (+)	50
Table 2. 13 Composition of reagents for plasmid isolation by alkaline lysis.	51
Table 2. 14 Restriction digestion system used for AMA1/pJET1.2.....	52
Table 2. 15 Restriction digestion system of AMA1/pET 30a (+).	53
Table 2. 16 Restriction digestion system for AMA1/pET30a (+).	53
Table 2. 17 Restriction digestion system for MSP7 /pJET 1.2	53
Table 2. 18 Restriction digestion system for MSP7/pET30a (+).	54
Table 2. 19 Restriction digestion system for MSP7/pET 30a	54

Table 2. 20 Restriction digestion system for WARP/pJET1.2.....	54
Table 2. 21 Restriction digestion system for WARP/pET30a (+).....	55
Table 2. 22 Restriction digestion system WARP/pET 30a (+) in putative clones.	55
Table 2. 23 Components and their composition used for 10% resolving gel.....	57
Table 2. 24 Components and their compositions used for 10% stacking gel.....	57
Table 2. 25 Components and their compositions used for silver staining.	58
Table 2. 26 Composition of lysis buffer 1 used for the solubilisation.....	60
Table 2. 27 Composition of lysis buffer 2 used to equilibrate the column.	60
Table 2. 28 Composition of wash buffer used to remove the unbound proteins present in the column.....	60
Table 2. 29 Composition of elution buffer used to elute out the protein that was bound to nickel beads present in the column.....	61
Table 2. 30 Primer sequences used for specific amplification of genes from <i>P. vivax</i> genomic DNA	62
Table 3. 1 Components and their composition used for the preparation of Bradford reagent.....	78
Table 3. 2 Composition of tank buffer used for western blot.....	79
Table 3. 3 Summary of western blot data.	91
Table 4. 1 Reaction system for semi-quantitative RT-PCR.....	109
Table 4. 2 RT-PCR (Reverse transcriptase) conditions for cDNA synthesis.....	109
Table 4. 3 Reaction system for β -actin PCR	110
Table 4. 4 System for Real time qPCR analysis of chemokine receptors.....	110
Table 4. 5 Primers used for qRT-PCR analysis.....	111
Table 4. 6 Reaction system for qRT-PCR.....	111