

References

- Angot, A.,** Peeters, N., Lechner, E., Vailleau, F., Baud, C., Gentzbittel, L., Sartorel, E., Genschik, P., Boucher, C. and Genin, S. (2006) *Ralstonia solanacearum* requires F-box-like domain-containing type III effectors to promote disease on several host plants. *Proc. Natl. Acad. Sci., USA* **103**: 14620–14625.
- Adachi, K.** and Hamer, J. E. (1998) Divergent cAMP signaling pathways regulate growth and pathogenesis in the rice blast fungus *Magnaporthe grisea*. *The Plant Cell* **10**: 1361–1374.
- Ariza, R. R.,** Keyse, S. M., Moggs, J. G. and Wood, R. D. (1996) Reversible protein phosphorylation modulates nucleotide excision repair of damaged DNA by human cell extracts. *Nucl. Acids Res.* **24**: 433–440.
- Bai, C.,** Sen, P., Hofmann, K., Ma, L. and Goebel, M. (1996) SKP1 connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell* **86**: 263–274.
- Bonman, J. M.** (1992) Durable resistance to rice blast disease-environmental influences. *Euphytica* **63**: 115–123.
- Bonman, J.M.** and Mackil, A. (1988) Effects of water deficit on rice blast. II. Disease development in field. *J. Plant Prot.* **5**: 67–73.

- Bolker**, M., Böhnert H. U., Braun K. H., Gorl J., and Kahmann R. (1995) Tagging pathogenicity genes in *Ustilago maydis* by restriction enzyme-mediated integration (REMI). *Mol. Gen. Genet.* **248**:5: 547–552.
- Bradford**, M. M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 248–254.
- Brace**, E. J., Parkinson L. P. and Fuller R. S. (2006) Skp1p regulates Soi3p/Rav1p association with endosomal membranes but is not required for vacuolar ATPase assembly. *Eukaryotic Cell.* **5**: 2104–2113.
- Brown**, J. S. and Holden D. W. (1998) Insertional mutagenesis of pathogenic fungi. *Curr. Opin. Microbiol.* **1**:4: 390–394.
- Burger**, A. M. and Seth, A. K. (2004) The ubiquitin-mediated protein degradation pathway in cancer: therapeutic implications. *Eur. J. Cancer.* **40**(15): 2217-29.
- Chen**, D. C., Yang, B. C. and Kuo, T. T. (1992) One-step transformation of yeast in stationary phase. *Curr. Genet.* **21**(1): 83-4.
- Choi**, W. and Dean, R. A. (1997) The adenylate cyclase gene MAC1 of *Magnaporthe grisea* controls appressorium formation and other aspects of growth and development. *The Plant Cell* **9**: 1973–1983.

- Ciechanover, A.** (1998) The ubiquitin-proteasome pathway: on protein death and cell life. *EMBO J.* **17(24)**: 7151-60.
- Connelly, C.** and Hieter, P. (1996) Budding yeast SKP1 encodes an evolutionarily conserved kinetochore protein required for cell cycle progression. *Cell* **86**: 275–285.
- Daniel, R.G.** and Wood, R.A. (2002) Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* **350**: 87–96.
- De Groot, M. J., Bundock P., Hooykaas P. J., and Beijersbergen A. G.** (1998) *Agrobacterium*-mediated transformation of filamentous fungi. *Nat. Biotechnol.* **16.9**: 839–842.
- DeBacker, M. D., Raponi, M. and Arndt, G. M.** (2002) RNA mediated gene silencing in non pathogenic and pathogenic fungi. *Curr. Opin. Microbiol.* **5**: 332-339.1
- Duyvesteijn, R. G. E., Wijk, R., Boer, Y., Rep, M. and Cornelissen, B.J.C.** (2005) Frp1 is a *Fusarium oxysporum* F-box protein required for pathogenicity on tomato. *Mol. Microbiol.* **57**: 1051–1063.
- Ebbole, D. J.** (2007) *Magnaporthe* as a model for understanding host-pathogen interactions. *Annu. Rev. Phytopathol.* **45**: 437–456.

- Fire, A., Xu S., Montgomery, M. K., Kostas, S. A., Driver, S. E., and Mello C. C. (1998)** Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*. **391.6669**: 806–811.
- Fitzgerald, A. M., Mudge A. M., Gleave A. P., and Plummer K. M. (2003)** *Agrobacterium* and PEG-mediated transformation of the phytopathogenic fungus *Venturia inaequalis*. *Mycol. Res.* **107**: 803–810.
- Fourrey, V. C., Barooah, M., Egan, M., Wakley, G. and Talbot, N.J. (2006)** Autophagic fungal cell death is necessary for infection by the rice blast fungus. *Science* **312**: 580–583.
- Feldman, R. M., Correll, C. C., Kaplan, K. B. and Deshaies, R. J. (1997)** A complex of Cdc4p, Skp1p, and Cdc53p/cullin catalyzes ubiquitination of the phosphorylated CDK inhibitor Sic1p. *Cell* **91**: 221–230.
- Gabert, J., Beillard, E., van der Velden, V. H. J., Bi, W., Grimwade, D., Pallisgaard, N., Barbany, G., Cazzaniga, G., Cayuela, J. M., Cavé, H., Pane, F., Aerts, J. L. E., De Micheli, D., Thirion, X., Pradel, V., González, M., Viehmann, S., Malec, M., Saglio, G. and van Dongen, J. J. M. (2003)** Standardization and quality control studies of 'real-time' quantitative reverse transcriptase polymerase chain reaction of fusion gene transcripts for residual disease detection in leukemia – A Europe Against Cancer Program. *Leukemia*. **17**: 2318–2357.

- Gilbert**, M.J., Thornton, C.R., Wakley, G.E. and Talbot, N.J. (2006) A P-type ATPase required for rice blast disease and induction of host resistance. *Nature* **440**: 535–539.
- Gilchrist**, E. J. and Haughn G. W. (2005) TILLING without a plough: a new method with applications for reverse genetics. *Curr. Opin. Plant Biol.* **8.2**:211–215.
- Goff**, S. A. (1999) Rice as model for cereal genomics. *Curr. Opin. Plant. Biol.* **2**: 86-89.
- Goto**, K., Kanazawa, A. and Kusaba, M. (2003) A Simple and Rapid Method to Detect Plant siRNAs Using Nonradioactive Probes. *Plant Mol. Biol. Rep.* **21**: 51–58.
- Gupta**, A. and Chattoo, B.B. (2007) A novel gene MGA1 is required for appressorium formation in *Magnaporthe grisea*. *Fungal Genet. Biol.* **44**: 1157–1169.
- Hattori**, T., Kitagawa, K., Uchida, C., Oda, T. and Kitagawa, M. (2003) Cks1 is degraded via the ubiquitin-proteasome pathway in a cell cycle-dependent manner. *Genes Cells* **8**: 889–896.
- Hammond**, S. M., Boettcher S., Caudy, A. A., Kobayashi, R. and Hannon, G. J. (2002) Argonaute2, a link between genetic and biochemical analyses of RNAi. *Science*. **293.5532**: 1146–1150.
- Hannon** G. J. (2002) RNA interference. *Nature*. **418**: 244–251.

- He**, M., Kershaw, M. J., Soanes, D.M., Xia, Y. and Talbot, N. J. (2012) Infection-associated nuclear degeneration in the rice blast fungus *Magnaporthe oryzae* requires non-selective macro-autophagy. *PloS One* **7**: e33270.
- Hermant**, D., Bamps, S., Tafforeau, L., Vandenhoute, J. and Makela, T. P. (2003) Skp1 and the F-box protein Pof6 are essential for cell separation in fission yeast. *J. Biol. Chem.* **278**: 9671–9677.
- Hieter**, P. and Boguski, M. (1997) Functional genomics: It's all how you read it. *Science*. **278**: 601-602.
- Irie**, T., Matsumura, H., Terauchi, R. and Saiton, H. (2003) Serial analysis of gene expression (SAGE) of *Magnaporthe grisea*: Genes involved in appressorium formation. *Mol. Genet. Genomics*. **270**: 181-189.
- Jackson** A. L., Pahl P. M. B., Harrison K., Rosamond J. and Sclafani R. A. (1993) Cell cycle regulation of the yeast Cdc7 protein Kinase by association with Dbf4 protein. *Mol. Cell. Biol.* **13**: 2899-2908.
- James** S. W., Bullock, K. A., Gyga, S. E., Kraynack, B. A., Matura, R. A., MacLeod, J. A., McNeal, K. K., Prasauckas, K. A., Scacheri, P. C., Shenefiel, H. L., Tobin, H. M. and Wade, S. D. (1999) nimO, an *Aspergillus* gene related to budding yeast Dbf4, is required for DNA synthesis and mitotic checkpoint control. *J. Cell Sci.* **112**: 1313-1324.

- Jantasuriyarat**, C., Gowda, M., Haller, K., Hatfield, J., Lu, G., Stahlberg, E., Zhou, B., Li, H., Kim, H., Yu, Y., Dean, R. A., Wing, R. A., Soderlund, C. and Wang, G. (2005) Large scale identification of expressed sequence tags involved in Rice and Rice blast fungus interaction. *Plant. Physiol.* **138**(1): 105-115.
- Jeon**, J., Park S. Y. and Chi M. H. (2007) Genome-wide functional analysis of pathogenicity genes in the rice blast fungus. *Nat. Genet.* **39**.4: 561–565.
- Jia**, L. and Sun, Y. (2011) SCF E3 ubiquitin ligases as anticancer targets. *Curr. Cancer Drug Targets.* **11**(3):347-56.
- Jin**, Q., Dong, H., Peng, Y., Chen, B., Shao, J., Deng, Y., Dai, C., Fang, Y., Lou, Y., Li, Y. and Li, D. (2007) Application of cDNA array for studying the gene expression profile of mature appressorium of *Magnaporthe grisea*. *J Zhejiang Univ Sci B.* **8**(2): 88-97.
- Jones**, D. T., Taylor, W. R. and T. J. M. (1992) The rapid generation of mutation data matrices from protein sequences. *Comput. Appl. Biosci.* **8**: 275–282.
- Kachroo**, P., Leong, S.A. and Chattoo, B. B. (1994) Pot2, an inverted repeat transposon from the rice blast fungus *Magnaporthe grisea*. *Mol. Gen. Genet.* **245**: 339–348.
- Kachroo**, P., Leong, S. A. and Chattoo, B. B. (1995) Mg-SINE: a short interspersed nuclear element from the rice blast fungus, *Magnaporthe grisea*. *Proc. Natl. Acad. Sci. USA* **92**: 11125–11129.

- Kadotani**, N., Nakayashiki H., Tosa Y., and Mayama S. (2003) RNA silencing in the phytopathogenic fungus *Magnaporthe oryzae*. *Mol. Plant Microb. Interact.* **16.9**: 769–776.
- Kamakura**, T., Xiao, J., Choi, W., Kochi, T., Yamaguchi, S., Teraoka, T. and Yamaguchi, I. (1999) cDNA subtractive cloning of gene expressed during early stage of appressorium formation by *Magnaporthe grisea*. *Biosci. Biotechnol. Biochem.* **63**: 1407-1413.
- Kaplan**, K. B., Hyman, A. A. and Sorger, P. K. (1997) Regulating the yeast kinetochore by ubiquitin-dependent degradation and Skp1p-mediated phosphorylation. *Cell* **91**: 491–500.
- Khang**, C. H., Park S. Y., Lee Y. H., and Kang S. (2005) A dual selection based, targeted disruption tool for *Magnaporthe grisea* and *Fusarium oxysporum*. *Fungal Genet. Biol.* **42.6**: 483–492.
- Kim**, D. E., Chivian, D. and Baker, D. (2004) Protein structure prediction and analysis using robetta server. *Nucleic Acids Res.* **32**: DOI: 10.1093/nar/gkh468.
- Kim**, S., Ahn, I. P. and Lee, Y. H. (2001) Analysis of gene expressed during Rice-*Magnaporthe grisea* interaction. *Mol. Plant. Microb. Interact.* **14**: 1340-1346.
- Kim**, N., Yoon, H., Lee, E. and Song, K. (2006) A new function of Skp1 in the mitotic exit of budding yeast *Saccharomyces cerevisiae*. *J. Microbiol. (Seoul, Korea)* **44**: 641–648.

- Koc**, A., Wheeler, L. J., Mathews, C. K. and Merrill, G. F. (2004) Hydroxyurea arrests DNA replication by a mechanism that preserves basal dNTP pools. *J. Biol. Chem.* **279**: 223–230.
- Komari**, T., Hiei, Y., Saito, Y., Murai, N. and Kumashiro, T. (1996) Vectors carrying two separate T-DNAs for co-transformation of higher plants mediated by *Agrobacterium tumefaciens* and segregation of transformants free from selection markers. *Plant J.* **10**: 165–174
- Lehmann**, A. and Toda, T. (2004) Fission yeast Skp1 is required for spindle morphology and nuclear membrane segregation at anaphase. *FEBS Lett.* **566**: 77–82.
- Lehmann**, A., Katayama, S., Harrison, C., Dhut, S. and Kitamura, K. (2004) Molecular interactions of fission yeast Skp1 and its role in the DNA damage checkpoint. *Genes Cells* **9**: 367–382.
- Lamour**, K. H., Finley L., Hurtado-Gonzales O., Gobena D., Tierney M., and Meijer H. J. G. (2006) Targeted gene mutation in *Phytophthora* spp. *Mol. Plant Microb. Interact.* **19.12**: 1359–1367.
- Lindsay**, D. S., Lenz S. D., Cole R. A., Dubey J. P., and Blagburn B. L. (1995) Mouse model for central nervous system *Neospora caninum* infections. *J. Parasitol.* **81.2**: 313–315.
- Liu**, T. B. and Xue, C. (2011) The Ubiquitin-Proteasome System and F-box Proteins in Pathogenic Fungi. *Mycobiology.* **39(4)**: 243-8.

- Latijnhouwers, M., Ligterink W., Vleeshouwers V. G. A. A., West P., and Govers F.**
(2004) A Galpha subunit controls zoospore motility and virulence in the potato late blight pathogen *Phytophthora infestans*. *Mol. Microbiol.* **51**: 925–936.
- Lyzena, W. J. and Stone, S. L.** (2011) Protein ubiquitination: An emerging theme in plant abiotic stress tolerance. *The Americans Journal of Plant Science and Biotechnology.* **5(2)**: 1-11.
- Lyzena, W. J. and Stone, S. L.** (2012) Abiotic stress tolerance mediated by protein ubiquitination. *J. Exp. Bot.* **63(2)**: 599-616.
- Magori, S. and Citovsky, V.** (2011) Hijacking of the Host SCF Ubiquitin Ligase Machinery by Plant Pathogens. *Front. Plant Sci.* **2**: 87.
- McCallum, C. M., Comai L., Greene E. A., and Henikoff S.** (2000) Targeting induced local lesions in genomes (TILLING) for plant functional genomics. *Plant Physiol.* **123.2**: 439–442.
- Maier, F. J. and Schafer W.** (1999) Mutagenesis via insertional or restriction enzyme-mediated-integration (REMI) as a tool to tag pathogenicity related genes in plant pathogenic fungi. *Biol. Chem.* **380.7-8**: 855–864.
- Meinke, D. W.** (1995) Molecular genetics of plant embryogenesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **46**: 369–394.

- Mercure**, E. W., Kunoh, H. and Nicholson, R. L. (1995) Visualization of materials released from adhered ungerminated conidia of *Colletotrichum graminicola*. *Physiol. Mol. Plant. Pathol.* **46**: 121-135.
- Mew**, T. (1991) Disease management in rice. CRC handbook of pest management in Agriculture. (2nd Ed.) V-III, D. Pimental ed. CRC press, Inc. Boca Raton, Ann Arbor, Boston. Biological studies. *Curr. Harm. Des.* **5**: 73-99.
- Michielse**, C. B., Hooykaas P. J. J., Van den Hondel C. A., and Ram A. F. J. (2005) *Agrobacterium*-mediated transformation as a tool for functional genomics in fungi. *Curr. Genet.* **48.1**: 1–17.
- Michielse**, C. B., Hooykaas P. J., Van den Hondel C. A., and Ram A. F. J. (2008) *Agrobacterium*-mediated transformation of the filamentous fungus *Aspergillus awamori*. *Nat. Protoc.* **3.10**: 1671–1678.
- Mitchell**, T. K. and Dean R. A. (1995) The cAMP-dependent protein kinase catalytic subunit is required for appressorium formation and pathogenesis by the rice blast pathogen *Magnaporthe grisea*. *Plant Cell.* **7.11**: 1869–1878.
- Moon**, J., Parry, G. and Estelle, M. (2004) The ubiquitin proteasome pathway and plant development. *Plant Cell.* **16**: 3181-3195.
- Mullins**, E. D., Chen X., Romaine P., Raina R., Geiser D. M., and Kang S. (2001) *Agrobacterium*-mediated transformation of *Fusarium oxysporum*: an efficient tool for insertional mutagenesis and gene transfer. *Phytopathology.* **91.2**: 173–180.

- Murray**, A.W. (2004) Recycling the cell cycle: cyclins revisited. *Cell* **116**: 221–234.
- Nakayama**, K. I. and Nakayama, K. (2006) Ubiquitin ligase: Cell cycle control and cancer. *Nat. Rev. Cancer*. **6**(5): 369-81.
- Nakayashiki**, H. (2005) RNA silencing in fungi: mechanisms and applications. *FEBS Lett.* **579**: 5950–5957.
- Nakayashiki**, H. and Nguyen, Q.B. (2008) RNA interference: roles in fungal biology. *Curr. Opin. Microbiol.* **11**:494-502.
- Nasmyth**, K. F. C. (1993) Yeast G1 cyclins CLN1 and CLN2 and a GAP-like protein have a role in bud formation Fatima. *EMBO J.* **1**: 5277–5286.
- Naujokat**, C. and Hoffmann, S. (2002) Role and function of the 26S proteasome in proliferation and apoptosis. *Lab Invest.* **82**(8): 965-80.
- Nguyen**, Q.B., Kadotani, N., Kasahara, S., Tosa, Y. and Mayama, S. (2008) Systematic functional analysis of calcium-signalling proteins in the genome of the rice-blast fungus, *Magnaporthe oryzae*, using a high-throughput RNA-silencing system. *Mol. Microbiol.* **68**: 1348–1365.
- Ogino**, K., Takeda, T., Matsui, E., Iiyama, H., Taniyama, C., Arai, K. and Masai, H. (2001) Bipartite Binding of a Kinase Activator Activates Cdc7-related Kinase Essential for S Phase. *J. Biol. Chem.* **276**: 31376–31387.

- Oh**, Y., Franck, W.L., Han, S.O., Shows, A. and Gokce, E. (2012) Polyubiquitin is required for growth, development and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *PloS One*. **7**: e42868.
- Ou**, S. H. (1985) Rice disease. Second ed., Commonwealth Mycological Institute Publication, Kew, Surrey. U.K. 280-282.
- Park**, G., Xue, C., Zhao, X., Kim, Y. and Orbach, M., (2006) Multiple upstream signals converge on the adaptor protein Mst50 in *Magnaporthe grisea*. *Plant Cell*. **18**: 2822–2835.
- Pukatzki**, S., Tordilla, N., Franke, J. and Kessin, R. H. (1998) A novel component involved in ubiquitination is required for development of *Dictyostelium discoideum*. *J. Biol. Chem.* **273**: 24131-24138.
- Rao**, K. M. (1994) Rice Blast Disease. Daya Publishing House, Delhi, India. 1-2.
- Rauyaree**, R., Choi, W., Fang, E., Blackmon, B. and Dean, R. A. (2001) Gene expressed during early stage of rice infection with the rice blast fungus *Magnaporthe grisea*. *Mol. Plant. Pathol.* **2**: 347-354.
- Redman**, R. S. and Rodriguez R. J. (1994) Factors affecting the efficient transformation of *Colletotrichum* species. *Exp. Mycol.* **18.3**: 230–246.
- Renski**, L., Clijsters, L. and Wolthuis, R. (2008) To cell cycle, swing the APC/C. *Biochem. Biophys. Acta* **1786**: 49–59.

- Rho**, H. S., Kang S., and Lee Y. H. (2001) *Agrobacterium tumefaciens*-mediated transformation of the plant pathogenic fungus, *Magnaporthe grisea*. *Molecules and cells*. **12.3**: 407–411.
- Riggle**, P. J. and Kumamoto C. A. (1998) Genetic analysis in fungi using restriction-enzyme-mediated integration. *Curr. Opin. Microbiol.* **1.4**: 395–399.
- Rolland**, S., Jobic C., Fevre M., and Bruel C. (2003) *Agrobacterium*-mediated transformation of *Botrytis cinerea*, simple purification of monokaryotic transformants and rapid conidia-based identification of the transfer-DNA host genomic DNA flanking sequences. *Curr. Genet.* **44.3**: 164-171.
- Rosebrock**, T. R., Zeng, L., Brady, J. J., Abramovitch, R. B., Xiao, F. and Martin, G. B. (2007) A bacterial E3 ubiquitin ligase targets a host protein kinase to disrupt plant immunity. *Nature*. **448**: 370-374.
- Rossman**, A. Y., Haward, R. J. and Valent, B. (1990) *Pyricularia grisea*, the current name for the rice blast fungus. *Mycologia*. **82**: 509-512.
- Salame**, T. M., Ziv, C., Hadar, Y. and Yarden, O. (2011) RNAi as a potential tool for biotechnological applications in fungi. *Appl. Microbiol. Biotechnol.* **89**: 501–512.
- Sambrook**, J., Fritsch, E.F. and Maniatis, T. (1989) Molecular cloning: A laboratory manual II. New York.

- Sara**, L. T. and Talbot, N. J. (2001) Surface attachment and pre-penetration stage development by plant pathogenic fungi. *Annu. Rev. Phytopathol.* **39**: 385-417.
- Saunders**, D. G. O., Aves, S. J. and Talbot, N. J. (2010) Cell cycle-mediated regulation of plant infection by the rice blast fungus. *Plant Cell.* **22**: 497–507.
- Saunders**, D. G. O., Dagdas, Y. F. and Talbot, N. J. (2010) Spatial uncoupling of mitosis and cytokinesis during appressorium-mediated plant infection by the rice blast fungus *Magnaporthe oryzae*. *Plant Cell.* **22**: 2417–2428.
- Schulman**, B. A., Carrano, A. C., Jeffrey, P. D., Bowen, Z. and Kinnucan, E. R. (2000) Insights into SCF ubiquitin ligases from the structure of the Skp1-Skp2 complex. *Nature.* **408**: 381–386.
- Scherer**, S., and Davis R. W. (1979) Replacement of chromosome segments with altered DNA sequences constructed in vitro. *Proc. Natl. Acad. Sci. USA.* **76.10**: 4951–4955.
- Schiestl**, R. H. and Petes T. D. (1991) Integration of DNA fragments by illegitimate recombination in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA.* **88.17**: 7585–7589.
- Seol**, J. H., Shevchenko, A. and Deshaies, R. J. (2001) Skp1 forms multiple protein complexes, including RAVE, a regulator of V-ATPase assembly. *Nat. Cell Biol.* **3**: 384–391.

- Shaner**, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N., Palmer, A. E. and Tsien, R. Y. (2004) Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat. Biotechnol.* **22(12)**: 1567-72.
- Shi**, Z., Christian D., and Leung H. (1995) Enhanced transformation in *Magnaporthe grisea* by restriction enzyme mediated integration of plasmid DNA. *Phytopathology*. **85.3**: 329–333.
- Siino**, J. S., Yau, P. M., Imai, B. S., Gatewood, J. M. and Morton, B. E. (2003) Effect of DNA length and H4 acetylation on the thermal stability of reconstituted nucleosome particles. *Biochem. Biophys. Res. Commun.* **302**: 885–891.
- Silva**, M. C. (2004) Signaling pathway in appressorium formation in *Magnaporthe grisea*. (dissertation). College Station: Texas A&M University.
- Singer**, A. U., Schulze, S., Skarina, T., Xu, X., Cui, H., Lippold, L. E., Egler, M., Sri Kumar, T., Raught, B., Lee, J., Scheel, D., Savchenko, A. and Bonas, U. A Pathogen Type III Effector with a Novel E3 Ubiquitin Ligase Architecture. *Plos Pathol. PLoS Pathog.* **9(1)**: e1003121. doi:10.1371/journal.ppat.1003121.
- Singer**, R.A. and Johnston, G.C. (1981) Nature of the G1 phase of the yeast *Saccharomyces cerevisiae*. *Cell Biol.* **78**: 3030–3033.
- Skamnioti**, P. and Gurr, S.J. (2009) Against the grain: safeguarding rice from rice blast disease. *Trends Biotech.* **27**: 141–150.

- Skowyra, D.,** Craig, K. L., Tyers, M., Elledge, S. J. and Harper, J. W. (1997) F-box proteins are receptors that recruit phosphorylated substrates to the SCF ubiquitin-ligase complex. *Cell*. **91**: 209–219.
- Smalle, J. and Vierstra, R. D.** (2004) The ubiquitin 26S proteasome proteolytic pathway. *Annu. Rev. Plant Biol.* **55**: 555-90.
- Staszczak, M.** (2008) The role of the ubiquitin-proteasome system in the response of the ligninolytic fungus *Trametes versicolor* to nitrogen deprivation. *Fungal Genet. Biol.* **45**(3): 328-37.
- Stemmann, O. and Lechner, J.** (1996) The *Saccharomyces cerevisiae* kinetochore contains a cyclin-CDK complexing homologue, as identified by *in vitro* reconstitution. *EMBO J.* **15**: 3611–3620.
- Sweigard, J. A., Carroll A. M., Farrall L., Chumley F. G. and Valent B.** (1998) *Magnaporthe grisea* pathogenicity genes obtained through insertional mutagenesis. *Mol. Plant Microbe Interact.* **11**: 404-12.
- Talbot, N. J.** (2003) On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*. *Annu. Rev. Microbiol.* **57**: 177–202.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S.** (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. Evol.* **28**: 2731–2739.

- Takano, Y.**, Choi, W., Mitchell, T. K., Okuno, T. and Dean, R. A. (2003) Large scale parallel analysis of gene expression during infection related morphogenesis of *Magnaporthe grisea*. *Mol. Plant. Pathol.* **4(5)**: 337-346.
- Tendulkar, S. R.**, Gupta, A. and Chattoo, B.B. (2003) A Simple protocol for isolation of fungal DNA. *Biotechnol. Lett.* **25**: 1941-1944.
- Thakur, S.**, Jha, S., Barman, R. S. and Chattoo, B.B. (2009) Genomic resources of *Magnaporthe oryzae* (GROMO): a comprehensive and integrated database on rice blast fungus. *BMC Genomics.* **10**: 316.
- Thines, E.**, Weber, R. W. and Talbot, N. J. (2000) MAP kinase and protein kinase A-dependent mobilization of triacylglycerol and glycogen during appressorium turgor generation by *Magnaporthe grisea*. *Plant Cell.* **12**: 1703–1718.
- Timberlake, W. E.**, and Marshall, M. A. (1989) “Genetic engineering of filamentous fungi” *Science.* **244**: 1313–1317.
- Tu, Y.**, Chen, C., Pan, J., Xu, J., Zhou, Z. G. and Wang, C. Y. (2012) The ubiquitin proteasome pathway (UPP) in the regulation of cell cycle control and DNA damage repair and it's implication in tumerogenesis. *Int. J. Clin. Exp. Pathol.* **5(8)**: 726-738.
- Valent, B.**, Farrall, L. and Chumley, F. G. (1990) *Magnaporthe grisea* genes for pathogenicity and virulence identified through a series of Backcross. *Genetics.* **127**: 87-107.

- van der Velden**, V. H. J., Hochhaus, A., Cazzaniga, G., Szczepanski, T., Gabert, J. and van Dongen, J. J. M. (2003) Detection of minimal residual disease in hematologic malignancies by real-time quantitative PCR: principles, approaches, and laboratory aspects. *Leukemia*. **17**: 1013–1034.
- van West**, P., Kamoun S., Van't Klooster J. W., and Govers F. (1999) Internuclear gene silencing in *Phytophthora infestans*. *Molec. Cell*. **3.3**: 339–348.
- Vijn**, I., and Govers F. (2003) *Agrobacterium tumefaciens*-mediated transformation of the oomycete plant pathogen *Phytophthora infestans*. *Mol. Plant Pathol*. **4.6**: 459–467.
- Vodermaier**, H.C. (2004) APC/C and SCF: controlling each other and the cell cycle. *Curr. Biol*. **14**: R787–96.
- Wang**, F. and Deng, X. W. (2011) Plant ubiquitin pathway and its role in gibberellin signalling. *Cell Res*. **21(9)**: 1286-1294.
- Wang**, Z. Y., Jenkinson, J. M., Holcombe, L. J., Soanes, D. M. and Fourrey, V. C. (2005) The molecular biology of appressorium turgor generation by the rice blast fungus *Magnaporthe grisea*. *Biochem. Soc. Transact*. **33**: 384–388.
- Waterhouse**, P. M., Graham M. W., and Wang M. (1998) Virus resistance and gene silencing in plants can be induced by simultaneous expression of sense and antisense RNA. *Proc. Natl. Acad. Sci. USA*. **95.23**: 13959–13964.

- Weber**, R. W. S., Wakley, G. E. and Talbot, N. J. (2001) The vacuole as central element of the lytic system and sink for lipid droplets in maturing appressoria of *Magnaporthe grisea*. *Protoplasma*. **216**: 101-112.
- Wendland**, J. (2003) PCR-based methods facilitate targeted gene manipulations and cloning procedures. *Curr. Genet*. **44**: 115–123.
- West**, P., Kamoun, S., Klooster, J. W. and Gover, F. (1999) Internuclear gene silencing in *Phytophthora infestans*. *Mol. Cell*. **3**: 339-348.
- Whisson**, S. C., Avrova A. O., West P. V., and Jones J. T. (2005) A method for double-stranded RNA-mediated transient gene silencing in *Phytophthora infestans*. *Mol. Plant Pathol*. **6.2**: 153–163.
- Wilson**, R.A. and Talbot, N.J. (2009) Under pressure: investigating the biology of plant infection by *Magnaporthe oryzae*. *Nat. Rev. Microbiol*. **7**: 185–195.
- Xu**, J. R. and Hamer, J. E. (1996) MAP kinase and cAMP signaling regulate infection structure formation and pathogenic growth in the rice blast fungus *Magnaporthe grisea*. *Genes Dev*. **10**: 2696–2706.
- Yam**, C. H., Ng, R. W., Siu, W. Y., Lau, A. W. and Poon, R. Y. (1999) Regulation of cyclin A-Cdk2 by SCF component Skp1 and F-box protein Skp2. *Mol. Cell. Biol*. **19**: 635–645.

- Yang, G.**, Rose M. S., Turgeon B. G., and Yoder O. C. (1996) A polyketide synthase is required for fungal virulence and production of the polyketide T-toxin. *Plant Physiol.* **8.11**: 2139–2150.
- Yate, G.** and Sadanandom, A. (2013) Ubiquitination in plant nutrient utilization. *Front Plant Sci.* **4**: 452.
- Yu, H.** (2007) Cdc20: a WD40 activator for a cell cycle degradation machine. *Molec. Cell.* **27**: 3–16.
- Zeng, L. R.**, Vega-Sánchez, M. E., Zhu, T. and Wang, G. L. (2006) Ubiquitination-mediated protein degradation and modification: an emerging theme in plant-microbe interactions. *Cell Research.* **16**: 413–426.
- Zhang, H.**, Kobayashi, R., Galaktionov, K. and Beach, D. (1995) p19Skp1 and p45Skp2 are essential elements of the cyclin A-CDK2 S phase kinase. *Cell.* **82**: 915–925.
- Zhao, X.**, Kim, Y., Park, G. and Xu, J. (2005) A Mitogen-Activated Protein Kinase Cascade Regulating Infection-Related Morphogenesis in *Magnaporthe grisea*. *Plant Cell.* **17**: 1317–1329.
- Zhao, X.**, Mehrabi, R. and Xu, J.R. (2007) Mitogen-activated protein kinase pathways and fungal pathogenesis. *Eukary. Cell.* **6**: 1701–1714.

- Zhao**, X. and Xu, J. R. (2007) A highly conserved MAPK-docking site in Mst7 is essential for Pmk1 activation in *Magnaporthe grisea*. *Mol. Microbiol.* **63**: 881–894.
- Zwiers**, L. H. and de Waard M. A. (2001) *Agrobacterium tumefaciens*-mediated gene disruption in the phytopathogen *Mycosphaerella graminicola*. *Curr. Genet.* **39.5-6**: 388–393.