

LIST OF FIGURES

S. No.	Title	Pg. No.
1	Structures of α -, β -, γ - and δ -HCH isomers	7
2	Various transportation and transformation processes involved in distribution of a pollutant in the environment	11
3	Anaerobic and aerobic degradation pathway of γ -HCH	27
4	Map showing soil sampling locations, Vadodara and Junagadh districts in Gujarat state, India.	53
5	Flow chart of procedure followed for the enrichment and isolation of γ -HCH degrader	54
6	Morphological characteristics and γ -HCH-clearance assay for isolate CGR-L1.	64
7	PCR amplification of partial 16S rDNA sequence of the bacterial isolates from the soil the soil samples	65
8	Phylogenetic relatedness of <i>Shewanella</i> sp. CGR-L1 based on 16S rDNA sequence	66
9	γ -HCH-clearance assay for isolate CGR-L2	69
10	Bacterial isolates obtained from γ -HCH degrading enrichment culture in MSM-L2 (presence γ -HCH) of soil Sample no. 4	70
11	Abundant of each bacterium in enrichment culture of soil sample no. 4	71
12	Phylogenetic tree of the bacterial partial 16S rDNA sequences of the cultivable isolates obtained from the γ -HCH degrading enrichment culture of the soil sample no. 4	73
13	Phylogenetic analysis of the isolate strain CGR-L2. The position of strain CGR-L2 is shown with respect to other species of the genus <i>Sphingobium</i>	74
14	Colorimetric determination for γ -HCH dechlorination using cell-free extracts of <i>Shewanella</i> sp. strain CGR-L1	89

15	Growth kinetics and biodegradation of γ -HCH by isolate <i>Shewanella</i> sp. CGR-L1	90
16	GC-MS analysis of γ -HCH degradation by <i>Shewanella</i> sp. strain CGR-L1 (analyzed from the clearance zone around culture of agar plate).	93
17	GC-MS analysis for the identification of metabolites produced during γ -HCH degradation using cell free extract of <i>Shewanella</i> sp. CGR-L2	95
18	PCR amplification of <i>linA</i> from isolate <i>Shewanella</i> sp. CGR-L1 and its phylogenetic analysis	97
19	Analysis of role of plasmid in γ -HCH biodegradation by <i>Shewanella</i> sp. CGR-L1	100
20	Southern blot hybridization of genomic DNA extracted from <i>Shewanella</i> sp. strain CGR-L1 and hybridised with non-radioactively labeled <i>linA</i> gene probe	101
21	Growth of <i>Shewanella</i> sp. CGR-L1 in MSM-L1 (supplemented with glucose) in presence and absence of different concentration manner of γ -HCH	102
22	SDS-PAGE analysis and colorimetric assay for γ -HCH dechlorination using cell-free extracts of strain CGR-L2	115
23	Growth kinetics and γ -HCH biodegradation analysis for <i>Sphingobium</i> sp. strain CGR-L2	118
24	Effect of initial inoculum on biodegradation of γ -HCH by isolate <i>Sphingobium</i> sp. CGR-L2	120
25	Effect of carbon source on growth of <i>Sphingobium</i> sp. strain CGR-L2	123
26	Effect of glucose on biodegradation of γ -HCH by isolate <i>Sphingobium</i> sp. CGR-L2	124
27	Effect of pH and temperature on biodegradation of γ -HCH by <i>Sphingobium</i> sp. strain CGR-L2	126
28	GC-analysis of γ -HCH biodegradation by <i>Sphingobium</i> sp. CGR-L2 in optimized parameters	128

29	Analysis of growth and γ -HCH residual concentration for <i>Sphingobium</i> sp. CGR-L2 under optimized parameters	129
30	GC-MS analysis of the metabolites producing during biodegradation of γ -HCH by <i>Sphingobium</i> sp. CGR-L2	130
31	GC analysis of biodegradation of technical HCH by isolate <i>Sphingobium</i> sp. CGR-L2	132
32	Reduction in residual concentration of HCH isomers in a time-dependent manner in MSM-L2, in presence and absence of <i>Sphingobium</i> sp. CGR-L2	133
33	Growth kinetics of <i>Sphingobium</i> sp. CGR-L2 in different concentration of γ -HCH, from 0 to 1000mg l ⁻¹	135
34	Biodegradation of γ -HCH by isolate <i>Sphingobium</i> sp. CGR-L2 at different initial concentration	136
35	PCR amplification of <i>lin</i> genes from <i>Sphingobium</i> sp. strain CGR-L2 genomic DNA using gene specific primers	138
36	Southern blot hybridization of genomic DNA of <i>Sphingobium</i> sp. strain CGR-L2 probed with nonradioactively labeled <i>linA</i> or <i>linB</i> or <i>linC</i>	141
37	Biodegradation of Hydroquinone by isolate CGR-L2 in MSM-L2	143
38	Microcosm (bioremediation) study by the isolate CGR-L2	144
39	Analysis of technical HCH biodegradation by <i>Sphingobium</i> sp. CGR-L2 during microcosm study	145
40	Amplification of <i>linA</i> (A), <i>linB</i> (B) and <i>linC</i> (C) from bacterial isolate of enrichment culture.	148