List of Figures

Fig No.	Figure Titles	Page No.
Fig. 1.1	Rhizodeposition root zones in the rhizosphere	5
Fig. 1.2	Positive and negative interactions among rhizobacteria and plants	7
Fig. 1.3	Cross-talk in the rhizosphere and step wise the molecular interactions involved	8
	during the establishment of PGPR strains in the rhizosphere	
Fig. 1.4	Model for suppression of root pathogens by siderophores of fluorescent Pseudomonas	13
Fig. 1.5	Chemical structures of antifungal metabolites produced by fluorescent	
	pseudomonads	18
Fig. 1.6	Overview of interactions between biocontrol strains, plants, pathogens,	
	predators, co-operators and soil.	21
Fig. 2.1	General strategy used for the isolation of biocontrol strains.	36
Fig. 2.2	Phylogenetic tree of Pseudomonas spp	37
Fig. 2.3	Location of sample collection sites on Gujarat map.	39
Fig. 2.4	Enrichment technique for the isolation of efficient root colonizing fluorescent	
	pseudomonads	42
Fig. 2.5	Schematic representation of the steps in PCR amplification of ITS.	47
Fig. 2.6	Representative plants from the enrichment assay.	56
Fig. 2.7	Representative strains of isolated fluorescent pseudomonads on Kings B plates.	56
Fig. 2.8	Antifungal activity against <i>R. bataticola</i> by representative fluorescent isolates.	57
Fig. 2.9	Radar plot for the inhibition of Rhizoctonia bataticola by fluorescent bacterial	
	isolates.	57
Fig. 2.10	Representative diagram of P-solubilization by isolates.	60
Fig. 2.11	Representative fluorescent isolates showing siderophore production on CAS	
	plates.	62
Fig. 2.12	Siderophore production by fluorescent bacterial isolates.	62

Fig. 2.13	Genomic DNA of representative fluorescent bacterial isolates.	64
Fig. 2.14	Amplification of fluorescent pseudomonads specific ITS (560bp) in the	
	bacterial isolates.	65
Fig. 2.15	IAA production by fluorescent <i>Pseudomonas</i> isolates.	67
Fig. 2.16	Bioassay using R. bataticola and S. aureus by fluorescent Pseudomonas	
	isolates.	70
Fig. 2.17	ACC deaminase activity of representative fluorescent Pseudomonas	
	Strains	72
Fig. 2.18	Precipitate formation method for the KMBA pathway in fluorescent	
	pseudomonads	73
Fig. 2.19	HCN production by fluorescent <i>Pseudomonas</i> strains.	75
Fig. 2.20	ITS-RFLP patterns of fluorescent Pseudomonas strains using Hae III.	76
Fig. 2.21	Dendrogram of fluorescent pseudomonad strains based on ITS-RFLP.	76
Fig. 3.1	Genetic organization of 2, 4-DAPG, Pyoluteorin, HCN, Phenazine and	
	Pyrrolnitrin biosynthesis genes.	85
Fig. 3.2	Biosynthetic pathway and genes involved for the biosynthesis of phenazines.	88
Fig. 3.3	Biosynthesis of PRN and PHZ from shikimic acid in strain PA23-63.	89
Fig. 3.4	Biotic and abiotic factors influencing DAPG production.	92
Fig. 3.5	General characteristics of the Gac/Rsm signal transduction pathway in g-	
	proteobacteria.	96
Fig. 3.6	Global regulatory pathway for the 2, 4-DAPG, Phz and HCN biosynthesis	
	regulation.	98
Fig. 3.7	Model for the regulatory cascade governing PCN production in P. chlororaphis	
	PCL1391.	99
Fig. 3.8	PCR amplification of <i>phlD</i> gene in fluorescent <i>Pseudomonas</i> strains.	113
Fig. 3.9	PCR amplification of hcnBC gene from fluorescent Pseudomonas strains.	114
Fig. 3.10	PCR amplification of pltC gene from fluorescent Pseudomonas strain	114
Fig. 3.11	PCR amplification of prnC gene from fluorescent Pseudomonas strains	115

Fig. 3.12	Pie graph for the number of ITS, <i>phlD</i> , <i>pltC</i> , <i>hcnBC</i> and <i>prnC</i> positive strains.	117
Fig. 3.13	Distribution of antibiotic synthesis genes among fluorescent Pseudomonas	
	strains.	117
Fig. 3.14	HPLC profile of pure DAPG.	118
Fig. 3.15	HPLC profile of 2,4-DAPG extracted from PfCHA0	119
Fig. 3.16:	HPLC profile for 2,4-DAPG production by fluorescent <i>Pseudomonas</i> isolates.	119
Fig. 3.17	2,4-DAPG production by fluorescent pseudomonad under different carbon	
	sources.	122
Fig. 3.18	Relatedness of fluorescent Pseudomonas isolates based on 2, 4- DAPG	
	production under different C-sources.	123
Fig. 3.19	Influence of phosphate levels on DAPG production in fluorescent	
	Pseudomonas.	124
Fig. 3.20	TLC for the separation and detection of PRN spot in the extract.	125
Fig. 3.21	Mechanism of MnO ₂ reduction.	128
Fig. 3.22	MnO ₂ reduction by representative fluorescent <i>Pseudomonas</i> isolates	129
Fig. 3.23	Clearance observed due to reduction of MnO ₂ precipitates in the presence of	
	phenazine.	130
Fig. 3.24	Quantification of phenazine production by fluorescent <i>Pseudomonas</i> isolates.	132
Fig. 3.25	Inhibition of R.bataticola by fluorescent Pseudomonas strains on MBV1-agar	
	plates.	133
Fig. 3.26	Inhibition of R.bataticola by PCN extracts of Isolate G44 grown on different	
	nitrogen sources.	134
Fig. 3.27	Fungal growth inhibition by G44 under 8mM and 16mM casamino acids	
	supplemented in MVB1-glucose-CAS.	135
Fig. 3.28	Growth Inhibition of R.bataticola by PCN extracts of Isolate G44 grown on	
	different C sources.	137
Fig. 3.29	Inhibition by PCN extract of MVB1 supplemented with NH4Cl and glycerol.	137
Fig. 3.30	UV absorption maxima for PCN extracts of Isolate G44 on different carbon	

	and nitrogen sources.	138
Fig. 3.31	Inhibition of <i>R. bataticola</i> by pure antibiotic std., ethyl acetate extracts of <i>Pf</i>	
	CHA0 grown in various nutritional combinations on 2d and 5d	143
Fig. 3.32	Effect of nutritional factors and their combinations on Pf CHA0 antifungal	144
	activity at 2d and 5d	
Fig. 3.33	Contribution of nutritional factors to PfCHA0 antifungal activity at 2d and 5d.	144
Fig. 3.34	Effect of nutritional factors and their combinations on PfCHA0 growth (OD	145
	600nm) at 2d and 5d.	
Fig. 3.35	Contribution of nutritional factors to OD600nm at 2d and 5d.	145
Fig. 3.36	Effect of nutritional factors and their combinations on PRN production at 2d	149
	and 5d.	
Fig. 3.37	Contribution of factors to PRN production at 2d and 5d	149
Fig. 3.38	Effect of nutritional factors and their combinations on PLT production at 2d	
	and 5d.	150
Fig. 3.39	Contribution of factors to the effect of factors on PLT production at 2d and 5d	150
Fig. 3.39 Fig. 3.40	Contribution of factors to the effect of factors on PLT production at 2d and 5d Effect of nutritional factors and their combinations on DAPG production at 2d	150
O	•	150 151
O	Effect of nutritional factors and their combinations on DAPG production at 2d	
Fig. 3.40	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d	151
Fig. 3.40 Fig. 3.41	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d	151 151
Fig. 3.40 Fig. 3.41 Fig. 3.42	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity	151 151 153
Fig. 3.40 Fig. 3.41 Fig. 3.42	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity Principal Component Analysis (PCA) for the effect of nutritional factors on	151 151 153
Fig. 3.40 Fig. 3.41 Fig. 3.42 Fig. 3.43	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity Principal Component Analysis (PCA) for the effect of nutritional factors on antibiotic production and antifungal activity of <i>Pf</i> CHAO.	151 151 153 154
Fig. 3.40 Fig. 3.41 Fig. 3.42 Fig. 3.43 Fig. 3.44	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity Principal Component Analysis (PCA) for the effect of nutritional factors on antibiotic production and antifungal activity of <i>Pf</i> CHAO. Antifungal activity by G20 extract under different nutrient compositions	151 151 153 154
Fig. 3.40 Fig. 3.41 Fig. 3.42 Fig. 3.43 Fig. 3.44	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity Principal Component Analysis (PCA) for the effect of nutritional factors on antibiotic production and antifungal activity of <i>Pf</i> CHAO. Antifungal activity by G20 extract under different nutrient compositions Radar graph for antifungal activity by G20 at 2 d and 5 d under different	151 151 153 154
Fig. 3.40 Fig. 3.41 Fig. 3.42 Fig. 3.43 Fig. 3.44 Fig. 3.45	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity Principal Component Analysis (PCA) for the effect of nutritional factors on antibiotic production and antifungal activity of <i>Pf</i> CHAO. Antifungal activity by G20 extract under different nutrient compositions Radar graph for antifungal activity by G20 at 2 d and 5 d under different nutrient combinations.	151 151 153 154 156
Fig. 3.40 Fig. 3.41 Fig. 3.42 Fig. 3.43 Fig. 3.44 Fig. 3.45	Effect of nutritional factors and their combinations on DAPG production at 2d and 5d Contribution of factors to DAPG production at 2d and 5d Contributions of PRN, PHL and PLT to the 2 d and 5 d antifungal activity Principal Component Analysis (PCA) for the effect of nutritional factors on antibiotic production and antifungal activity of <i>Pf</i> CHAO. Antifungal activity by G20 extract under different nutrient compositions Radar graph for antifungal activity by G20 at 2 d and 5 d under different nutrient combinations. Evolutionary distance between standard rhizobial strains used in this study	151 151 153 154 156

Fig. 4.4	Dual culture interaction of model fluorescent Pseudomonas and rhizobia	188
Fig. 4.5	Inhibition of rhizobial strains by the ethyl acetate extracts of fluorescent	
	Pseudomonas strains	190
Fig. 4.6	Clustering of fluorescent Pseudomonas strains based on their inhibitory effect	
	to rhizobial strains.	191
Fig. 4.7	Clustering of rhizobial strains based on their inhibition pattern by fluorescent	
	Pseudomonas strains.	191
Fig. 4.8	Inhibition of rhizobium spp by antibiotic extraction from different DAPG mediums by Pf CHA0 and G20.	192
Fig. 4.9	Inhibition of rhizobia by Pf CHA0 antibiotic extracts from culture supernatants from the biocontrol supportive media (A&B)	194
Fig. 4.10	Inhibition of rhizobia by antibiotic extracts of fluorescent <i>Pseudomonas</i> strains	
8	from cultures grown under biocontrol supportive nutrient combinations	194
Fig. 4.11	Interaction of fluorescent <i>Pseudomonas</i> and rhizobia in presence of	
	R.bataticola.	196
Fig 5. 1	Plant growth promotion effect of fluorescent pseudomonads on Vigna radiata	
	under synthetic MS medium.	211
Fig. 5.2	Effect of fluorescent pseudomonads on Vigna radiata growth	211
Fig. 5.3	Effect of fluorescent pseudomonads on root /shoot dry weight of Vigna radiata	212
Fig. 5.4	Seed germination assay for protection against fungus R. bataticola.by	
	fluorescent pseudomonad strains	213
Fig. 5.5	Plant assay for the protection of Vigna radiata by fluorescent pseudomonads	
	against R.bataticola.	213
Fig. 5.6	Pot study for the effect of ST1, Pf CHA0 and E.asburiae PSI3 on Cajanaus	
	cajan.	216
Fig. 5.7	Plant growth promotion effect of ST1, Pf CHA0 and E.asburiae PSI3 on	
	Cajanaus cajan	216
Fig. 5.8	Pot study for the effect of ST1, Pf CHA0 and E.asburiae PSI3 on Cajanaus	

	cajan.	217
Fig. 5.9	Plant growth promotion effect of IC3109, Pf CHA0 and E.asburiae PSI3 on	
	Cajanaus cajan.	217
Fig. 5.10	Pot study for the effect of IC3123, Pf CHA0 and E.asburiae PSI3 on Cajanaus	
	cajan.	218
Fig. 5.11	Plant growth promotion effect of IC3123, Pf CHA0, E.asburiae PSI3 on	218
	Cajanus cajan.	
Fig. 5.12	Histogram for the effect of PGP strains on root length.	219
Fig. 5.13	Box plot for the effect of different combination of PGP strains on root length.	219
Fig. 5.14	Histogram for the effect of PGP strains on shoot length.	219
Fig. 5.15	Box plot for the effect of different combination of PGP strains on shoot length.	219
Fig. 5.16	Effect of fluorescent pseudomonad strains under biocontrol supporting	
	nutritional combinations	221