## **List of Tables**

Table 1.1:	Microbial abundance reported in different subsurface environments8
Table 1.2:	Microbial diversity and distribution in different subsurface sedimentary
	ecosystems11
Table 1.3: I	Methods used for the measuring microbial activity of soil or sediment samples21
Table 1.4:	Phenotypic and genotypic methods used for functional microbial diversity
;	assessment
Table 1.5: H	Heavy metals bioremediation studies in the laboratory (i.e., microcosm study)28
Table 2.1: S	sediment enzymes assay protocols for measuring microbial activities41
Table 2.2: P	Primers and thermal profiles used in qPCR46
Table 3.1:	Genus level identification of bacterial isolates obtained from Chokari (CRD)
:	sediment profile67
Table 3.2: \$	Sequence analysis of selected DGGE bands obtained from CRD core samples
1	based on topmost best match sequence acquired in NCBI-BLASTn similarity
;	search70
Table 3.3:	Mean, SD, and Max values of enzyme activities and 16S rRNA gene copy
1	numbers of CRD core samples selected for metabarcoding and 0.2-0.3 m core
:	sample (CRD 1)
Table 3.4: T	Cotal number of reads and OTUs obtained after preprocessing71
Table 5.1:	Genus level identification of bacterial isolates obtained from Rayka (RYD)
:	sediment profile
Table 5.2: 0	Genus level identification of bacterial isolates obtained from Rampura (RMD)
:	sediment profile
Table 5.3:	MIC of bacterial strains obtained from RYD and RMD cores for Cr(VI).
	Bacterial strains showing MIC >40 ppm are represented in bold112
Table 5.4:	pH and EC of leachate fraction-1 obtained from control and experimental
	columns treated with Bacillus sp. RYD 15.1A and Streptomyces sp. RMD
•	42.2B116