

Bibliography

Bibliography

- Agarwal, R. M., Tomar, N. S., Jatav, K. S., & Sharma, G. L. (2009) Potassium induced changes in flowering plants. In: *Flower Retrospect and Prospect* (Professor Vishwambhar Puri Birth Centenary Volume). SR Scientific Publication, Delhi, pp. 158–186.
- Arazoe, T., Miyoshi, K., Yamato, T., Ogawa, T., Ohsato, S., Arie, T., & Kuwata, S. (2015). Tailor-made CRISPR/Cas system for highly efficient targeted gene replacement in the rice blast fungus. *Biotechnology and bioengineering*, 112(12), 2543–2549.
- Bacete, L., Mélida, H., Miedes, E., & Molina, A. (2018). Plant cell wall-mediated immunity: cell wall changes trigger disease resistance responses. *The Plant journal: for cell and molecular biology*, 93(4), 614–636.
- Balcerzak, M., Harris, L. J., Subramaniam, R., Ouellet, T. (2012). The feruloyl esterase gene family of *Fusarium graminearum* is differentially regulated by aromatic compounds and hosts. *Fungal Biology*, 116, 478–488.
- Battaglia, E., Klaubauf, S., Vallet, J., Ribot, C., Lebrun, M. H., & de Vries, R. P. (2013). Xlr1 is involved in the transcriptional control of the pentose catabolic pathway, but not hemi-cellulolytic enzymes in *Magnaporthe oryzae*. *Fungal genetics and biology: FG & B*, 57, 76–84.
- Beliën, T., Van Campenhout, S., Robben, J., & Volckaert, G. (2006). Microbial endoxylanases: effective weapons to breach the plant cell-wall barrier or, rather, triggers of plant defense systems? *Molecular plant-microbe interactions: MPMI*, 19(10), 1072–1081.
- Benoit, I., Navarro, D., Marnet, N., Rakotomanomana, N., Lesage-Meessen, L., Sigoillot, J. C., Asther, M., & Asther, M. (2006). Feruloyl esterases as a tool for the release of phenolic compounds from agro-industrial by-products. *Carbohydrate Research*, 341, 1820–1827.
- Biswas, S., Dijck, P.V., & Datta, A. (2007). Environmental sensing and signal transduction pathways regulating morphopathogenic Determinants of *Candida albicans*. *Microbiology and molecular biology reviews*, 71, 348–376.
- Black, R. L. B., & Dix, N. J. (1976). Utilization of ferulic acid by microfungi from litter and soil. *Transactions of the British mycological society*, 66, 313–317.
- Boutrot, F., & Zipfel, C. (2017). Function, Discovery, and Exploitation of Plant Pattern Recognition Receptors for Broad-Spectrum Disease Resistance. *Annual review of phytopathology*, 55, 257–286.

- Broekaert, W. F., Cammue, B. P., De Bolle, M. F., Thevissen, K., De Samblanx, G. W., Osborn, R. W., et al. (1997). Antimicrobial Peptides from Plants. *Critical reviews in plant sciences*, 16(3), 297–323.
- Broekaert, W. F., Terras, F. R., Cammue, B. P., & Osborn, R. W. (1995). Plant defensins: novel antimicrobial peptides as components of the host defense system. *Plant physiology*, 108(4), 1353–1358.
- Brunner, K., Lichtenauer, A. M., Kratochwill, K., Delic, M., & Mach, R. L. (2007). Xyr1 regulates xylanase but not cellulase formation in the head blight fungus *Fusarium graminearum*. *Current genetics*, 52(5-6), 213–220.
- Bunzel, M., Ralph, J., Marita, J. M., Hatfield, R. D., Steinhart, H., (2001). Diferulates as structural components in soluble and insoluble cereal dietary fibre. *J. Sci. Fd. Agric.* 81: 653–660.
- Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25, 1972–1973.
- Chanclud, E., Kisiala, A., Emery, N. R., Chalvon, V., Ducasse, A., Romiti-Michel, C., Gravot, A., Kroj, T., & Morel, J. B. (2016). Cytokinin Production by the Rice Blast Fungus Is a Pivotal Requirement for Full Virulence. *PLoS pathogens*, 12(2), e1005457.
- Cheng, Y. T., Zhang, L., & He, S. Y. (2019). Plant-Microbe Interactions Facing Environmental Challenge. *Cell host & microbe*, 26(2), 183–192.
- Cheval, C., & Faulkner, C. (2018). Plasmodesmal regulation during plant-pathogen interactions. *The New phytologist*, 217(1), 62–67.
- Cosgrove D. J. (2005). Growth of the plant cell wall. *Nature reviews. Molecular cell biology*, 6(11), 850–861.
- Cosgrove, D. J. (2001). Wall structure and wall loosening. A look backwards and forwards. *Plant Physiology*, 125, 131–134.
- Couch, B. C., & Kohn, L. M. (2002). A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia*, 94(4), 683–693.
- Crepin, V. F., Faulds, C. B., & Connerton, I. F. (2003). A non-modular type-B feruloyl esterase from *Neurospora crassa* exhibits concentration dependent substrate inhibition. *Biochemical Journal*, 370, 417–427.
- Crepin, V. F., Faulds, C. B., & Connerton, I. F. (2004). Functional recognition of new classes of feruloyl esterases. *Applied Microbiology & Biotechnology*, 63, 647–652.

- Dangl, J. L., Horvath, D. M., & Staskawicz, B. J. (2013). Pivoting the plant immune system from dissection to deployment. *Science (New York, N.Y.)*, 341(6147), 746–751.
- de Jong, J., McCormack, B., Smirnov, N., Talbot, N. (1997). Glycerol generates turgor in rice blast. *Nature*, 389, 244–245.
- De Lorenzo, G., D'Ovidio, R., & Cervone, F. (2001). The role of polygalacturonase-inhibiting proteins (PGIPs) in defense against pathogenic fungi. *Annu. Rev. Phytopathol.*, 39, 313–35.
- De Vries, R. P., Michelsen, B., Poulsen, C. H., Kroon, P. A., van den Heuvel, R. H. H., et al. (1997). The faeA genes from *Aspergillus niger* and *Aspergillus tubingensis* encode ferulic acid esterases involved in degradation of complex cell wall polysaccharides. *Applied environmental microbiology*, 63, 4638–4644.
- de Vries, R. P., vanKuyk, P. A., Kester, H. C., & Visser, J. (2002). The *Aspergillus niger* faeB gene encodes a second feruloyl esterase involved in pectin and xylan degradation and is specifically induced in the presence of aromatic compounds. *The Biochemical journal*, 363(Pt 2), 377–386.
- Dean, R. A., Talbot, N. J., Farman, M. L., Mitchell, T. K., Orbach, M. J., et al. (2005). The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature*, 434, 980–986.
- Dellaporta, S. L., Wood, J., & Hicks, J. B. (1983). A plant DNA miniprep: Version II. *Plant Molecular Biology Reports*, 1, 19–21.
- Dilokpimol, A., Makela, M. R., Anguilar-Pontes, M. V., Benoit-Gelber, I., Hidden, K. S., et al. (2016). Diversity of fungal feruloyl esterases: updated phylogenetic classification, properties, and industrial applications. *Biotechnology & Biofuels*, 9, 231.
- Durrant, W. E., & Dong, X. (2004). Systemic acquired resistance. *Annual review of phytopathology*, 42, 185–209.
- Ebbole, D. J. (2007). *Magnaporthe* as a model for understanding host-pathogen interactions. *Annual review of phytopathology*, 45, 437–456.
- Eddy, S. R. (2011). Accelerated profile HMM searches. *PLoS computational biology*, 7, e1002195.
- Egan, M. J., Wang, Z. Y., Jones, M. A., Smirnov, N., & Talbot, N. J. (2007). Generation of reactive oxygen species by fungal NADPH oxidases is required for rice blast disease. *Proceedings of the National Academy of Sciences of the United States of America*, 104(28), 11772–11777.

- Faulds, C. B. & Williamson, G. (1995). Release of ferulic acid from wheat bran by a ferulic acid esterase (FAE-III) from *Aspergillus niger*. *Applied microbiology and biotechnology*, 43, 1082–1087.
- Faulds, C. B., & Williamson, G. (1994). Purification and characterization of a ferulic acid esterase (FAEIII) from *Aspergillus niger*: specificity for the phenolic moiety and binding to micro-crystalline cellulose. *Microbiology*, 140, 779–787.
- Faulkner, C., Petutschnig, E., Benitez-Alfonso, Y., Beck, M., Robatzek, S., Lipka, V., & Maule, A. J. (2013). LYM2-dependent chitin perception limits molecular flux via plasmodesmata. *Proceedings of the National Academy of Sciences of the United States of America*, 110(22), 9166–9170.
- Fernandez, J., & Orth, K. (2018). Rise of a Cereal Killer: The Biology of *Magnaporthe oryzae* Biotrophic Growth. *Trends in microbiology*, 26(7), 582–597.
- Fernandez, J., & Wilson, R. A. (2014). Cells in cells: morphogenetic and metabolic strategies conditioning rice infection by the blast fungus *Magnaporthe oryzae*. *Protoplasma*, 251, 37–47.
- Ferrari, S., Sella, L., Janni, M., De Lorenzo, G., Favaron, F., & D'Ovidio, R. (2012). Transgenic expression of polygalacturonase-inhibiting proteins in Arabidopsis and wheat increases resistance to the flower pathogen *Fusarium graminearum*. *Plant biology (Stuttgart, Germany)*, 14 Suppl 1, 31–38.
- Freeman, B.C. and Beattie, G.A. (2008). An Overview of Plant Defenses against Pathogens and Herbivores. *The Plant Health Instructor*.
- Galhano, R., & Talbot, N. J. (2011). The biology of blast: Understanding how *Magnaporthe oryzae* invades rice plants. *Fungal biology reviews*, 25(1), 61–67.
- Gietz, R. D., & Woods, R. A. (2002). Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods in enzymology*, 350, 87–96.
- Giraldo, M. C., & Valent, B. (2013). Filamentous plant pathogen effectors in action. *Nature reviews. Microbiology*, 11(11), 800–814.
- Giraldo, M. C., Dagdas, Y. F., Gupta, Y. K., Mentlak, T. A., Yi, M., Martinez-Rocha, A. L., Saitoh, H., Terauchi, R., Talbot, N. J., & Valent, B. (2013). Two distinct secretion systems facilitate tissue invasion by the rice blast fungus *Magnaporthe oryzae*. *Nature communications*, 4, 1996.
- Glazebrook J. (2005). Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annual review of phytopathology*, 43, 205–227.

- Ham, J. H., Kim, M. G., Lee, S. Y., & Mackey, D. (2007). Layered basal defenses underlie non-host resistance of *Arabidopsis* to *Pseudomonas syringae* pv. *phaseolicola*. *The Plant journal: for cell and molecular biology*, 51(4), 604–616.
- Hamel, L. P., Nicole, M. C., Duplessis, S., & Ellis, B. E. (2012). Mitogen-activated protein kinase signaling in plant-interacting fungi: distinct messages from conserved messengers. *The Plant cell*, 24(4), 1327–1351.
- Hamel, L. P., Nicole, M. C., Duplessis, S., & Ellis, B. E. (2012). Mitogen-activated protein kinase signaling in plant-interacting fungi: distinct messages from conserved messengers. *The Plant cell*, 24(4), 1327–1351.
- Harris, P. J., & Hartley, R. D. (1977). Detection of bound ferulic acid in cell walls of the Gramineae by ultraviolet fluorescence microscopy. *Nature*, 259, 508–510.
- Hegde, S., & Muralikrishna, G. (2009). Isolation and partial characterization of alkaline feruloyl esterases from *Aspergillus niger* CFR 1105 grown on wheat bran. *World Journal of Microbiology and Biotechnology*, 25, 1963–1969.
- Hermoso, J. A., Aparicio, J. S., Molina, R., Juge, N., Gonzalez, R., & Faulds, C. B. (2004). The crystal structure of Feruloyl esterase A from *Aspergillus niger* suggests evolutive functional convergence in feruloyl esterase family. *Journal of molecular biology*, 338(3), 495–506.
- Howard, R. J., & Valent, B. (1996). Breaking and entering: host penetration by the fungal rice blast pathogen *Magnaporthe grisea*. *Annual review of microbiology*, 50, 491–512.
- Howard, R. J., Ferrari, M. A., Roach, D. H., & Money, N. P. (1991). Penetration of hard substrates by a fungus employing enormous turgor pressures. *Proceedings of the National Academy of Sciences of the United States of America*, 88(24), 11281–11284.
- Igarashi, S., Utimada, C., Igarashi, L., Kazuma, A., & Lopes, R. (1986). *Pyricularia* sp. em trigo. I. Ocorrência de *Pyricularia* sp. no Estado do Paraná. *Fitopatologia Brasileira*, 11, 351–352.
- Islam, M. S., Haque, M. S., Islam, M. M., Emdad, E. M., Halim, A., Hossen, Q. M., et al. (2012). Tools to kill: genome of one of the most destructive plant pathogenic fungi *Macrophomina phaseolina*. *BMC genomics*, 13, 493.
- Islam, M. T., Croll, D., Gladieux, P., Soanes, D. M., Persoons, A., Bhattacharjee, P., et al. (2016). Emergence of wheat blast in Bangladesh was caused by a South American lineage of *Magnaporthe oryzae*. *BMC biology*, 14(1), 84.
- Jacobs, A. K., Lipka, V., Burton, R. A., Panstruga, R., Strizhov, N., Schulze-Lefert, P., & Fincher, G. B. (2003). An *Arabidopsis* Callose Synthase, GSL5, Is Required for Wound and Papillary Callose Formation. *The Plant cell*, 15(11), 2503–2513.

- Jeon, J., Lee, G. W., Kim, K. T., Park, S. Y., Kim, S., Kwon, S., et al. (2020). Transcriptome Profiling of the Rice Blast Fungus *Magnaporthe oryzae* and Its Host *Oryza sativa* During Infection. *Molecular plant-microbe interactions: MPMI*, 33(2), 141–144.
- Jones, J., & Dangl, J. (2006). The plant immune system. *Nature*, 444, 323–329.
- Juge N. (2006). Plant protein inhibitors of cell wall degrading enzymes. *Trends in plant science*, 11(7), 359–367.
- Kachroo, P., Lee, K. H., Schwerdel, C., Bailey, J. E., & Chattoo, B. B. (1997). Analysis of host-induced response in the rice blast fungus *Magnaporthe grisea* using two-dimensional polyacrylamide gel electrophoresis. *Electrophoresis*, 18(1), 163–169.
- Kachroo, P., Leong, S. A., & Chattoo, B. B. (1994). Pot2, an inverted repeat transposon from the rice blast fungus *Magnaporthe grisea*. *Molecular & general genetics: MGG*, 245(3), 339–348.
- Kankanala, P., Czymmek, K., & Valent, B. (2007). Roles for rice membrane dynamics and plasmodesmata during biotrophic invasion by the blast fungus. *The Plant cell*, 19(2), 706–724.
- Kihoro, J., Bosco, N. J., & Murage, H. (2013). Suitability analysis for rice growing sites using a multicriteria evaluation and GIS approach in great Mwea region, Kenya. *SpringerPlus*, 2(1), 265.
- Koseki, T., Hori, A., Seki, S., Murayama, T., & Shiono, Y. (2009). Characterization of two distinct feruloyl esterases, AoFaeB and AoFaeC, from *Aspergillus oryzae*. *Applied Microbiology and Biotechnology*, 83, 689–696.
- Kouzai, Y., Mochizuki, S., Nakajima, K., Desaki, Y., Hayafune, M., Miyazaki, H., Yokotani, N., Ozawa, K., Minami, E., Kaku, H., Shibuya, N., & Nishizawa, Y. (2014). Targeted gene disruption of *OsCERK1* reveals its indispensable role in chitin perception and involvement in the peptidoglycan response and immunity in rice. *Molecular plant-microbe interactions: MPMI*, 27(9), 975–982.
- Kroon, P. A., & Williamson, G. (1996). Release of ferulic acid from sugar-beet pulp by using arabinanase, arabinofuranosidase and an esterase from *Aspergillus niger*. *Biotechnology and applied biochemistry*, 23(3), 263–267.
- Kubicek, C. P., Starr, T. L., & Glass, N. L. (2014). Plant cell wall-degrading enzymes and their secretion in plant-pathogenic fungi. *Annual review of phytopathology*, 52, 427–451.
- Lee, S. J., Kelley, B. S., Damasceno, C. M., St John, B., Kim, B. S., Kim, B. D., & Rose, J. K. (2006). A functional screen to characterize the secretomes of eukaryotic pathogens

- and their hosts *in planta*. *Molecular plant-microbe interactions: MPMI*, 19(12), 1368–1377.
- Lee, S., Choi, H., Suh, S., Doo, I. S., Oh, K. Y., Choi, E. J., Schroeder Taylor, A. T., Low, P. S., & Lee, Y. (1999). Oligogalacturonic acid and chitosan reduce stomatal aperture by inducing the evolution of reactive oxygen species from guard cells of tomato and *Commelina communis*. *Plant physiology*, 121(1), 147–152.
- Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic acids research*, 49(W1), W293–W296.
- Lionetti, V., Raiola, A., Camardella, L., Giovane, A., Obel, N., Pauly, M., Favaron, F., Cervone, F., & Bellincampi, D. (2007). Overexpression of pectin methylesterase inhibitors in *Arabidopsis* restricts fungal infection by *Botrytis cinerea*. *Plant physiology*, 143(4), 1871–1880.
- Liu, W., Liu, J., Triplett, L., Leach, J. E., & Wang, G. L. (2014). Novel insights into rice innate immunity against bacterial and fungal pathogens. *Annual review of phytopathology*, 52, 213–241.
- Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta Ct}$ Method. *Methods (San Diego, Calif.)*, 25(4), 402–408.
- Lorrai, R., & Ferrari, S. (2021). Host Cell Wall Damage during Pathogen Infection: Mechanisms of Perception and Role in Plant-Pathogen Interactions. *Plants (Basel, Switzerland)*, 10(2), 399.
- Ma, H., Zhang, B., Gai, Y., Sun, X., Chung, K. R., & Li, H. (2019). Cell-Wall-Degrading Enzymes Required for Virulence in the Host Selective Toxin-Producing Necrotroph *Alternaria alternata* of Citrus. *Frontiers in microbiology*, 10, 2514.
- Malaker, P. K., Barma, N. C. D., Tiwari, T. P., Collis, W. J., Duveiller, E., Singh, P.K., et al. (2016). First Report of Wheat Blast Caused by *Magnaporthe oryzae* Pathotype *triticum* in Bangladesh. *Plant Disease*, 100:11, pp. 2330.
- Marchler-Bauer, A., Derbyshire, M. K., Gonzales, N. R., Lu, S., Chitsaz, F., Geer, L. Y., et al. (2015). CDD: NCBI's conserved domain database. *Nucleic acids research*, 43(Database issue), D222–D226.
- Marroquin-Guzman, M., Hartline, D., Wright, J. D., Elowsky, C., Bourret, T. J., & Wilson, R. A. (2017). The *Magnaporthe oryzae* nitrooxidative stress response suppresses rice innate immunity during blast disease. *Nature microbiology*, 2, 17054.
- Martínez-Soto, D., Robledo-Briones, A. M., Estrada-Luna, A. A., & Ruiz-Herrera, J. (2013). Transcriptomic analysis of *Ustilago maydis* infecting *Arabidopsis* reveals

- important aspects of the fungus pathogenic mechanisms. *Plant signaling & behavior*, 8(8), e25059.
- Mastihuba, V., Kremnický, L., Mastihubová, M., Willett, J. L., & Côté, G. L. (2002). A spectrophotometric assay for feruloyl esterases. *Analytical biochemistry*, 309(1), 96–101.
- Mathioni, S. M., Belo, A., Rizzo, C. J., Dean, R. A., & Donofrio, N. M. (2011). Transcriptome profiling of the rice blast fungus during invasive plant infection and *in vitro* stresses. *BMC Genomics*, 12, 49.
- Mentlak, T. A., Kombrink, A., Shinya, T., Ryder, L. S., Otomo, I., Saitoh, H., Terauchi, R., Nishizawa, Y., Shibuya, N., Thomma, B. P., & Talbot, N. J. (2012). Effector-mediated suppression of chitin-triggered immunity by *Magnaporthe oryzae* is necessary for rice blast disease. *The Plant cell*, 24(1), 322–335.
- Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular biology and evolution*, 37(5), 1530–1534.
- Mitchell, T. K., & Dean, R. A. (1995). The cAMP-dependent protein kinase catalytic subunit is required for appressorium formation and pathogenesis by the rice blast pathogen *Magnaporthe grisea*. *The Plant cell*, 7(11), 1869–1878.
- Moreira, L. R., Milanezi, N. G., & Filho, E. X. (2011). Enzymology of Plant Cell Wall Breakdown: An Update. In: Buckeridge, M. S. & Goldman, G. H. (eds.), *Routes to Cellulosic Ethanol*, Springer, pp.73–96.
- Mosquera, G., Giraldo, M. C., Khang, C. H., Coughlan, S., & Valent, B. (2009). Interaction transcriptome analysis identifies *Magnaporthe oryzae* BAS1-4 as Biotrophy-associated secreted proteins in rice blast disease. *The Plant cell*, 21(4), 1273–1290.
- Nguyen, Q. B., Itoh, K., Van Vu, B., Tosa, Y., & Nakayashiki, H. (2011). Simultaneous silencing of endo- β -1,4 xylanase genes reveals their roles in the virulence of *Magnaporthe oryzae*. *Molecular microbiology*, 81(4), 1008–1019.
- Olivares-Hernández, R., Sunner, H., Frisvad, J. C., Olsson, L., Nielsen, J., & Panagiotou, G. (2010). Combining substrate specificity analysis with support vector classifiers reveals feruloyl esterase as a phylogenetically informative protein group. *PloS one*, 5(9), e12781.
- Park, C. H., Chen, S., Shirsekar, G., Zhou, B., Khang, C. H., Songkumarn, P., Afzal, A. J., Ning, Y., Wang, R., Bellizzi, M., Valent, B., & Wang, G. L. (2012). The *Magnaporthe oryzae* effector AvrPiz-t targets the RING E3 ubiquitin ligase APIP6 to suppress

- pathogen-associated molecular pattern-triggered immunity in rice. *The Plant cell*, 24(11), 4748–4762.
- Patkar, R. N., & Naqvi, N. I. (2017). Fungal manipulation of hormone-regulated plant defense. *PLoS pathogens*, 13(6), e1006334.
- Patkar, R. N., Benke, P. I., Qu, Z., Chen, Y. Y., Yang, F., Swarup, S., & Naqvi, N. I. (2015). A fungal monooxygenase-derived jasmonate attenuates host innate immunity. *Nature chemical biology*, 11(9), 733–740.
- Patkar, R. N., Ramos-Pamplona, M., Gupta, A. P., Fan, Y., & Naqvi, N. I. (2012). Mitochondrial β -oxidation regulates organellar integrity and is necessary for conidial germination and invasive growth in *Magnaporthe oryzae*. *Molecular microbiology*, 86(6), 1345–1363.
- Patkar, R. N., Suresh, A., & Naqvi, N. I. (2010). MoTea4-mediated polarized growth is essential for proper asexual development and pathogenesis in *Magnaporthe oryzae*. *Eukaryotic cell*, 9(7), 1029–1038.
- Perez-Nadales, E., Nogueira, M. F., Baldin, C., Castanheira, S., El Ghalid, M., Grund, E., Lengeler, K., et al. (2014). Fungal model systems and the elucidation of pathogenicity determinants. *Fungal genetics and biology: FG & B*, 70(100), 42–67.
- Petersen, T. N., Brunak, S., von Heijne, G., & Nielsen, H. (2011). SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nature methods*, 8(10), 785–786.
- Pham, C. L., Rey, A., Lo, V., Soulès, M., Ren, Q., Meisl, G., Knowles, T. P., Kwan, A. H., & Sunde, M. (2016). Self-assembly of MPG1, a hydrophobin protein from the rice blast fungus that forms functional amyloid coatings, occurs by a surface-driven mechanism. *Scientific reports*, 6, 25288.
- Pham, K. T., Inoue, Y., Vu, B. V., Nguyen, H. H., Nakayashiki, T., Ikeda, K., & Nakayashiki, H. (2015). MoSET1 (Histone H3K4 Methyltransferase in *Magnaporthe oryzae*) Regulates Global Gene Expression during Infection-Related Morphogenesis. *PLoS genetics*, 11(7), e1005385.
- Pinto, C. (2005). Feruloyl Esterase: A Principal Biodegradative Enzyme. In: Borkar, S. (ed.), *Bioprospects of Coastal Eubacteria*, Springer, Switzerland, pp. 209–224.
- Prakash, C., Manjrekar, J., & Chattoo, B. B. (2016). Skp1, a component of E3 ubiquitin ligase, is necessary for growth, sporulation, development and pathogenicity in rice blast fungus (*Magnaporthe oryzae*). *Molecular plant pathology*, 17(6), 903–919.
- Puchart, V., Vršanská, M., Mastihubová, M., Topakas, E., Vafiadi, C., Faulds, C. B., et al. (2007). Substrate and positional specificity of feruloyl esterases for monoferuloylated

- and monoacetylated 4-nitrophenyl glycoside. *Journal of Biotechnology*, 127, 235–243.
- Quoc, N. B., & Chau, N. (2017). The Role of Cell Wall Degrading Enzymes in Pathogenesis of *Magnaporthe oryzae*. *Current protein & peptide science*, 18(10), 1019–1034.
- Sakulkoo, W., Osés-Ruiz, M., Oliveira Garcia, E., Soanes, D. M., Littlejohn, G. R., Hacker, C., Correia, A., Valent, B., & Talbot, N. J. (2018). A single fungal MAP kinase controls plant cell-to-cell invasion by the rice blast fungus. *Science (New York, N.Y.)*, 359(6382), 1399–1403.
- Sasaki, T., & Burr, B. (2000). International Rice Genome Sequencing Project: the effort to completely sequence the rice genome. *Current opinion in plant biology*, 3(2), 138–141.
- Seiboth, B., Herold, S., & Kubicek, C. P. (2012). Metabolic engineering of inducer formation for cellulase and hemicellulase gene expression in *Trichoderma reesei*. *Sub-cellular biochemistry*, 64, 367–390.
- Shi, Y., Mosser, D. D., & Morimoto, R. I. (1998). Molecular chaperones as HSF1-specific transcriptional repressors. *Genes & development*, 12(5), 654–666.
- Sievers, F., & Higgins, D. G. (2018). Clustal Omega for making accurate alignments of many protein sequences. *Protein science: a publication of the Protein Society*, 27(1), 135–145.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular systems biology*, 7, 539.
- Singh, R., Dangol, S., Chen, Y., Choi, J., Cho, Y. S., Lee, J. E., et al. (2016). *Magnaporthe oryzae* Effector AVR-Pii Helps to Establish Compatibility by Inhibition of the Rice NADP-Malic Enzyme Resulting in Disruption of Oxidative Burst and Host Innate Immunity. *Molecules and cells*, 39(5), 426–438.
- Skamnioti, P., & Gurr, S. J. (2007). *Magnaporthe grisea* cutinase2 mediates appressorium differentiation and host penetration and is required for full virulence. *The Plant cell*, 19(8), 2674–2689.
- Skamnioti, P., & Gurr, S. J. (2009). Against the grain: safeguarding rice from rice blast disease. *Trends in biotechnology*, 27(3), 141–150.
- Skamnioti, P., Henderson, C., Zhang, Z., Robinson, Z., & Gurr, S. J. (2007). A novel role for catalase B in the maintenance of fungal cell-wall integrity during host invasion in the rice blast fungus *Magnaporthe grisea*. *Molecular plant-microbe interactions: MPMI*, 20(5), 568–580.

- Soundararajan, S., Jedd, G., Li, X., Ramos-Pamplona, M., Chua, N. H., & Naqvi, N. I. (2004). Woronin body function in *Magnaporthe grisea* is essential for efficient pathogenesis and for survival during nitrogen starvation stress. *The Plant cell*, 16(6), 1564–1574.
- Spence, C. A., Lakshmanan, V., Donofrio, N., & Bais, H. P. (2015). Crucial Roles of Absciscic Acid Biogenesis in Virulence of Rice Blast Fungus *Magnaporthe oryzae*. *Frontiers in plant science*, 6, 1082.
- Talbot N. J. (2003). On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*. *Annual review of microbiology*, 57, 177–202.
- Talbot, N. J., Ebbole, D. J., & Hamer, J. E. (1993). Identification and characterization of MPG1, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *The Plant cell*, 5(11), 1575–1590.
- Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular biology and evolution*, 38(7), 3022–3027.
- Tanaka, E., Koga, H., Mori, M., & Mori, M. (2011). Auxin production by the rice blast fungus and its localization in host tissue. *Journal of Phytopathology*, 159(7-8), 522–530.
- Tanaka, S., Yamada, K., Yabumoto, K., Fujii, S., Huser, A., Tsuji, G., et al. (2007). *Saccharomyces cerevisiae* SSD1 orthologs are essential for host infection by the ascomycete plant pathogens *Colletotrichum lagenarium* and *Magnaporthe grisea*. *Molecular microbiology*, 64(5), 1332–1349.
- Tonukari, N. J., Scott-Craig, J. S., & Walton, J. D. (2000). The *Cochliobolus carbonum* SNF1 gene is required for cell wall-degrading enzyme expression and virulence on maize. *The Plant cell*, 12(2), 237–248.
- Torres, M. A., & Dangl, J. L. (2005). Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development. *Current opinion in plant biology*, 8(4), 397–403.
- Valent, B., & Khang, C. H. (2010). Recent advances in rice blast effector research. *Current opinion in plant biology*, 13(4), 434–441.
- Vincken, J. P., Schols, H. A., Oomen, R. J., McCann, M. C., Ulvskov, P., Voragen, A. G., & Visser, R. G. (2003). If homogalacturonan were a side chain of rhamnogalacturonan I. Implications for cell wall architecture. *Plant physiology*, 132(4), 1781–1789.
- Vu, B., Itoh, K., Nguyen, Q. B., Tosa, Y., & Nakayashiki, H. (2012). Cellulases belonging to glycoside hydrolase families 6 and 7 contribute to the virulence of *Magnaporthe oryzae*. *Molecular plant-microbe interactions: MPMI*, 25(9), 1135–1141.

- Walton J. D. (1994). Deconstructing the Cell Wall. *Plant physiology*, 104(4), 1113–1118.
- Wang, Z. Y., Soanes, D. M., Kershaw, M. J., & Talbot, N. J. (2007). Functional analysis of lipid metabolism in *Magnaporthe grisea* reveals a requirement for peroxisomal fatty acid beta-oxidation during appressorium-mediated plant infection. *Molecular plant-microbe interactions: MPMI*, 20(5), 475–491.
- Wegner, A., Casanova, F., Loehrer, M., Jordine, A., Bohnert, S., Liu, X., Zhang, Z., & Schaffrath, U. (2022). Gene deletion and constitutive expression of the pectate lyase gene 1 (*MoPL1*) lead to diminished virulence of *Magnaporthe oryzae*. *Journal of microbiology (Seoul, Korea)*, 60(1), 79–88.
- Wilson R. A. (2021). Microbe of the Month: *Magnaporthe oryzae*. *Trends in Microbiology*, 29(7), 663–664.
- Wilson, R. A., & Talbot, N. J. (2009). Under pressure: investigating the biology of plant infection by *Magnaporthe oryzae*. *Nature reviews. Microbiology*, 7(3), 185–195.
- Xu, M., Gao, X., Chen, J., Yin, Z., Feng, H., & Huang, L. (2018). The feruloyl esterase genes are required for full pathogenicity of the apple tree canker pathogen *Valsa mali*. *Molecular plant pathology*, 19(6), 1353–1363.
- Yang, C., Yu, Y., Huang, J., Meng, F., Pang, J., Zhao, Q., Islam, M. A., Xu, N., Tian, Y., & Liu, J. (2019). Binding of the *Magnaporthe oryzae* Chitinase MoChia1 by a Rice Tetratricopeptide Repeat Protein Allows Free Chitin to Trigger Immune Responses. *The Plant cell*, 31(1), 172–188.
- Yang, F., & Naqvi, N. I. (2014). Sulfonyleurea resistance reconstitution as a novel strategy for *ILV2*-specific integration in *Magnaporthe oryzae*. *Fungal genetics and biology: FG & B*, 68, 71–76.
- Yu, J. H., Hamari, Z., Han, K. H., Seo, J. A., Reyes-Domínguez, Y., & Scazzocchio, C. (2004). Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi. *Fungal genetics and biology: FG & B*, 41(11), 973–981.
- Zhang, B., Gao, Y., Zhang, L., & Zhou, Y. (2021). The plant cell wall: Biosynthesis, construction, and functions. *Journal of integrative plant biology*, 63, 251–272.
- Zhang, H., Zheng, X., & Zhang, Z. (2016). The *Magnaporthe grisea* species complex and plant pathogenesis. *Molecular plant pathology*, 17(6), 796–804.
- Zhang, N., Li, S., Xiong, L., & Hong, Y. (2015). Cellulose-hemicellulose interaction in wood secondary cell-wall. *Modelling and Simulation in Materials Science and Engineering*, 23, 085010.
- Zhao, Z., Liu, H., Wang, C., & Xu, J. R. (2013). Comparative analysis of fungal genomes reveals different plant cell wall degrading capacity in fungi. *BMC genomics*, 14, 274.

- Zheng, X., Zhou, J., Lin, C., Lin, X., Lan, L., Wang, Z., et al. (2009). Secretion property and gene expression pattern of a putative feruloyl esterase in *Magnaporthe grisea*. In: Xiaofan, W., Valent, B. (eds), *Advances in genetics, genomics and control of rice blast disease*, 1st edn. Springer, Netherlands, pp. 41–50.
- Zipfel, C., & Oldroyd, G. E. (2017). Plant signalling in symbiosis and immunity. *Nature*, 543(7645), 328–336.
- Ziv, C., Zhao, Z., Gao, Y. G., & Xia, Y. (2018). Multifunctional Roles of Plant Cuticle During Plant-Pathogen Interactions. *Frontiers in plant science*, 9, 1088.