

ACKNOWLEDGEMENTS

The author very gratefully acknowledges the constant encouragement and the valuable guidance received from Shri S.D. Desai, Reader in Geology, Faculty of Science, M.S. University of Baroda.

The author is grateful to Prof. S.S. Merh, Head of the Geology Department and Dean of the Faculty of Science for taking interest and providing research facilities.

The author's grateful thanks are also due to Shri G.V.U. Rao, Head of the Ore Dressing Section, B.A.R.C., Trombay, Bombay for extending Pilot Plant testing facilities.

The author is also grateful to Shri Malkan, Head of the Chemical Laboratory, Directorate of Geology and Mining, Ahmedabad.

The author sincerely acknowledges the assistance offered by Dr. S.J. Desai, Shri D.M. Shringarpure, Dr. M.P. Patel of the Department of Geology.

The author sincerely acknowledges the manifold help received from Shri N.D. Desai, Shri S.R. Nayak and P.K. Patel.

The printing of photographs by Shri D.M. Chauhan, and decent typing of this thesis by Shri V.F. Bhavsar deserve special thanks.

The author is grateful to Shri Ajay Kumar and Shri Mulgibhai Patel whose hospitality and generous help made the author's stay and field work.

Financial assistance from University Grants Commission is also acknowledged.