

LIST OF TABLES

| Table no. | Title | Page no. |
|-----------|---|----------|
| 2.1 | Prevalence of lactose intolerance in various countries | 2.5 |
| 2.2 | Summary of tests for lactose intolerance | 2.12 |
| 2.3 | Milk production and per capita availability of milk in India | 2.24 |
| 2.4 | State-wise estimates of milk production | 2.26 |
| 2.5 | Plant categories that serve as basic foodstuff for plant drinks or milk substitutes | 2.29 |
| 2.6 | Number of Panelists and samples required for sensory test | 2.35 |
| 2.7 | Various sources of β -galactosidase including their optimum pH and temperature for hydrolyzing lactose | 2.37 |
| 4.2.1.1 | BMI range (Asia pacific) | 4.12 |
| 4.3.3 | Ingredients used along with their brand in each food preparation | 4.18 |
| 5.1.1 | General information of the respondents | 5.4 |
| 5.1.2 | Number of adults, children and elderly suffering from gastrointestinal symptoms experienced upon consumption of various food | 5.6 |
| 5.1.2.1 | Number of adults, children and elderly suffering from gastrointestinal symptoms experienced upon consumption of various milk and milk products food | 5.7 |
| 5.1.3 | Presence of lactose intolerance detected post HBT | 5.8 |

| | | |
|----------------|---|------------|
| 5.1.4 | Severity of lactose intolerance amongst the subjects | 5.9 |
| 5.1.5 | Mean values of HBAT readings of lactose intolerant and non-lactose intolerant subjects (children) | 5.11 |
| 5.1.5.1 | Mean values of HBAT readings of lactose intolerant and non-lactose intolerant subjects (adults) | 5.12 |
| 5.1.5.2 | Mean values of HBAT readings of lactose intolerant and non-lactose intolerant subjects elderly | 5.12 |
| 5.1.6 | Pearson's correlation between age and HBT reading | 5.15 |
| 5.1.7 | Pearson's correlation between age and gastrointestinal symptoms upon consumption of various food | 5.15 |
| 5.2.1 | Percent nutrient intake of RDA by adults, children and elderly population before and after supplementation of LHM | 5.19 |
| 5.2.1.1 | Mean value of nutrient intake of the lactose intolerant subjects before and after supplementation | 5.21 |
| 5.2.4 | Feedback upon acceptability of lactose hydrolysed milk | 5.27 |
| 5.3.1 | Organoleptic evaluation of beverages made out of standard milk and lactose hydrolysed milk | 5.30 |
| 5.3.2 | Organoleptic evaluation of white sauce pasta and veg au gratin made out of standard milk and LHM | 5.33 |
| 5.3.3 | Organoleptic evaluation of desserts (<i>kheer</i> and <i>Sandesh</i>) made out of standard milk and Lactose hydrolysed milk | 5.36 |