

List of Tables

<i>Table 2. 1: Samples prepared for the study</i>	<i>49</i>
<i>Table 2. 2: Composition of zirconia-alumina bi-component supports</i>	<i>53</i>
<i>Table 2. 3: Calculated Na/Zr and Zr/Al atomic ratio of zirconia-alumina bi-component supports from EDAX data</i>	<i>62</i>
<i>Table 2. 4: CHN analysis of GA and ZrO₂ series</i>	<i>79</i>
<i>Table 2. 5: Molar fraction of zirconium as carbonate and hydroxide and molar ratio of NO₃:Na derived from combined results of ICP-OES, Flame photometry and CHN analysis</i>	<i>80</i>
<i>Table 2. 6: Crystallite size of GA-7 Cal, GA-10 Cal, ZrO₂-7 Cal, ZrO₂-10 Cal, AM-7 Cal, AM-10 Cal, CP-7 Cal and CP-10 Cal</i>	<i>91</i>
<i>Table 2. 7: Comparison of reactivity of neat zirconium carbonate with different Na₂O content</i>	<i>99</i>
<i>Table 2. 8: Product selectivity of MBOH decomposition over substrate materials</i>	<i>100</i>
<i>Table 3. 1 : Bi-component alumina-zirconia supports with varying molar ratio</i>	<i>119</i>
<i>Table 3. 2: Composition of carriers used in the study</i>	<i>121</i>
<i>Table 3. 3: Crystallite size of t-zirconia and γ-Al₂O₃ in the supports.....</i>	<i>125</i>
<i>Table 3. 4: Values of specific surface area, pore volume and average pore diameter of the bicomponent zirconia-alumina supports.</i>	<i>125</i>
<i>Table 3. 5: Values of weak, strong and total acidity and corresponding peak temperature of desorption of the supports.</i>	<i>132</i>
<i>Table 3. 6 :Series of final catalysts prepared</i>	<i>137</i>
<i>Table 3. 7: % composition of catalysts by ICP-OES</i>	<i>140</i>
<i>Table 3. 8: Trend of relative intensity of XRD peaks for Co, Ni and Cu catalysts</i>	<i>145</i>
<i>Table 3. 9: Raman transitions of pure metal oxides</i>	<i>159</i>
<i>Table 4. 1: Components of products of styrene oxide reaction along with retention time</i>	<i>175</i>