

LIST OF TABLES

Table No.	Title	Page No.
Chapter 1	Introduction	
1.1	Potential contributors to nutritional problems in elderly people	8
Chapter 2	Review of Literature	
2.1	Proportion of older people in India (1901-2016)	20
2.2	Factors influencing nutritional inadequacy in elderly population	31
Chapter 3	Materials and Methods	
3.1	Parameters for processed food items	83
Chapter 4	Results and Discussion - Phase I	
4.1.1	Socio demographic status of subjects belonging to different living arrangements	103
4.1.2	Daily activity pattern of the elderly women as per their living arrangements and age group	106
4.1.3	Percentage of elderly women showing source of purchase of various food items (n=180)	110
4.1.4	Percentage of elderly women showing frequency of purchase of various food items (n=180)	111
4.1.5	Percentage of the elderly women showing influence of help in purchase activity of various foods	112
4.1.6	Percentage of subjects facing difficulties in purchasing of foods	113
4.1.7	Percentage of subjects with different living arrangements and age groups facing various difficulties in handling stored foods	115

4.1.8	Percentage of elderly women with different living arrangements and age groups facing different type of difficulties while using electrical appliances	117
4.1.9	Percentage of elderly women with different living arrangements and age groups facing different types of difficulties during cooking	119
4.1.10	Percentage of elderly belonging to different living arrangements and age groups showing different dietary practices	123
4.1.11	Percentage of elderly women showing data on frequency of substitutes for the meal with reasons (N=180)	124
4.1.12	Percentage of elderly women with different fasting pattern and their reasons (n=180)	125
4.1.13	Gastrointestinal tract related problems among elderly women belonging to different living arrangements and age groups as percentages	126
4.1.14	Percentage of elderly subjects with different living arrangements and age groups showing different adjustments with evening meal	127
4.1.15(a)	Percentage of elderly female subjects showing frequency of consumption of uncooked foods (N=180)	128
4.1.15(b)	Percentage of elderly women belonging to different living arrangements and age groups showing frequency of consumption of uncooked foods	129
4.1.16	Percentage of elderly women belonging to different living arrangement and age groups showing frequency of consumption of foods that require elaborate preparation	130
4.1.17	Percentage of elderly women with different living arrangements and age groups showing consumption of water per day	131

4.1.18	Mean nutrient intake of the elderly women with different living arrangements and age groups (mean \pm SD)	134
4.1.19	Mean anthropometric measurements of elderly women with different living arrangements and age groups	138
4.1.20	Percentage of subjects with different living arrangements and age groups showing prevalence of weight distribution	139
4.1.21	Mean hemoglobin values of the elderly women with different living arrangements and age groups (mean \pm SD)	139
4.1.22 (a)	Percent subjects showing prevalence of anemia	142
4.1.22 (b)	Percent prevalence of anemia in elderly subjects with different living arrangements and age groups	142
4.1.23	Mean blood glucose values and percentage of normal and abnormal blood glucose levels of the elderly women with different living arrangements and age groups (mean \pm SE)	144
4.1.24	Mean blood pressure values and percentage of normal and abnormal levels of the elderly women with different living arrangements and age groups (mean \pm SD)	145
4.1.25	Percentage of elderly women with different living arrangements and age groups showing self perceived minor health problems occurring frequently using checklist	147
4.1.26	Percentage of elderly with different living arrangements and age groups showing presence of major illnesses	150
Phase II		
4.2.1	Mean nutrient intake of the elderly anemic subjects in two groups before IFA supplementation (Mean \pm SD)	153
4.2.2	Mean nutrient intake of elderly anemic women before and after IFA supplementation (Mean \pm SD)	155
4.2.3	Percentage of elderly anemic women with IFA supplementation showing nutrient intake as percent RDA before and after intervention (n=45)	156

4.2.4	Mean hemoglobin levels of anemic women before and after intervention (mean \pm SE)	159
4.2.5	Percentage of elderly anemic subjects showing impact of IFA supplementation on the prevalence of degrees of anemia	160
4.2.6	Physical performance test score of anemic subjects before and after intervention with IFA supplementation for 6 weeks	163
4.2.7	Mental function test score of anemic subjects before and after intervention with IFA supplementation for 6 weeks	168
Phase III		
4.3.1	Mean scores of various soy and quick cooking food items on a 9 – point Hedonic scale	175
4.3.2	Protein, Fat and percent of RDA supplied by various food items (per serving)	178
4.3.3	Total Ash and Crude Fibre content of various food items	179
4.3.4	Calcium, Iron and percent RDA as supplied by various food items (per serving)	180
4.3.5	Beta carotene content and percent of RDA of quick cooking food items	181
4.3.6	Comparison of all the developed food items based on their sensory evaluation and nutritive composition	183
Phase IV		
4.4.1	Percentage of institutionalized elderly showing socio-demographic status	186
4.4.2	Percentage of institutionalized elderly visited by relatives and friends (n=20)	188
4.4.3(a)	Percent distribution of activity pattern of institutionalized elderly	189
4.4.3(b)	Percentage of institutionalized elderly involved in different activities in past and at present (n=20)	190

4.4.4	Percentage of institutionalized elderly showing past and present addiction pattern (n=20)	191
4.4.5	Mean anthropometric measurements of institutionalized elderly subjects before intervention	192
4.4.6	Percentage of subjects showing prevalence of weight distribution	192
4.4.7	Mean nutrient intake of institutionalized elderly male and female before intervention	193
4.4.8	Percentage consumption of nutrient intake as percent RDA by institutionalized elderly before intervention. (n=18)	194
4.4.9	Mean hemoglobin and serum total protein values of institutionalized elderly before intervention	195
4.4.10	Percentage of institutionalized elderly subjects showing prevalence of degree of anemia before intervention	196
4.4.11	Percent of institutionalized elderly subjects falling under normal and below normal serum total protein values	197
4.4.12	Mean lipid profile values of the institutionalized elderly before intervention	197
4.4.13	Percent of institutionalized elderly with normal and abnormal lipid profile before intervention (n=18)	198
4.4.14	Percentage of institutionalized elderly subjects showing major disease profile	199
4.4.15	Percentage of elderly institutionalized subjects using various prosthetic aids	201
4.4.16	Percentage of institutionalized elderly facing different psychological problems before intervention	202
4.4.17	Percentage of institutionalized elderly men and women reporting minor illnesses before intervention	204

4.4.18	Percentage of institutionalized elderly affected by mental health related factors in the institution	205
4.4.19	The meal pattern of institutionalized subjects	207
4.4.20	Mean anthropometric measurements of institutionalized elderly subjects before and after intervention of 6 weeks and 3 months	208
4.4.21	Percent prevalence of weight distribution in the institutionalized elderly before and after intervention of six weeks and three months (n=18)	209
4.4.22(a)	Mean nutrient intake of institutionalized elderly male before and after 6 weeks and 3 months of intervention (n=12)	212
4.4.22(b)	Mean nutrient intake of institutionalized elderly female before and after 6 weeks and 3 months of intervention (n=6)	212
4.4.23	Nutrient intake as percent RDA of institutionalized elderly before and after 6 weeks and 3 months of intervention (n=18)	214
4.4.24	Mean hemoglobin and serum total protein values of institutionalized elderly subjects before and after intervention (mean \pm sd)	219
4.4.25	Percentage of institutionalized elderly subjects showing prevalence of degree of anemia before and after three months of intervention	220
4.4.26	Percent of institutionalized elderly falling under normal and below normal total serum protein values before and after intervention of 3 months (n=18)	222
4.4.27	Mean values of the lipid profile of institutionalized elderly before and after intervention for 6 weeks and three months (n=18)	223
4.4.28	Mean random blood glucose levels of institutionalized elderly before and after intervention (n=18)	224

4.4.29	Mean blood pressure measurements of institutionalized elderly before and after intervention (n=18)	225
4.4.30	Percentage of elderly men and women reporting minor illnesses before and after three months of intervention	226
4.4.31	Percentage of institutionalized elderly showing psychological problems before and after three months of intervention	228