

List of Plates

Plate No.	Title	Page No.
Plate 3.1	Field photographs of the Cheriya Bet Conglomerate Member of Khadir Formation. A. Steeply inclined sandstone showing fining upward sequence with conglomerate at the base. (b) thinly laminated ferruginous sandstone (c) Conglomeratic sandstone showing with planar and cross-stratification (d) and (e) lumps of a polymictic clast dominated conglomerate.	40
Plate 3.2	Hadibhadang Shale Member of Khadir Formation. (a) Ferruginous Shale is exposed at the foothills of the north-facing scarp of Khadir Island. (b) Cross-bedded micritic Sandstone at the base of the north-facing scarp of Khadir Island. (c) Close up view of the micritic sandstone showing cross-bedding and liquefaction structure. (d) Micritic sandstone with palaeo-channel in the western tip of Khadir Island (e) Hadibhadang Shale Member exposed at the northern tip of Bela Island. (f) Hadibhadang Shale Member exposed at the core of an elliptical dome near Aaval village in Chorar Island.	42
Plates 3.3	Hadibhadang Sandstone Member of Khadir Formation. (a) Panoramic view of the north-facing scarp of Hadibhadang Sandstone Member of Khadir Formation in Khadir Island. Micritic sandstone at the base of Hadibhadang Sandstone Member in Khadir Island. (c) Thickly bedded micritic sandstone is exposed on the banks of the north-flowing Lelagar river in Bela Island. (d and e) Hard limestone at the top of micritic sandstone marked the top of the Hadibhadang Sandstone Member in Bela Island. (f) Sandy allochemic limestone with well developed (e) corals at the top of the member in Chorar Island.	44
Plate 3.4	Ratanpur Sandstone Member of the Gadhada Formation. (a) Micritic sandstone forms the back slope of Khadir Island. (b) Vertical cliff section formed by fluvial erosion on the back slope of Khadir Island. (c) Thickly bedded, friable micritic sandstone with trough cross-stratification in Bela Island. (d) Thick shale succession of Ratanpur Sandstone Member exposed near Dhabda village. (e) Thickly bedded friable sandstone of Ratanpur Sandstone Member in Chorar Island. Ferruginous sandstone containing (f) fossils wood and (g) iron concretion in Ratanpur Sandstone Member of Chorar.	48
Plate 3.5	Bambhanka Member of Gadhada Formation exposed only at the southern tip of Khadir Island and Kakinda Bet. (a) Sandy allochemic limestone at the southern tip of Khadir Island (b) Ferruginous sandstone containing abundant trace fossils. (c) Panoramic view of the hard bands forming ridges.	50

Plate 4.1	Field photographs of the bioclastic grainstone facies. (a), (b) and (c) exposed in the eastern part of Khadir Island near Umrapur village (d) Photomicrographs of bioclastic packstone grainstone facies with micritized bryozoan, (e) micritized algae (al), bioclast/Shell fragments (sf) and pellets (pl) and (f) highly micritized bioclast.	53
Plate 4.2	(a) Bioclastic wakestone-packstone facies of Hadibhadang Sandstone Member exposed at the northern top of Khadir island, dotted line separating bioclastic wakestone-packstone and peloidal packstone-wakestone (b) Bioclastic packstone-wackestone facies of Bambhanka Member exposed at Kakinda Bet and (c) Closed up view of event bands by the highly fossiliferous in nature. (d) Photomicrographs of bioclastic wakestone-packstone facies, bryozoans at the centre. (e) Photomicrographs showing gastropods (ga) and sparitised shell fragments (ssf) in the micritic medium. (f) Photomicrographs showing highly micritized shell fragments and gastropods (ga) where the internal cavities are filled with quartz, pellets and micrite.	54
Plate 4.3	Micritic sandstone facies of Hadibhadang Shale Member exposed (a) at the base of the northern cliff of Khadir Island, (b) Close up view of micritic sandstone show cross-stratification and (c) containing fossil wood in Hadibhadang Shale Member of Khadir Island. (d) Highly friable micritic sandstone showing prominent cross stratification and parallel lamination in Ratanpur Sandstone Member of Khadir Island. (e) Thickly bedded ferruginous micritic sandstone of Bambhanka Member exposed at the southern slope of Kakinda Bet. (f) Photomicrographs of ferruginised micritic sandstone containing, fine, well sorted quartz grains and pellets. (g) Micritic sandstone with quartz grain with point contact and echinoid spines. Photomicrographs of micritic sandstone with partially micritized algae.	56
Plate 4.4	(a) Photomicrographs of pelloidal wakestone-packstone containing algae, forams, onchoids polycrystalline quartz and reworked pellets. (b) Photomicrographs of pelloidal wakestone-packstone containing abundant shell fragments and algae.	57
Plate 4.5	(a) Field photograph of golden oolites in the form of pelloidal packstone-grainstone (b) Close up view of golden oolites observed in Ratanpur Sandstone Member on Khadir Island. (c) Photomicrographs of pelloidal packstone-grainstone with angular to subangular quartz (q) grains with abundant pellets (pl) and few shell fragments (sf).	57
Plate 4.6	(a) Close-up view of matrix supported polymictic conglomerate. (b) Field photograph of the clast of weathered polymictic conglomerate exposed in Cheriya Bet. (c) Thickly bedded, showing fining upward sequence with conglomerate at the base Showing 4-5 cycles of fining	58

upward sequence. (d) Photomicrographs of micro-faulting observed in quartz grains. (e) Photomicrographs of the clast of the conglomerate, basalt and (f) spherulite in rhyolite.

- | | | |
|------------|--|----|
| Plate 4.7 | Sandy allochemic limestone facies observed in Ratanpur Sandstone Member at the top of the northern cliff of Khadir Island. (b) Field photograph of Sandy allochemic limestone of Bambhanka Member exposed at the southern tip of Khadir Island. (c) Photomicrographs of sandy allochemic limestone facies with abundant shell fragment (sf) and angular to sub angular quartz (q) grains. (d) Photomicrographs of sandy allochemic limestone facies showing along the transverse section of the tubes with distinct burrow fill and the host rock separated by distinct dotted burrow wall as shown in dash line; (m=micrite) (e) Photomicrographs of sandy allochemic limestone facies with abundant pellets (pl) and shell fragments (sf). | 59 |
| Plate 4.8 | Greyish gypseous shale facies of Bambhanka Member observed at the (a) core below the hard band of highly weathered micritic sandstone and (b) the back slope of Kakinda bet. | 60 |
| Plate 4.9 | (a) Allochemic sandstone facies of Ratanpur Sandstone Member exposed near Neelagar river b) Photomicrograph of allochemic sandstone showing pellets (pl) with few shell fragments (sf) (c) Thickly bedded allochemic sandstone facies of Ratanpur Sandstone Member exposed on the eastern back slope of the Bela Island and its photomicrograph showing (d) with well sorted, fine grained quartz grains and abundant shell fragments (sf). | 65 |
| Plate 4.10 | Bioclastic packstone facies exposed at the top of eastern Bela Island at the top of Hadibhadang Sandstone Member. (b) Close up view of (a) showing abundant bivalve fossils. (c) Abundant well rounded bioclastic grains. (d) Photomicrograph shows highly sparitized shell fragment (ssf) with micritized shell fragments boundaries. | 66 |
| Plate 4.11 | Field photograph of micritic sandstone facies Ratanpur Sandstone Member exposed near Bela village showing planar laminated sandstone sandwich between bedded micritic sandstone. (b) Field photograph of thickly bedded micritic sandstone of Hadibhadang Sandstone Member exposed in the north facing scarp of Bela Island. (c) Field photograph highly friable, trough cross-stratified micritic sandstone facies exposed in the north facing scarp of Bela Island. (d) Photomicrograph of micritic sandstone containing polycrystalline quartz (pq) with sub-angular quartz grains floating in micrite (m). | 67 |
| Plate 4.12 | (a) Thinly bedded mudstone facies intercalated with shale of Ratanpur Sandstone Member exposed near Dhabda village, East of Bela. (b) Thickly bedded mudstone of Ratanpur Sandstone Member exposed on the back slope of Eastern Bela. (c) Photomicrograph of well- | 68 |

developed sparite in mudstone facies. (d) Photomicrograph of mudstone with negligible siliceous content.

- | | | |
|------------|---|----|
| Plate 4.13 | (a) Field photograph of sandy allochemic limestone exposed on the back slope of Bela Island showing ripple marks, (b) Highly bioturbated sandy allochemic limestone including <i>Diplocraterion</i> (di), <i>Skolithos</i> (sk) and <i>Taenidium</i> (ta) (c) Photomicrograph of sandy allochemic limestone with abundant pellets (pl), shell fragments (sf) and algae (al). (d) Photomicrograph of sandy allochemic limestone with abundant pellets, shell fragments (sf) and well developed elliptical ooids (o) with distinct growth rings. | 69 |
| Plate 4.14 | (a) Panoramic view of the north facing scarp of Bela Island (b) Shale facies of Hadibhadang Shale Member exposed at the base, showing flat topography of the north facing scarp of Bela Island. | 70 |
| Plate 4.15 | Allochemic Sandstone facies (a) Field photograph of thickly bedded allochemic limestone of Hadibhadang Sandstone Member. (b) Photomicrograph of allochemic sandstone containing abundant pellets (pl), and shell fragments (sf) with micrite (m) observed in Hadibhadang Sandstone Member (c) Field photograph of allochemic sandstone of Ratanpur Sandstone Member intercalated with shales and its corresponding Photomicrograph (d) containing algae (al), abundant pellets (pl) and micrite. | 73 |
| Plate 4.16 | Coralline limestone facies (a) Vertical section of a recrystallized corals embedded in the host rock. (b) Semi-circular broken boulder of corals. (c) Vertical section of a highly recrystallized corals embedded in the host rock. (d) Photomicrograph of cross section of the corals. The internal structures are not preserved due to diagenetic effect of recrystallization. (e) Photomicrograph of the host rock containing abundant dolomite and well-preserved echinoids spine (es). (f) Close up view of well-developed dolomite crystal. | 74 |
| Plate 4.17 | Cross-bedded white sandstone and ferruginous sandstone facies. (a) Thickly cross-bedded white sandstone facies of Ratanpur Sandstone Member expose on the south western margin of the Chorar Dome, showing faint mega trough cross-stratification. (b) Photomicrograph shows fine to medium grain, subrounded to rounded quartz in ferruginous matrix. | 75 |
| Plate 4.18 | (a) Field photograph of ferruginous sandstone facies showing concretionary structures. (b) Photomicrograph of ferruginous sandstone facies where the clasts are masked by ferruginous matrix | 76 |
| Plate 4.19 | Micritic sandstone facies (a) Micritic sandstone facies of Hadibhadang Shale Member exposed at the base of Chorar Dome showing cross-stratification. (b) Photomicrograph of micritic | 77 |

sandstone with polycrystalline quartz (pq), microcline (kf) and sub rounded quartz (q) grains. (c) Symmetrical ripples observed in micritic sandstone facies of Hadibhadang Sandstone Member (d) Petrographic photograph of micritic sandstone with shell fragments (sf), Plagioclast feldspar (pf) and with few occurrences of pellets (pl).

- | | | |
|------------|---|----|
| Plate 4.20 | Mudstone and sandy micrite facies. (a) Field photograph of mudstone facies of Ratanpur Sandstone Member occurring as thinly bedded hard band within shales. (b) Photomicrograph shows domination of the micritic material. | 78 |
| Plate 4.21 | (a) Field photograph of thickly bedded sandy micrite in Hadibhadang Sandstone Member of Khadir Formation exposed in the core of Chorar Dome (b) Photomicrograph of sandy micrite with sub-rounded to well-rounded quartz grain floating in micritic matrix. | 79 |
| Plate 4.22 | Sandy allochemic limestone facies (a) Field photograph of thickly bedded sandy allochemic limestone of Hadibhdang Sandstone Member exposed along the eastern side of Chorar Dome. (b) Thickly bedded sandy allochemic limestone with layers of sudden increase in chaotic fossil content representing sudden change in energy condition. (c) Close up view of liquefaction within the fossiliferous horizon. (d) Photomicrograph of sandy allochemic limestone showing abundant shell fragments (sf) and well-developed algae (al) with well sorted quartz (q) grains | 80 |
| Plate 4.23 | Field photograph of thick gypseous shale at the core of Chorar Dome. | 81 |
| Plate 5.1 | (a) Sub-horizontal protruding bulbus tubes of <i>Aesterosoma</i> isp. in sandy allochemic limestone facies of Hadibhadang Sandstone Member of Khadir Formation in the Chorar Island (Scale: coin diameter = 2.3 cm). (b) Faintly preserved trilobate <i>Curvolithus multiplex</i> (arrow) in sandy allochemic limestone of Hadibhadang Sandstone Member in Chorar Island (Scale: coin diameter = 2.3 cm). (c) <i>C. multiplex</i> (arrow) in sandy allochemic limestone of Ratanpur Sandstone Member in Khadir Island. (d) trilobate <i>Curvolithus simplex</i> in micritic sandstone of Bambhanka Member in Kakinda Bet. (e) Partially preserved trilobate <i>Curvolithus</i> isp. in sandy allochemic limestone of Hadibhadang Sandstone Member in Chorar Island (Scale: coin diameter = 2.3 cm). | 90 |
| Plate 5.2 | (a) <i>Didymaulichnus lyelli</i> with faint median groove observed in sandy allochemic limestone facies of Hadibhadang Sandstone Member of Khadir Formation in Chorar Island. (b) <i>D. lyelli</i> (Dy) with distinct median groove observed in association with <i>Rhizocorallium</i> (Rh) in micritic sandstone facies of Bambhanka Member. (c) Sinous, <i>Didymaulichnus alternatus</i> in micritic sandstone facies of Bambhanka Member in Kakinda Bet showing faintly preserved undulating | 95 |

bilobate structure as shown by the dashed line (Scale: coin diameter = 2.3 cm). and (d) Bilobate *Gyrochorte comosa* in Hadibhadang Sandstone Member of Khadir Formation in Chorar Island (Scale: coin diameter = 2.3 cm).

- Plate 5.3 (a) Irregularly wrinkled *Halopoa imbricata* in sandy micrite facies of Hadibhadang Sandstone Member in Chorar Island (Scale: coin diameter = 2.3 cm). (b) Equidimensional grooves arranged in linear fashion at equidistant of *Helicolithus sampelayoi*, in micritic sandstone facies of Ratanpur Sandstone Member in Bela Island (Scale: coin diameter = 2.3 cm). 97
- Plate 5.4 (a) *Megagraption irregulare* branching almost perpendicularly in sandy allochemic limestone facies of Hadibhadang Sandstone Member in Chorar Island (represented by dotted lines). (b) Horizontal thin lining *Palaeophycus tubularis* (pa) in sandy allochemic limestone facies in Ratanpur Sandstone Member in Bela Island. (c) *P. tubularis* in ferruginous sandstone facies in Ratanpur Sandstone Member in Chorar Island. (d). *P. tubularis* (Pa) in association with *Curvolithus simplex* (Cu) in micritic sandstone facies of Bambhanka Member in Kakinda Bet. (e) Thickly lined *P. heberti* in sandy allochemic limestone of Hadibhadang Sandstone Member in Chorar Island (Scale: coin diameter = 3.2 cm). (f) Horizontal *P. heberti* with thick lining which is preserved as a groove around the central tube, observed in peloidal packstone/grainstone of Ratanpur Sandstone Member, in Khadir Island (Scale: coin diameter = 3.2 cm). 100
- Plate 5.5 (a) Short and curvy bifurcating burrow *Phycodes curvipalmatum* in micritic sandstone facies of Bambhanka Member in Kakinda Bet. (b) Large digitating burrow *Phycodes palmatus* in sandy allochemic limestone of Ratanpur Sandstone Member in Bela Island. Horizontal, unlined burrows of *Planolites baverleyensis* in (c) sandy allochemic limestone facies of Hadibhadang Sandstone Member of Khadir Formation in Chorar Island (Scale: coin diameter = 3.2 cm) (d) micritic sandstone in association with *Skolithos* (Sk) and *Arenicolites* (Ar) and (e) sandy allochemic limestone facies (Scale: coin diameter = 3.2 cm) Ratanpur Sandstone Member in Bela Island and (f) micritic sandstone facies of Bambhanka Member in Khadir Island (Scale: coin diameter = 3.2 cm). (g) Contorted burrows of *P. montanus* in sandy allochemic limestone facies of Ratanpur Sandstone Member in Khadir Island. 104
- Plate 5.6 (a) Horizontal *Protovirgularia dichotoma* with prominent deep median furrow in micritic sandstone facies of Bambhanka Member in Kakinda Bet (bar = 5cm). (b) *Protovirgularia obliterata* showing faint median groove in sandy allochemic limestone facies of Ratanpur Sandstone Member in Bela Island. (c) *Protovirgularia* isp. in sandy 107

allochemic limestone facies of Hadibhadang Sandstone Member of Khadir Formation in Chorar Island (Scale: coin diameter = 3.2 cm).

- Plate 5.7 (a) *Rhizocorallium commune* var. *auriform* showing faint spreiten structure in thinly bedded peloidal packstone/grainstone of Ratanpur Sandstone Member in Khadir Island. (b) Twisted *Rhizocorallium commune* var. *auriform* in thickly bedded micritic sandstone of Bambhanka Member in Kakinda Bet showing prominent spreiten structure. (c) Straight, elongated *R. commune* var. *irregularie* in sandy allochemic limestone facies of Hadibhadang sandstone Member in Chorar Island where the spreiten structures between the tubes have been obscured due to erosion. (d) *R. commune* var. *irregularie* horizontally preserved in sandy allochemic limestone facies of Ratanpur Sandstone Member in Bela Island. (e) Highly ferruginised *R. commune* var. *irregularie* observed in micritic sandstone facies of Bambhanka Member in Khadir Island. (f) Horizontal, partially preserved *R. commune* isp. in micritic sandstone of Hadibhadang Shale Member in Khadir Island. 109
- Plate 5.8 (a) Horizontal and sinuous *Taenidium serpentinum* in thinly bedded sandy allochemic limestone facies of Ratanpur Sandstone Member in Bela Island where the meniscate structure is perpendicular to the tube and is often obscure by erosion. (b) *T. serpentinum* in peloidal packstone/grainstone of Ratanpur Sandstone Member in Khadir Island with distinctly preserved meniscate structure while it is poorly preserved in micritic sandstone facies of Bambhanka Member. (c) in Khadir Island (d) Highly sinous *Taenidium*. isp. where the meniscate structure is obscure by ferruginous material in micritic sandstone of Ratanpur Sandstone Member in Khadir Island (Scale: coin diameter = 3.2 cm). 111
- Plate 5.9 Vertical U-shapped burrows without spreiten structure *Arenicola carbonaria* (Ar). (a) in association with *Skolithos* (Sk) in ferruginous sandstone facies of Ratanpur Sandstone Member in Chorar Island. (b) micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island. (c) *Arenicola carbonaria* (Ar) in micritic sandstone facies in association with *Skolithos verticalis* (Sk) and *Lockeia* (Lo) of Ratanpur Sandstone Member in Bela Island. (d) Vertical lined pared tube *Arenicolites statheri* (indicated by an arrow) where the lining to the vertical is closely spaced or touch each other, observed in sandy allochemic limestone facies of Ratanpur Sandstone Member in Bela Island. (e) Very small size *Arenicolites* isp. represented by a circle in association with *Skolithos* (Sk), observed in micritic sandstone facies of Hadibhadang Shale Member Island and (f) Ratanpur Sandstone Member in Khadir. 114

- Plate 5.10 (a) Radiating branches of *Chondrites intricatus* in micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island. (b) Meandering burrows *Chondrites targionii* in sandy allochemic limestone of Ratanpur Sandstone Member (Scale: coin diameter = 3.2 cm). (c) Vertical view of *C. targionii* in sandy allochemic limestone of Ratanpur Sandstone Member. (d) Close-up view of *C. targionii* bearing sandy allochemic limestone facies in Ratanpur Sandstone Member in Khadir Island (Scale: length of pen = 14 cm). 116
- Plate 5.11 (a) Vertical paired burrows of *Diplocraterion parallelum* (Di) in association with *Monocraterion* (Mo) in micritic sandstone facies of Ratanpur Sandstone Member in Bela Island (Scale: coin diameter = 2.3 cm). (b) *Laevicyclus parvus* showing the concentric ring in micritic sandstone of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm) (c) Almond-shaped *L. siliquaria* observed in ferruginised sandy allochemic limestone of Ratanpur Sandstone Member in Khadir Island. (Scale: coin diameter = 2.3 cm). (d) Funnel-shaped vertical tube, *Monocraterion tentaculatum* (Mo) in association with *Arenicolites statheri* (Ar) in sandy allochemic limestone of Ratanpur Sandstone in Bela Island. (e) Funnel-shaped *M. tentaculatum* (Mo) observed in micritic sandstone of Ratanpur Sandstone Member in Khadir Island in association with *Diplocraterion* (Di) and *Skolithos* (Sk). 119
- Plate 5.12 (a) Vertical *Skolithos linearis* in cross-bedded white sandstone facies of Ratanpur Sandstone Member in Chorar Island (Scale: Bar = 5 cm). (b) Straight and vertical *S. linearis* in cross-bedded white sandstone facies of Ratanpur Sandstone Member in Chorar Island (Scale: Bar = 5 cm). (c) Short, curved, and densely populated *Skolithos verticalis* in sandy allochemic limestone facies of Ratanpur Sandstone Member in Bela Island (Scale: Bar = 5 cm). (d) Vertical *Skolithos* isp. in micritic sandstone facies of Bambhanka Member in Kakinda Bet (Scale: coin diameter = 3.2 cm). 123
- Plate 5.13 (a). Level A of *Hillichnus lobosensis* shows the basal segmented structure in sandy allochemic limestone of the Bela Island. (b). *H. lobosensis* shows the basal tubes of Level B. (c) Level B of *H. lobosensis* shows dark brown lined, undulated, wrinkled, irregular, and elongated tubes overlapping with leaf-like branches of Level C. 126
- Plate 5.14 (a). Level C of *H. lobosensis* shows well-developed horizontal leaf or feather branches with prominent undulated and wrinkled basal tubes (Scale: coin diameter = 3.2 cm). (b). Leaf-like branches are smoothened by erosion in the Chorar Island. (c and d) Less prominent leaf-like Level C of Khadir (Scale: coin diameter = 3.2 cm). (e). Overlapping of Level C of *H. lobosensis* with a prominent basal tube 127

with the overlying sediments and the crowded nature masking the leaf-like branches in the Bela Island (f) Level C with remnants of a broken basal tube.

- Plate 5.15 (a) Level D of *H. lobosensis* (a) the rising tube arises in pairs and (b) with prominent basal tube where the rising tubes arise from the side in an alternate pattern in the Khadir Island. (c) Crowded Level D of *H. lobosensis* with the prominent basal tube but eroded inclined tube, (d) Level D of *H. lobosensis* occurring parallel with a series of vertical tubes belonging to Level E. (e) Level D of *H. lobosensis* showing prominent basal tube with the inclined tube arising from the side in the Bela Island. (f) Crowded Level D of *H. lobosensis* where the basal tube is concealed while the inclined tube occurs in pair or zigzag. (g). The transition of level C to Level D. (h) Vertical view of *H. Lobosensis* in Chorar Island showing level D on top with the basal tube appearing as horizontal tubes and the feather-like protrusion from Level C appearing as thin curved lines. 129
- Plate 5.16 (a) Isolated (i) or series (dash line) of small vertical tubes belonging to Level E of *H. lobosensis* in Khadir Island. (b) Series of vertical tubes occurring in a straight line or in pairs, Bela Island (Scale: Coin diameter = 2.3 cm). 130
- Plate 5.17 (a) Elongated pellets of *Ophiomorpha annulata* in micritic sandstone of Ratanpur Sandstone Member with elongated pellets (Scale: coin diameter = 2.3 cm). (b) T or Y-shaped *Ophiomorpha irregulaire* showing less dense pellets in micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island. (c) *Ophiomorpha nodosa* showing regularly dense pellets in micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island (Scale: coin diameter = 2.3 cm). (d) Highly ferruginised *Ophiomorpha* isp. in micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island (Scale: coin diameter = 2.3 cm). 133
- Plate 5.18 (a) T/Y shaped *Thalassinoides horizontalis* in sandy allochemic limestone facies of Ratanpur Sandstone Member in Khadir Island and (Scale: coin diameter = 3.2 cm). (b) Networks of *T. horizontalis* in sandy allochemic limestone facies of Ratanpur Sandstone Member in Chorar Island (length of hammer = 32 cm). (c) Horizontal Y-shaped *T. paradoxicus* with swollen nodes in micritic sandstone facies of Hadibhadang Sandstone Member in Chorar Island (Scale: coin diameter = 3.2 cm). (d) Field photograph of *Thalassinoides* isp. in Ratanpur Sandstone Member of Bela Island (Scale: coin diameter = 3.2 cm). 136
- Plate 6.1 Trace fossils of *Diplo craterion* assemblage: (a) Association of *Diplo craterion* (Di), *Arenicolites* (Ar) and *Skolithos* (Sk), (b) 146

Arenicolites (Ar) and *Skolithos* (Sk) in a thinly bedded micritic sandstone of Hadibhadang Shale Member.

- Plate 6.2 Trace fossils of *Skolithos* assemblage. (a) An association of *Skolithos* (Sk) and *Arenicolites* (Ar) in a thickly bedded micritic sandstone facies of Hadibhadang Shale Member exposed at the base of the north facing scarp of Khadir Island. (b) Monodominant *Skolithos* in micritic sandstone facies of Hadibhadang Shale Member. 147
- Plate 6.3 Trace fossils of *Lockeia* assemblage. (a) Monodominant *Lockeia* in micritic sandstone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (b) *Planolites* in highly ferruginised micritic sandstone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (c) *Rhizocorallium* in ferruginised micritic sandstone where the spreiten structures have been destroyed due to erosion (Scale: coin diameter = 2.3 cm). 148
- Plate 6.4 Trace fossils of *Planolites-Palaeophycus* assemblage. (a) *Palaeophycus* observed in sandy allochemic limestone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (b) *Planolites* in association with *Arenicolites* and *Skolithos* (Scale: coin diameter = 2.3 cm). (c) *Ophiomorpha* observed in sandy allochemic limestone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (d) *Chondrites* (Ch) in association with *Arenicolites* (Ar) and *Skolithos* (Sk). (Scale: coin diameter = 2.3 cm). (e) *Thalassinoides* in association with *Lockeia* in thinly bedded sandy allochemic limestone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (f) Faintly preserved trilobate *Curvolithus* in association with *Thalassinoides* thinly bedded sandy allochemic limestone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). 150
- Plate 6.5 Trace fossils of *Rhizocorallium* assemblage. (a) *Rhizocorallium* observed in micritic sandstone facies of Bambhanka Member. (b) *Taenidium* (Ta) in association with *Arenicolites* (Ar) in peloidal packstone-grainstone facies of Ratanpur Sandstone Member exposed towards the southern tip of Khadir Island. (c) *Planolites* (Pl) in association with *Diplocraterion* (Di) and *Monocraterion* (Mo). (d) *Phycodes*, (e) *Ophiomorpha*, (f) *Laevicyclus*, (g) *Thalassinoides* and (h) *Skolithos* observed in micritic sandstone facies of Bambhanka Member in Kakinda Bet (Scale: coin diameter = 2.3 cm). 152
- Plate 6.6 Trace fossils of *Protovirgularia* assemblage. (a) *Protovirgularia* (Pr) in association with *Gyrochorte* (Gy) in sandy allochemic limestone facies of Bambhanka Member. (b) *Didymaulichnus* observed in sandy allochemic limestone facies of Bambhanka Member (Scale: coin diameter = 2.3 cm). (c) *Protovirgularia* (Scale: hammer = 35 cm). (d) 154

Planolites (Pl) in association with *Diplocraterion* (Di), (e) *Gyrochorte* and (f) *Taenidium* are observed in micritic sandstone facies of Bambhanka Member.

- Plate 6.7 Trace fossils of *Hillichnus* assemblage. (a) Level C of *Hillichnus* in micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island (Scale: coin diameter = 2.3 cm). (b) *Arenicolites* observed in micritic sandstone facies of Ratanpur Sandstone Member in Khadir Island (Scale: coin diameter = 2.3 cm). (c) *Chondrites* observed in micritic sandstone facies of Ratanpur Sandstone Member. (d) *Planolites* in thinly bedded micritic sandstone facies in Ratanpur Sandstone Member. (e) *Thalassinoides* observed in micritic sandstone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (f) *Taenidium* in micritic sandstone facies exposed in the western part of Khadir Island (Scale: coin diameter = 2.3 cm). (g) *Laevicyclus* observed in micritic sandstone facies exposed in the back slop of Khadir Island (Scale: coin diameter = 2.3 cm). (h) *Gyrochorte* observed in thinly bedded micritic sandstone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (i) *Diplocraterion* (Di) in association with *Monocraterion* (Mo), *Arenicolites* (Ar) and *Skolithos* (Sk). 155
- Plate 6.8 Trace fossils of *Monocraterion* assemblage in Bela Island. (a) *Diplocraterion* observed in association with *Monocraterion* in micritic sandstone facies in Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (b) *Rhizocorallium* (Rh) observed in association with *Monocraterion* (Mo) and *Skolithos* (Sk) in a rippled micritic sandstone facies in Ratanpur Sandstone Member. (c) *Helicolithus* (Scale: coin diameter = 2.3 cm). (d) *Arenicolites* (Scale: coin diameter = 2.3 cm). (e) *Lockeia* (Scale: coin diameter = 2.3 cm). (f) *Planolites* in association with *Skolithos* (g) *Palaeophycus*, (h) *Thalassinoides* (Scale: coin diameter = 2.3 cm). (i) *Monocraterion* (Mo) observed in association with *Arenicolites* (Ar) and *Skolithos* (Sk) in micritic sandstone facies in Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). 158
- Plate 6.9 *Hillichnus* assemblage observed in Bela Island. (a) Level C of *Hillichnus* in association with Level E. (b) Pared tubes *Arenicolites* in association with *Diplocraterion*. (c) *Planolites* (d) *Palaeophycus* with ferruginised lining in association with *Diplocraterion*. (e) *Taenidium*. (f) *Phycodes*. 160
- Plate 6.10 Mono occurrence of *Thalassinoides* in micritic sandstone facies of Hadibhadang Shale Member in Bela Island. 161
- Plate 6.11 Trace fossils for *Hillichnus* assemblage (Scale: coin diameter = 2.3 cm). (a) Level C of *Hillichnus* and (b) *Protovirgularia* observed in 162

sandy allochemic limestone of Hadibhadang Sandstone Member
(Scale: coin diameter = 2.3 cm).

- | | | |
|------------|---|-----|
| Plate 6.12 | Trace fossils for <i>Rhizocorallium</i> assemblage. (a) <i>Rhizocorallium</i> (Scale: coin diameter = 2.3 cm) (b) <i>Curvolithus</i> in association with <i>Planolites</i> (Scale: coin diameter = 2.3 cm) (c) <i>Gyrochorte</i> in sandy allochemic limestone of Hadibhadang Sandstone Member of Chorar Island (Scale: coin diameter = 2.3 cm). | 163 |
| Plate 6.13 | Trace fossils for <i>Gyrochorte</i> assemblage. (a) <i>Gyrochorte</i> (Scale: coin diameter = 2.3 cm) (b) Paired tubes of <i>Arenicolites</i> (Ar) in association with <i>Lockeia</i> (Lo) and <i>Planolites</i> . | 164 |
| Plate 6.14 | <i>Thalassinoides</i> assemblage. (a) Predominant occurrence of <i>Thalassinoides</i> at the sediment-sediment interface of sandy allochemic limestone (Scale: hammer=15 cm). (b) Wrinkled with discontinuous striation of <i>Halopoa</i> in sandy allochemic limestone of Hadibhadang Sandstone Member in Chorar Island (Scale: coin diameter = 2.3 cm). | 165 |
| Plate 6.15 | Trace fossils of <i>Skolithos</i> assemblage. (a) Highly weathered <i>Planolites</i> (Scale: coin diameter = 2.3 cm). (b) Highly weathered <i>Thalassinoides</i> burrows are observed in friable cross-bedded white sandstone facies of Ratanpur Sandstone Member (Scale: coin diameter = 2.3 cm). (c) <i>Skolithos</i> (Sk) in association with <i>Arenicolites</i> (Ar) (Scale: bar = 5 cm) and <i>Diplocraterion</i> (Di). (d) <i>Palaeophycus</i> observed in ferruginous facies of Ratanpur Sandstone Member in Chorar Island. | 167 |