ACKNOWLEDGEMENT

I wish to express my profound gratitude to my guide, **Prof. H.L. Bhalla**, Professor of Pharmaceutics, and Founder Director of the B.V. Patel PERD Centre. He diligently took the pivotal role to accomplish this work by furnishing excellent guidance, portentous suggestions, unstinted help and critical criticisms.

I am profoundly indebted to *Dr.(Ms)*. Anisha Pargal, Director, B.V. Patel PERD Centre, for her encouragement, valuable comments and for providing excellent laboratory facilities to carry out my research, throughout my graduate studies.

I am grateful to Dr. Bakulesh Khamar, for his help in the in vivo studies and V.P. Shedbalkar for his support to carry out instrumental analysis. I would like to thank Dr. Shubha Rani who very kindly has gone through the data and suggested the most suitable statistical treatment to the problem.

I would like to thank my colleagues and staff of the B.V. Patel PERD Centre for their co-operation and goodwill throughout the tenure of the study.

I am thankful to **Syntex Research**, U.S.A for the gift samples of degradation products of ketorolac. The excellent library facilities and gift samples of drugs provided by **Cadila** *Laboratories Ltd.* is gratefully acknowledged.

I wish to thank BASF (I) Ltd. and BF-Goodrich Co Ltd. U.S.A, for providing gift samples of polymers.

Finally, I can not in words thank my parents. To them I dedicate my thesis.