

LIST OF FIGURES

Figure	Page
1. The relative size of body iron compartments in a 75 kg adult male.	14
2. Schematic representation of the intestinal absorption of iron.	16
3. A concept of control of iron absorption by the intestinal mucosa, in normal, iron deficient and iron loaded subjects.	19
4. Iron requirements in man.	25
5. Schematic diagram of the absorption of the two forms of food iron; heme iron and nonheme iron.	42
6. Magnitude of increase in absorption of nonheme iron with ascorbic acid.	51
7. Dose effect of tannin on relative absorption of iron, calculated as the ratio of with to without addition of tannin to the meal.	71
8. Structural formulae of a) gallic acid and b) catechin.	72
* 9. Decreased absorption of nonheme iron in healthy adult men who were presented with 700-kcal semi-synthetic meals with the addition of calcium and/or phosphate salts.	90
10. Effect of calcium (CaCl_2) on iron absorption from wheat rolls, expressed as the ratio of with to without added calcium to the rolls.	92
11. Correlation between analysed in vitro iron availability vis-a-vis in vivo iron absorption from 12 selected meals, as reported by the respective authors (Phase I).	157
12. Dose relationship of enhancers and inhibitors with % iron availability (%IA) from pure system (Phase II).	161
13. Regression analysis of the interaction effect of various enhancers and inhibitors on % iron availability from the pure system (Phase III).	165

14.	Dose relationship of enhancers and inhibitors with % iron availability (%IA) from standard meal (Phase IV).	170
15.	Effect of bran, dephytinised bran or mono-ferric-phytate on % iron availability from pure system (sub study under Phase IV).	176
16.	Regression analysis of the interaction effect of various enhancers and inhibitors on % iron availability from the standard meal (Phase V).	179
17.	Correlation between analysed % iron availability (% IA) vs computed % iron availability (% CIA) using equation 1 and 2, incorporating analysed (A) or food table values (B) of enhancers and inhibitors.	186
18.	Correlation between the analysed in vitro iron availability vs bioavailability of iron, computed from the model of Monsen and Balintfy (1982).	190
19.	Comparison of dose effect of ascorbic acid on iron availability in the present study with that reported by other in vitro and in vivo studies.	200
20.	Comparison of dose effect of tannin on iron availability in the present study with that reported by other authors.	208
21.	Comparison of dose effect of calcium and phosphate added, to FeCl ₃ / STD meal in the present study with that of calcium per se, reported by other authors.	212