

REFERENCES

- Acharyya, S. K., Kayal, J. R., Roy, A. and Chaturvedi, R. K. (1998) Jabalpur earthquake of May 22, 1997: constraint from a aftershock study. *J. Geol. Soc. Ind.*, v. 5, pp. 295-304.
- Acharyya, S. K. and Roy, A., (2000) Tectonothermal history of the central Indian tectonic zone and reactivation of major faults/shear zones. *J. Geol. Soc. Ind.*, v. 55, pp. 239-256.
- Agarwal, B. N. P., Das, L. K., Chakraborty, K. and Sivaji, C. H. (1995) Analysis of the bouguer anomaly over coastal India: a regional perspective. *Mem. Geol. Soc. Ind.*, v. 31, pp. 469-493.
- Agarwal, G. C. (1986) Structure and tectonics of exposed Tertiary rocks between Narmada and Kim rivers in South Gujarat. *J. Geol. Soc. Ind.*, v. 27, pp. 531-542.
- Ahmad, F. (1968) Distribution and causes of gully erosion in India. Twenty first I.G.U. selected papers, v. 1, pp.103.
- Allchin, B., Goudie, A. and Hegde, K. T. M. (1978) The prehistory and palaeogeography of the Great Indian Thar Desert. Academic Press, London, 370pp.
- Allen, G. P. and Posamentier, H. W. (1993) Sequence stratigraphy and facies model of an incised valley fill, the Gironde estuary, France. *J. Sedimen. Petrol.*, v. 63, pp. 378-391.
- Allen, J. R. L. (1963) The classification of Cross-Stratified units, with notes on their origin. *Sedimentology*, v. 2, pp. 93-114.
- Allen, J. R. L. (1965) A review of the origin and characteristics of recent alluvial sediments. *Sedimentology*, v. 5, pp. 89-191.
- Al-Sarawi, M. (1988) Morphology and facies of alluvial fans in Kadhmah Bay, Kuwait. *J. Sediment. Petrol.*, v. 58, pp. 902-907.
- Andrews, J. E., Singhvi, A. K., Kaliath, A. J., Kuhn, R., Dennis, P. F., Tandon, S. K. and Dhir, R. P. (1998) Do stable isotope data from calcrete record Late Pleistocene monsoonal climate variation in the Thar desert of India? *Quat. Res.*, v. 50, pp. 240-251.
- Ashley, G. M. and Renwick, W. H. (1983) Channel morphology and processes at the riverine-estuarine transition, the Raritan river, New Jersey, in Collinson, J.D. and Lewin, J., eds., *Modern and ancient fluvial systems: International Association of Sedimentologists, Special Publication*, v. 6, pp. 207-218.
- Auden, J. B. (1949) Dykes in western India. *Trans. Nat. Inst. Sci. Ind.*, v. 3, pp. 123-157.
- Autin, W. J. (1996) Pleistocene stratigraphy in the southern Lower Mississippi Valley. *Eng. Geol.*, v. 45, pp. 87-112.
- Badam, G. L., Ganjoo, R. K. and Salahuddin (1986) Preliminary taphonomical studies of some Pleistocene Fauna from the Central Narmada Valley, Madhya Pradesh, India. *Palaeogeog. Palaeoclimat. Palaeoecol.*, v. 53, pp. 335-348.

- Bedi, N. and Vaidyanadhan, R. (1982) Effects of neotectonics on the morphology of Narmada river in Gujarat, Western India, Z.F. Geomorphologie, v. 26, pp. 87-102.
- Behrensmeyer, A. K. (1987) Miocene fluvial facies and vertebrate taphonomy in Northern Pakistan. In: Ethridge, F.G., Flores, R. M. and Harvey, M.D. (Eds.), Recent developments in Fluvial Sedimentology: SEPM Special Publication, v. 39, pp. 169-176.
- Bhandari S., Raj, R., Maurya, D. M. and Chamyal, L. S. (2001) Formation and Erosion of Holocene Alluvial fans along Narmada – Son Fault near Rajpipla in lower Narmada basin, western India. Jour. Geol. Soc. Ind., v. 58, pp. 519-531.
- Bhandari, S., Maurya, D. M. and Chamyal, L. S. (2004) Late Pleistocene alluvial plain sedimentation in lower Narmada valley, Western India: Palaeoenvironmental implications. Jour. of Asian Earth Sciences (in press).
- Bhattacharji, S., Chatterji, N., and Wampler, J. M. (1996) Zones of Narmada-Tapi rift reactivation and Deccan volcanism: geo-chronological and geochemical evidence. In: Deshmukh, S. S., Nair, K.K.K. (ed), Deccan Basalts. Gondwana Geological Society, pp. 329-340.
- Biswas, S. K. and Deshpande, S. V. (1983) Geology and Hydrocarbon prospects of Kutch, Saurashtra and Narmada Basins. In. L.L.Bhandari, B.S.Venkatachala, R.Kumar, S.N.Swamy, P. Garga and D.C. Srivastava (Eds.) Petroliferous Basins of India. Pet. Asia J., v. 6(4), pp. 111-126.
- Biswas, S. (1997) Fossil Mammalia of the Quaternary sequence of the Narmada Valley: Their affinity, age and ecology. GSI Spl. Publ., v. 46, pp. 91-104.
- Biswas, S. K. (1982) Rift basins in western margin of India and their hydrocarbon prospects with special reference to Kutch Basin. J. Am. Assoc. of Petrol. Geol., v. 10, pp. 1497-1513.
- Biswas, S. K. (1987) Regional tectonic framework, structure and evolution of western marginal basins of India. Tectonophysics, v. 135, pp. 307-327.
- Blair, T. C. and Bilodeau, W. L. (1988) Development of tectonic cyclothsems, in rift, pull-apart and foreland basins: sedimentary response to episodic tectonism. Geology, v. 16, pp. 517-520.
- Blair, T. C. and McPherson, J. G. (1994) Alluvial fans and their natural distinction from rivers based on morphology, hydraulic processes, sedimentary processes and facies. Jour. Sed. Res., v. A 64, pp. 451-490.
- Blanford, W. T. (1869) On the geology of the Tapti and Lower Narmada valleys and some adjoining districts. Mem. Geol. Surv. Ind., v. 6, pp. 163-384.
- Bloom, A. L. (2002) Geomorphology A systematic analysis of Late Cenozoic landforms. Prentice-Hall of India, pp. 482.
- Blum, M. D., Guccione, M. J., Wysocki, D. A., Robnett, P. C. and Rutledge, E. M. (2000) Late Pleistocene evolution of the lower Mississippi valley, southern Missouri to Arkansas. Geol. Soc. Amer. Bull., v. 112, pp. 221-235.

- Blum, M. D. and Tornquist, T. E. (2000) Fluvial responses to climate and sea level change: a review and look forward. *Sedimentology*, v. 47, pp. 2-48.
- Blum, M. D. Toomey, R. S. and Valastro, S. (1994) Responses of fluvial systems to Late Quaternary climatic changes. Edwards Plateau, Texas. *Palaeogeog. Palaeoclimat. Palaeoecol.*, v. 108, pp. 1-21.
- Bose, M. K. (1980) Magnetically defined tectonic framework of the Deccan Volcanic province. *News Mineral.*, v. 8, pp. 373-384.
- Bown, T. W. and Kraus, M. J. (1987) Integration of channel and floodplain suites, I. Developmental sequence and lateral relations of alluvial palaeosols. *Jour.Sed. Petr.*, v. 57, pp. 587-601.
- Bridge, J. S. (1984) Large-scale facies sequences in alluvial overbank environments. *Jour. Sed. Petr.*, v. 54, pp. 583-588.
- Bridge, J. S., Jalfin, G. A. and Georgieff, S. M., (2000) Geometry, lithofacies, and spatial distribution of Cretaceous Fluvial Sandstone Bodies, San Jorge Basin, Argentina: Outcrop analog for the hydrocarbon-bearing Chubut Group. *J. Sed. Res.*, v. 70 (2), pp. 341-359.
- Brierley, G. J., Keyu Liu and Crook, K. A. W. (1993) Sedimentology of coarse grained alluvial fans in Markham valley, Papua New Guinea. *Sediment. Geol.*, v. 86, pp. 297-324.
- Brown, G. H. and Plint, A. G. (1994) Alternating braid plain and lacustrine deposition in a strike-slip setting: the Pennsylvanian Boss Point Formation of the Cumberland basin, Maritime Canada. *Jour. Sediment. Petrol.*, v. 64, pp. 40-59.
- Bryson, R. A. (1989) Modelling the NW India Monsoon for the last 40000 years. *Clim Dyn*, v. 3, pp. 169-177
- Bull, W. B. (1964) Alluvial fans. *J. Geol. Edu.*, v. 16, pp. 101-106.
- Bull, W. B. (1984) Tectonic Geomorphology. *J. Geol. Edu.*, v. 32, pp. 310-324.
- Bull, W. B. (1991) Geomorphic responses to climate change. Oxford University Press, New York.
- Bull, W. B. (1978) Geomorphic tectonic activity classes of the south front of the San Gabriel Mountains, CA.-Unpubl. Final Rep., U.S. Geol. Surv., Contract No.,14-08-0001-g-394: p. 59.
- Bull, W. B. and McFadden, L. M. (1977) Tectonic geomorphology north and south of Garlock Fault, California. In: D.O. Doehring (ED.), *Geomorphology in Arid regions*. Eighth Annual Geomorphology Symp. SUNY, Binghampton, pp. 127-137.
- Burbank, D.W. (1992) Causes of recent Himalayan uplift deduced from deposited patterns in the Ganges basin, *Nature*, v. 357, pp. 680-683.

- Burke, K. and Dewey, J. F. (1973) Plume generated triple junctions: key indicators in applying plate tectonics to old rocks. *J. Geol.*, v. 81, pp. 406-433.
- Burnett, A. W. and Schumm, S. A. (1983) Alluvial-river response to neotectonic deformation in Louisiana and Mississippi, *Science*, v. 222, pp. 49-50.
- Centamore, E., Ciccacci, S., Monte Del, M., Fredi, P. and Lupia Palmieri, E. (1996) Morphological and morphometric approach to the study of the structural arrangement of northeastern Abruzzo (central Italy). *Geomorphology*, v. 16, pp. 127-137.
- Chadha, R. K. (1992) Geological contacts, thermal springs and earthquakes in Peninsular India. *Tectonophysics*, v. 213, pp. 367-374.
- Chamley, H. (1979a) Les successions agrileuses de l'Atlantique Nord, echo des changements Mesozoiques et Cenozoiques de l'environnement. Exemple du basin de Cap Vert. *C.R. Acad. Sci. D*, v. 289, pp. 769-772.
- Chamley, H. (1979b) North Atlantic clay sedimentation and palaeoenvironment since the Late Jurasic. In: M. Talwani, W.Hay, and W.B. Ryan (Ed.) *Deep Drilling Results in the Atlantic Ocean: Continental Margins and palaeoenvironment*. Maurice Ewing Ser. 3, Am. Geophys. Union, pp. 342-361.
- Chamley, H. (1980) Clay sedimentation and palaeoenvironment in the Shikoku Basin since the middle Eocene (DSDP leg 58, Northern Philippine Sea). In G. deVries Klien, K. Kobayashi, et al., *Initial Reports DSDP*, 58. U.S. Gov. Print. Off. Washington, DC, pp. 669-678.
- Chamyal, L. S. and Merh, S. S. (1992) Sequence stratigraphy of the surface Quaternary deposits in the semi arid basin of Gujarat. *Man and Environment*, V.17, pp. 33-40.
- Chamyal, L. S. and Merh, S. S. (1995) The Quaternary formations of Gujarat. *Mem. Geol. Soc. Ind.*, v. 32, pp. 246-257.
- Chamyal, L. S., Sharma, B., Merh, S. S. and Karami, H. (1994) Significance of bank material at Tilakwada in lower Narmada Valley. *Current Science*, v. 66, pp. 306-307.
- Chamyal, L. S., Khadkikar, A. S., Malik, J. N. and Maurya, D. M. (1997) Sedimentology of the Narmada Alluvial Fan, Western India. *Sed. Geol.*, v. 109, pp. 263-279.
- Chamyal, L. S., Maurya, D. M., Bhanadari, S., and Raj, R. (2002) Late Quaternary geomorphic evolution of the Lower Narmada Valley, Western India: Implications for neotectonic activity along the Narmada-Son Fault. *Geomorphology*, v. 46 (3-4), pp. 177-202.
- Chamyal, L. S., Maurya, D. M., and Raj, R. (2003) Fluvial systems of the drylands of western India: a sysnthesis of Late Quaternary environmental and tectonic changes. *J. Quaternary International*, v. 104, pp. 69-86.
- Chandra, U. (1977) Earthquakes of Peninsular India: A seismotectonic study. *Bull. Seis. Soc. Amer.*, v. 65, pp. 1387-1413.

- Chappel, J., and Shackleton, N. J. (1986) Oxygen isotopes and sea level. *Nature*, v. 324, pp. 137-140.
- Cheetham, G. H. (1980) Late Quaternary palaeohydrology: the Kennet valley case-study. In Jones DKC(ed), *The shaping of southern England. (Inst. Br geogr Spec Publ. No.11)*. Academic press, London, New York, pp. 203-223.
- Choubey, V. D. (1971) Narmada-Son lineament, India. *Nature*, v. 232, pp. 38-40.
- Clemens, S., Prell, W., Murray, D., Shimmield, G. and Weedon, G. (1991) Forcing mechanisms of the Indian Ocean monsoon. *Nature*, v. 368, pp. 533-536.
- Coleman, J. M. (1969) Brahmaputra River: Channel Processes and sedimentation. *Sed. Geol.*, v. 3, pp. 129-239.
- Coulombe, C. E., Wilding, C. P. and Dixon, J. B. (1996) Overview of vertisols: Characteristics and impacts on society. *Adv. Agronomy*, v. 57, pp. 289-375.
- Crawford, A. R. (1978) Narmada-Son Lineament of India traced into Madagascar. *J. Geol. Soc. of India*, v. 19, pp. 144-153
- Darymple, R. W., Knight, R. J., Zaitlin, B. A. and Middleton, G. V. (1990) Dynamics and facies model of a macrotidal sand-bar complex, Cobiquid Bay-Salmon river estuary (Bay of Fundy). *J. Sedimentology*, v. 37, pp. 577-612.
- Darymple, R. W., Zaitlin, B. A. and Boyd, R. (1992) Estuarine facies models conceptual basis and stratigraphic implications. *J. Sed. Petrol.*, v. 62, pp. 1130-1146.
- DeCelles, P. G., Tolson, R. B., Graham, S. A., Ingersoll, R. V., White, J., Schmidt, C. J., Rice, R., Moxon, I., Lemke, L., Follo, J. W., Edwards, M. F., Cavazza, W., Caldwell, M. and Bargar, E. (1987) Laramide thrust-generated alluvial-fan sedimentation, Sphinx Conglomerate, southwestern Montana: American Association of Petroleum Geologists Bulletin, v. 71, pp. 135-155.
- DeCelles, P. G., Gray, M. B., Ridgway, K. D., Cole, R. B., Pivnik, D. A., Pequera, N. and Srivastava, P. (1991) Controls on syn-orogenic alluvial fan architecture, Beartooth Conglomerate (Palaeocene), Wyoming and Montana. *Sedimentology*, v. 38, pp. 567-590.
- Dilli, K. and Paint, R. K. (1994) Clay minerals as indicators of the provenance and palaeoclimatic record of the Kashmir Loess. *J. Geol. Soc. of Ind.*, v. 44, pp. 563-574.
- Dorn, R. I., DeNiro, M. J. and Ajie, H. O. (1987) Isotopic evidence of climatic influence of alluvial-fan development in Death valley, California, *Geology*, v. 15, pp. 108-110.
- Dudal, R. and Eswaran, H. (1988) Distribution, properties and classification of Vertisols. In: *Vertisols: Their distribution, properties, classification and management.* (Ed. Wilding, L.P. and Puentes) Tech. Mono. No. 18., Texas A & M Printing Center, pp. 1-22.
- Dumont, J. F. (1996) Neotectonics of Subandes-Brazilian craton boundary using geomorphological data: the Maranou and Ben basin. *Tectonophysics*, v. 257, pp. 137-151.

- Duplessy, J. C., (1982) Glacial to interglacial contrast in the northern Indian Ocean. *Nature*, v. 295, pp. 494-498.
- Dury, G. H. (1970) General theory of meandering valleys and underfit streams. In: *Rivers and River terraces* (Ed. Dury, G. H.). Praeger Publishers, N. Y., pp. 264-275.
- Elliott, T. (1974) Interdistributary bay sequences and their genesis. *Sedimentology*, v. 21, pp. 611-621.
- Ely, L. L.; Yehouda, E., Baker, V. R., Kale, V. S. And Mishra, S. (1996) Changes in the magnitude and frequency of late Holocene monsoon floods on the Narmada river, central India. *Geol. Soc. Am. Bull.*, v. 108, pp. 1134-1148.
- Ethridge, F. G. and Schumm, S. A. (1978) Reconstructing palaeochannel morphologic and flow characteristics – methodology, limitations and assessment. *Mem. Can. Soc. Pet. Geol.*, v. 5, pp. 703-721.
- Evans, J. E. (1991) Facies relationship, alluvial architecture, and paleohydrology of a Paleogene, humid-tropical alluvial-fan system: Chumstick Formation, Washington State, USA. *Jour. Sed. Petrol.* v. 61, pp. 732-755.
- Farrell, K. M. (1987) Sedimentology and facies architecture of overbank deposits of the Mississippi River, False River region, Louisiana. In: Ethridge, F.G., Flores, R. M. and Harvey, M.D. (Eds.), *Recent developments in Fluvial Sedimentology: SEPM Special Publication*, v. 39, pp. 111-120.
- Foote, R. B. (1898) *The Geology of the Baroda State*. Baroda State Press, 122p.
- Fuller, I. C., Macklin, M. G., Lewin, J., Passmore, D. G. and Wintle, A. G. (1998) River response to high frequency climate oscillations in southern Europe over the past 200 k.y. *Geology*, v. 26, pp. 275-278.
- Gardner, T. W. (1983) Palaeohydrology and palaeomorphology of a Carboniferous, meandering fluvial sandstone. *Jour. Sediment. Petrol.*, v. 53(3), pp. 991-1005.
- Gordon, I. and Heller, P. L. (1993) Evaluating major controls on basinal stratigraphy, Pine valley, Nevada: implications for syntectonic deposition. *Geol. Soc. Am. Bull.*, v. 105, pp. 47-55.
- Gradusov, B. P. (1974) A tentative study of clay mineral distribution in soils of the world. *Gerderma*, v. 12, pp. 49-55.
- Gupta, A. (1995) Fluvial geomorphology and Indian rivers. *Mem. Geol. Soc. of Ind.*, v. 32, pp. 52-60.
- Gupta, A., Kale, V. S. and Rajaguru S. N. (1999) The Narmada River, India, through space and time. Varieties of fluvial form, (Ed.) Miller, A. J. and Gupta A. pp. 113-143.
- Gupta, H. K., Indra Mohan and Hari Narain, (1972) The Broach earthquake of March 23, 1970. *Bull. Seis. Soc. Am.*, v. 62, pp. 47-61.

- Gupta, H. K., Chadha, R. K. , Rao, M. N., Narayna, B. L., Mandal, P., Ravikumar, M. and Kumar, N. (1997) The Jabalpur earthquake of May 22, 1997. Jour. Geol. Soc. India, v. 50. pp. 85-91.
- Gupta, M. L. (1991) Geothermal energy and its prospects in India. In: Moraes, C.P. (ed.) Power Directory 1991. Astra Publ., Bombay, India, pp. 157-165.
- Geological Survey of India (1978) Geological and mineralogical atlas of India. Sheet No. 24.
- Hagen, E. S., Schuster, M. W. and Furlong, K. P. (1985) Tectonic loading and subsidence of intermontane basins: Wyoming foreland province. Geology, v. 13, pp.585-588.
- Harris, P. T. (1988) Large-scale bedforms as indicators of mutually evasive sand transport and the sequential infilling of wide-mouthed estuaries. Sediment. Geol., v. 57, pp. 273-298.
- Harvey, A. M. (1984) Debris flow and fluvial deposits in Spanish Quaternary alluvial fans: implications for fan morphology. In: E.H., Koster, and R. J., Steel, (Eds) Sedimentology of gravels and conglomerate. Can. Soc. Pet. Geol. Mem. No. 10, pp. 123-132.
- Harvey, A. M. (1988) Controls on alluvial fan development, the alluvial fans of Sierra de Carrascoy, Murcia, Spain. Catena Supplement, A. 13, pp. 123-137.
- Hashimi, N. H., Nigam, R., Nair, .R. R. and Rajagopalan, G. (1995) Holocene sea level fluctuations on western Indian continental margin: An update. Jour. Geol. Soc. India, v. 46, pp. 157-162.
- Hegde, K. T. M. and Switsur, S. R. (1973) Radiocarbon dates on the buried soil in the lower Narmada valley. Current Science, v. 42(17), pp. 607-609.
- Heward, A. B. (1978) Alluvial fan sequence and megasequence models: with examples from Westphalian D-Stephanian B-coalfields, Northern Spain. In : A. D. Miall (Editor), Fluvial sedimentology, Can. Soc. Petrol. Geol. Mem., v. 5, pp. 669-702.
- Hooke, R. L. (1968) Steady state relationship on arid region alluvial fans in closed basins. Amer. Jour. Sci., v. 266, pp. 609-629.
- Hopkins, J. C. (1985) Channel-Fill deposits formed by aggradation in deeply scoured, superimposed distributaries of the Lower Kootenai Formation(Cretaceous). Jour. Sed. Petr., v. 55 (1), pp. 42-52.
- Horton, R. E. (1945) Erosional development of streams and their drainage basins, hydrophysical approach to quantitative morphology. Geol. Soc. Amm., v. 83, pp. 1649-1670.
- Hubert, J. F. and Filipov, A. J. (1989) Debris-flow deposits in alluvial fans on the west flank of the white mountains, Owens valley, California. J. Sediment. Geol., v. 61, pp. 177-205.
- Hutington, E. (1907) Some characteristics of the glacial period in non-glaciated regions. Geol. Soc. of Am. Bull., v. 18, pp. 351-388.
- Jackson, R. G. (1978) Preliminary evaluation of lithofacies models for meandering alluvial streams. In: Miall, A. D. (Ed.), Fluvial sedimentology: Mem. Can. Soc. Petrol. Geol., v. 5, pp. 543-576.

- Jain, S. C., Nair, K. K. K. and Yedekar, D. B. (1995) Geology of the Son-Narmada-Tapi lineament zone in central India. *Geoscientific studies of the Son-Narmada-Tapi lineamnet zone*. Project crumansonata. Geol. Surv. Ind. Spl. Publ., v. 10, pp. 1-154.
- Jain, M., Woodcock, N. H. and Tandon, S. K. (1998) Neotectonics of western India: evidence from deformed fluvial sequences, Mahi River, Gujarat. *J. Geol. Soc. London*, v. 155, pp. 897-901.
- Jain, M., Tandon, S. K., Bhatt, S. C., Singhvi, A. K. and Mishra, S. (1999) Alluvial and aeolian sequences along the river Luni, Barmer district: Physical stratigraphy and feasibility of Luminescence chronology methods. *Geol. Soc. Ind. Mem.*, v. 42, pp. 273-295.
- Johnson, A. M. (1970) *Physical processes in Geology*. San Francisco: Freeman Cooper and Co., 577 p.
- Jones, S. J., Frostick, L. E., and Tomothy, R. A. (1999) Climatic and tectonic controls on fluvial incision and aggradation in the Spanish Pyrenees. *J. Geol. Soc. London*, v. 56, pp. 761-769.
- Jordan, T. E. (1981) Thrust loads and foreland basin evolution, Cretaceous, western United States. *Bull. Am. Assoc. Petrol. Geol.*, v. 15, pp. 2500-2520.
- Juyal, N., Pant, R. K., Bhushan, R. and Somayajulu, B. L. K. (1995) Radiometric dating of the Late Quaternary sea-levels of the Saurashtra coast, western India: an experiment with oyster and clam shells. *Geological Society India Memoir*, v. 32, pp. 372-379.
- Juyal, N., Raj, R., Maurya, D. M., Chamyal, L. S. and Singhvi, A.K. (2000) Chronology of Late Pleistocene environmental changes in the Lower Mahi basin, Western India. *Jour. Quaternary Sci.*, v. 15, pp. 501-508.
- Juyal, N., Chamyal, L. S., Bhandari, S., Maurya, D. M. and Singhvi, A. K. (in press) Environmental changes during Late Pleistocene in the Orsang River basin, Western India. *Mem. Geol. Soc. of India*.
- Kaila, K. L. (1986) Tectonic framework of Narmada-Son lineament: a continental rift system in central India from deep seismic sounding. In: *Reflection seismology: a global perspective*. M. Barazangi and L. D. Brown (ed), Geodynamics series, 13: AGU Washington D. C., U.S., pp. 133-150.
- Kaila, K. L. (1988) Mapping the thickness of Deccan Trap flow in India from DSS studies and inferences about a hidden Mesozoic basin in the Narmada-Tapi region. In: *Deccan flood basalts*, K. V. Subbarao (ed), *Mem. Geol. Soc. Ind.*, no. 10, pp. 91-116.
- Kaila, K. L., Krishna, V. G. and Mall, D. (1981) Crustal structure along Mehmabad-Billimora profile in the Cambay basin, India from deep seismic soundings. *Tectonophysics*, v. 76, pp. 99-130.
- Kailasam, L. N., Murthy, B.G.K. and Chayanulu, A.V.S.R., (1972) Regional gravity studies of the Deccan Trap areas of peninsular India. *Curr. Sci.*, v. 41, pp. 403-407.
- Kale, V.S. (1999) Long-period Fluctuations in Monsoon Floods in the Deccan Peninsula, India. *Jour. Geol. Soc. Ind.*, v. 53, pp. 5-15.

- Kale, V. S. and Rajaguru, S. N. (1987) Late Quaternary alluvial history of the North Western Deccan Upland Region. *Nature*, v. 325, pp. 612-614.
- Kale, V. S., Mishra, S., Baker, V. R., Rajaguru, S. N., Enzel, Y. and Ely, L. (1993) Prehistoric flood deposits on the Choral River, Central Narmada Basin, India. *Current Science*, v. 65, pp. 877-878.
- Kale, V. S., Ely, L. L., Enzel, Y. and Baker, V. R. (1994) Geomorphic and hydrologic aspects of monsoon floods on the Narmada and Tapi Rivers in central India. *Geomorphology*, v.10, pp.157-168.
- Kale, V. S., Baker, V. R. and Mishra, S. (1996) Multi-channel patterns of bed rock rivers: An example from the Central Narmada Basin, India. *Catena*, v. 26, pp. 85-96.
- Kale, V. S., Mishra, S., and Baker, V. R. (1997) A 2000-Year palaeoflood record from Sakarghat on Narmada, Central India. *Jour. Geol. Soc. Ind.*, v. 50, pp.283-288.
- Kale, V. S., Mishra, S. and Baker, V. R. (2003) Sedimentary records of palaeofloods in the bedrock gorges of the Tapi and Narmada rivers, central India. *Current Science*, v. 84, pp. 1072-1079.
- Kar, A., Singhvi, A. K., Rajaguru, S. N., Juyal, N., Thomas, J. V., Banerjee, D. and Dhir, R. P. (2001) Reconstruction of Late Quaternary environment of the lower Luni plain, Thar desert, India. *j. Quat. Sci.*, v. 16, pp. 61-58.
- Kershaw, A. P., and Nanson, G. C. (1993) The last full glacial cycle in the Australian region. *Global and Planetary Changes* v. 7, pp. 1-9.
- Khadkikar, A. S., Chamyal, L. S. and Ramesh, R. (2000) The character and genesis of Calcrete in Late Quaternary alluvial deposits, Gujarat, Western India, and its bearing on the interpretation of Ancient Climates. *Palaeogeog. Palaeoclimat. Palaeoecol.*, v. 162, pp. 239-261.
- Khattri, K. N., Rogers, A. M., Perkins, D. M. and Algermissen, S. T. (1984) A seismic hazard map of India and adjacent area. *Tectonophysics*, v. 108, pp. 93-134.
- Kotlia, B. S., Sharma, C., Bhalla, M. S., Rajagopalan, G., Subrahmanyam, K., Bhattacharyya, and Valdiya, K. S. (2000) Palaeoclimatic conditions in the late Pleistocene Wadda Lake, eastern Kumaun Himalaya (India). *Palaeogeog. Palaeoclimat. Palaeoecol.*, v. 162, pp. 105-118.
- Kraus M. J. and Middleton, L. T. (1987) Contrasting architecture of two alluvial suites in different structural settings. *The Society of Economic Palaeontologists and Mineralogists*, pp. 253-262.
- Kraus, M. J. and Aslan, A. (1993) Eocene hydromorphic palaeosols: significance for interpreting ancient floodplain processes. *Jour. Sed. Petr.*, v. 63, pp. 453-463.
- Kraus, M. J. (1992) Alluvial response to differential subsidence: sedimentological analysis aided by remote sensing. Willwood Formation (Eocene), Bighorn Basin, Wyoming, USA. *Sedimentology*, v. 39, pp. 455-470.

- Krishnamachary, T. (1972) Dykes around Dadiapada, Broach district, Gujarat. Bull. Volcanologique, v. 35, pp. 947-956.
- Kusumgar, S., Raj, R., Chamyal, L. S. and Yadav, M. G. (1998) Holocene palaeoenvironment in the Lower Mahi basin, Western India. Radiocarbon, v. 40, pp. 819-823.
- Lawson, D. E. (1981) Mobilization, movement and deposition of subaerial sediment flows, Matanuska glacier, Alaska. Jour. Sed. Geol., v. 90, pp. 279-300.
- Lean, J. and Rind, D. (1994) Solar variability: Implications for global change. EOS. Trans. Amm. Geophys. Union, v. 75, pp. 5-7.
- Lean, J., Skumanich, A., and White, D. (1992) Estimating the sun's radiative output during the Maunder minimum. Geophy. Res. Letters, v. 19, pp. 1591-1594.
- Macklin, M. G., Fuller, I. C., Lewin, J., Maas, G. S., Passmore, D. G., Rose, J., Woodward, J. C., Black, S., Hamlin, R. H. B. and Rowan, J. S. (2002) Correlation of fluvial sequences in the Mediterranean basin over the last 200 ka and their relationship to climate change. Quat. Sci. Rev. v. 21, pp. 1633-1641.
- Maddy, D., Bridgland, D.R. and Westway, R. (2001) Uplift- driven valley incision and climate controlled river terrace development in the Thames valley, U.K. Quat. Int., v. 79, pp. 23-36.
- Maher, B. A. and Thompson, R. (1995) Paleorainfall reconstructions from pedogenic magnetic susceptibility variations in the Chinese loess and palaeosols. Quat. Res., v. 44, pp. 383-391.
- Maizels, J. (1993) Lithofacies variation within sandur deposits: the role of run off regime, flow dynamics and sediment supply characteristics. Sediment. Geol., v. 85, pp. 299-325.
- Malik, J. N., Khadkikar, A. S. and Merh, S. S. (1999) Allogenic control on Late Quaternary Continental Sedimentation in the Mahi River basin, Western India. Jour. Geol Soc. of Ind., v. 53, pp. 299-314.
- Marriott, S. B. and Wright, V. P. (1993) Palaeosols as indicators of geomorphic stability in two Old Red Sandstone alluvial suites, south Wales. J. Geol. Soc. London, v. 150, pp. 1109-1120.
- Massop, G. D. and Flach, P. D. (1983) Deep Channel Sedimentation in the Lower Cretaceous McMurray Formation, Athabasca Oil Sands, Alberta. Sedimentology, v. 30, pp. 493-509.
- Mather, A. E. (1993) Basin inversion: some consequences for drainage evolution and alluvial architecture. Sedimentology, v. 40, pp. 1069-1089.
- Mathur, L. P., Rao, K. L. N. and Chaube, A. N. (1968) Tectonic framework of Cambay basin, India. Bull., O.N.G.C., v. 5, pp. 7-28.
- Maurya, D. M., Chamyal, L. S. and Merh, S. S. (1995) Tectonic evolution of the Central Gujarat plain, Western India. Current Science, v. 69, pp. 610-613.
- Maurya D. M., Malik, J. N., Raj, R. and Chamyal, L. S. (1997a) Soft sediment deformation in the Quaternary sediments of the Lower Mahi river basin, Western India. Current Science, V. 72, pp. 519-522.

- Maurya D. M., Malik, J. N., Raj, R. and Chamyal, L. S. (1997b) The Holocene valley fill terraces in the Lower Mahi valley, Western India. Current Science, v. 73, No.6, pp.539-542.
- Maurya, D.M., Raj, R. and Chamyal, L.S. (1998) Seismically induced deformational structures from Mid-Late Holocene terraces in the lower Mahi valley, Gujarat, J. Geol. Soc. Ind., v. 51, pp. 755-758.
- Maurya, D. M., Raj, R. and Chamyal, L. S. (2000) History of tectonic evolution of Gujarat alluvial plains, Western India, during Quaternary: a review. Jour. Geol. Soc. India, v. 55, pp. 343-366.
- Mayer, L. (1985) Tectonic geomorphology of escarpments and mountain fronts. In: R.E. Wallace (Ed.), Active Tectonic. Natl. Res. Counc., Washington DC, pp.125-135.
- McCabbe, P. J., (1977) Deep distributary channels and giant bedforms in the Upper Carboniferous of the Central Pennines, northern England. Sedimentology, v. 24, pp. 271-290.
- McPherson, J. G., Shanmugam, G. and Miola, R. J. (1987) Fan deltas and braid deltas : Varieties of coarse grained deltas. Geol. Soc. Am. Bull., v. 99, pp.331-340.
- Meckel, L. D. (1972) Anatomy of distributary channel-fill deposits in recent mud deltas (Abstract). Am. Asso. Petroleum Geologists Bull., v. 56, pp. 639.
- Merh, S. S. (1993) Neogene-Quaternary sequence in Gujarat : A review . Jour. Geol.Soc.Ind., v. 41, Mar.1993, pp. 259-276.
- Merh, S. S. (1995) Geology of Gujarat. Geol. Soc. India, Bangalore, 222 pp.
- Merh, S. S. and Chamyal, L. S. (1993) The Quaternary sediments of Gujarat. Curr.Sci., v. 64, pp. 823-827.
- Merh, S. S. and Chamyal, L. S. (1997) The Quaternary Geology of Gujarat Alluvial Plains. Proc. Indian Natn. Sci. Acad., v. 63, pp. 1-98.
- Miall, A. D. (1977) A review of the braided river depositional environment. Earth Sci. Rev., v. 13, pp. 1-62.
- Miall, A. D. (1978) Lithofacies types and vertical profile models. In: A. D. Miall (Editor) Fluvial sedimentology. Can. Soc. Pet. Geol. Mem., v.5, pp.597-604.
- Miall, A. D. (1985) Architectural element analysis: A new method of analysis applied to fluvial deposits. Earth Sci. Rev., v. 22, pp. 261-308.
- Miall, A. D. (1996) The Geology of Fluvial Deposits. Springer-Verlag, Berlin. 582pp.
- Mishra S., and Rajaguru, S. N. (2001) Late Quaternary palaeoclimates of Western India: A geoarchaeological approach. Mausam, v. 52, pp. 285-296.
- Muller, J. E. (1968) An introduction to the hydraulic and topographic sinuosity indices. Annals Ass. Am. Geog., v. 58, pp. 371-385.

- Murthy, M. V. N. (1981) Late Mesozoic- Early Tertiary volcanism in the trans-Deccan trap areas of the Indian shield: a synthesis. In: K. V. Subbarao and R. N. Sukeshwala(eds) Deccan Volcanism. Mem. Geol. Soc. Ind., v. 3, pp. 93-100.
- Nanson, G. C., Price, D. M. and Short, S. A. (1992) Wetting and drying of Australia over the past 300 ka. *Geology*, v. 20, pp. 791-794.
- Nemec, W. and Postma, G. (1993) Quaternary alluvial fans in southwestern Crete: sedimentation processes and geomorphic evolution, in Marzo, M. and Puigdefabregas, C., (Editors), Alluvial Sedimentation: International Association of Sedimentologists Special Publication, v. 17, pp. 235-276.
- Nigam, R. (1984) Living benthonic foraminifera in a tidal environment : Gulf of Khambhat (India). *Marine Geology*, v. 58, pp. 415-425.
- Nilsen, J.H. (1982) Alluvial fan deposits. In sandstone depositional environments, P. Scholle and D. Spearing (ed.), 49-86. Tulsa: American Association of Petroleum Geologists Memoir, v. 31, pp. 49-86.
- Ouchi, S. (1985) Response of alluvial rivers to slow active tectonic movement. *Geol. Soc. Am. Bull.*, v. 96, pp. 504-515.
- Overpeck, J., Anderson, D., Trumbore, S. and Prell, W. (1996) The southwest Indian monsoon over the last 18000 years. *Climate Dynamics*, v. 12, pp. 213-225.
- Pant, R. K. and Chamyal, L. S. (1990) Quaternary sedimentation pattern and terrain evolution in Mahi river basin, Gujarat, India. *Proc. Indian Nat. Sci. Acad.*, v. 56, pp. 501-511.
- Pant, R. K. and Juyal, N. (1993) Late Quaternary coastal instability and sea-level changes: new evidences from Saurashtra coast, Western India. *Zeit fur geomorphologie*, v. 37, pp. 29-40.
- Pattanayak, S. K. and Shrivastava, J. P. (1999) Petrography and major oxide geochemistry of basalts from the eastern Deccan Volcanic Province, India. *Deccan Volcanic Province. Mem. Geol. Soc. Ind.* No.43(1), pp. 219-233.
- Pierson, T. C. (1980) Erosion and deposition by debris flow at Mt. Thomas, north Canterbury, New- Zealand. *Sedimentology*, v. 5, pp. 227-247.
- Pierson, T. C. and Scott, K. M. (1985) Downstream dilution of a lahar: transition from debris flow to hyperconcentrated stream flow. *Water Resources Res.*, v. 21, pp. 1511-1524.
- Powar, K. B. (1980) Lineament fabric and dyke pattern in the western part of the Deccan Volcanic Province. *Geol. Soc. Ind.*, v. 3, pp. 45-57.
- Prell, W. L. (1984) Variations of monsoonal upwelling: a response to changing solar radiation. In. Hansen, J.E. Takahashi, T. (eds) Climate processes and climate sensitivity, *Geophysical monograph* v. 29, pp. 48-57. Washington, D.C. American Geophysical Union.
- Prell, W. L., Marvil, R. E. and Luther, M. E. (1990) Variability in upwelling fields in the north western Indian Ocean 2. data-model comparisonat 9000 years B.P. *Palaeoceanography*, v. 5(3), pp. 447-457.

- Quittmeyer, R. C. and Jacob, R. F. (1976) Historical and modern seismicity of Pakistan, Afghanistan, northwest India and southeastern Iran. *Bull. Seis. Soc. Am.*, v. 69, pp. 773-823.
- Raj R. and L.S.Chamyal (1998a) Microfauna from a Holocene valley fill terrace, Lower Mahi basin, Gujarat. *J. Palaeont. Soc. Ind.*, v. 43, pp. 55-67.
- Raj R., Maurya, D. M. and Chamyal, L. S. (1998b) Late Quaternary sea level changes in western India : Evidence from lower Mahi valley. *Curr. Sci.*, v. 74, pp. 910-914.
- Raj R., Maurya, D. M. and Chamyal, L. S. (1999a) Tectonic control on distribution and evolution of Ravines in the lower Mahi valley, Western India. *J. Geol. Soc. India*, v. 52, pp. 669-674.
- Raj R., Maurya, D. M. and Chamyal, L. S. (1999b) Tectonic Geomorphology of the Mahi river basin, Western India, *J. Geol. Soc. Ind.*, v. 54, pp. 387-398.
- Raj R., Bhandari, S. Maurya, D. M. and Chamyal, L. S. (2003) Geomorphic indicators of Active Tectonics in the Karjan River basin, Lower Narmada Valley, Western India. *J. Geol. Soc. of Ind.*, v. 62, pp. 739-752.
- Radhakrishna, B. P. and Ramakrishnan, M. (1998) Archean-Proterozoic boundary in India. *J. Geol. Soc. India*, v. 32, pp. 263-278.
- Rajaguru, S. N., Gupta, A., Kale, V. S., Ganjoo, R. K., Ely, L. L., Enzel, Y. and Baker, V. R. (1995) Channel form and processes of the flood-dominated Narmada River, India. *Earth Surf. Process. and Landforms*, v. 20, pp. 407-421.
- Rao, P. C., Veerayya, M., Thamban, M. and Wagle, B. G. (1996) Evidences of late Quaternary neotectonic activity and sea-level changes along the Western Continental margin of India. *Curr. Sci.*, v. 71, pp. 213-219.
- Rao, R. P. and Talukdar, S. N. (1980) Petroleum geology of Bombay high field, India. In. M.T. Halbouty (ed.) *Giant Oil and Fields of the decade 1968-1978*. Amar. Asso. Pet. Geol., Mem., v. 30, pp. 387-506.
- Ravishankar (1991) Thermal and crustal structure of 'SONATA'. A zone of mid-continental rifting in Indian shield. *J. Geol. Soc. India*, v. 37, pp. 211-220.
- Reddey, P. R., Murthy, P. R. K., Rao, I. B. P., Khare, P., Kesava Rao, G., Mall, D. M., Koteshwara Rao P., Raju, S., Sridher, V. and Reddy, M. S. (1995) Seismic reflection fabric pattern in Central India : Preliminary results. *Mem. Geol. Soc. Ind.*, v. 31, pp. 537-544.
- Rhea, S. (1993) Geomorphic observation of rivers in the Oregon Coast Range from a regional reconnaissance perspective. *Geomorphology*, v. 27, pp.135-150.
- Rhee, C. W., Ryang, W. H., and Chough, S. K. (1993) Contrasting development patterns of crevasse channel deposits in Cretaceous alluvial successions, Korea. *Sed. Geol.*, v. 85, pp. 401-410.
- Rind, D. and Overpeck, J. (1993) Hypothesized causes of decade-to-century-scale climate variability: climate model results. *Quaternary Science Reviews*, v. 12, pp. 357-374.

- Roy, T. K. (1990) Structural styles in southern Cambay basin and role of Narmada geofracture in the formation of giant hydrocarbon accumulation. Bull. O.N.G.C., v. 27, pp. 15-38.
- Royden, L., Selator, J. G. and Herzen, R. P. (1980) Continental margin subsidence and heat flow: important parameter in formation of petroleum hydrocarbons, AAPG Bull, v. 64, pp. 173-187.
- Ruhe, R. V. (1956) Geomorphic surfaces and nature of soils. Soil Science, v. 82, pp. 441-455.
- Rust, B. R. and Koster, E. H. (1984) Coarse alluvial deposits. In : R. G. Walker (Editor) Facies models. Geoscience Canada Reprint Series 1, pp. 53-69.
- Sankalia, H. D. (1946) Investigations into the prehistoric Archaeology of Baroda, Gujarat, Baroda State Press.
- Sareen, B. K., Tandon, S. K. and Bhola, A. M. (1993) Slope deviatory alignment, stream network and lineament orientation of the Sabarmati river system-neotectonic activity in the Mid to Late Quaternary. Current Science, v. 64, pp. 258-276.
- Schubert, C. (1982) Neotectonics of Bocono Fault, Western Venezuela, Tectonophysics, v. 85, pp. 205-220.
- Schultz, A. W. (1984) A subaerial debris flow deposition in the Upper Palaeozoic Cutler Formation, Western Colorado. Jour. Sediment. Petrol., v. 54, pp. 759-772.
- Schuster, M. W. and Steidtmann, J. R. (1987) Fluvial sandstone architecture and thrust induced subsidence, northern Green river, Wyoming. The Society of Economic Palaeontologists and Mineralogists, pp. 279-285.
- Seeber, L. and Gornitz, V. (1983) River profiles along Himalayan arc as indicators of active tectonics. Tectonophysics, v. 92, pp. 335-365.
- Singer, A. (1984) The palaeoclimatic interpretation of clay minerals in sediments – a review. Earth Science Reviews, v. 21, pp. 251-293.
- Singh, A. D. (1998) Late Quaternary oceanographic changes in the eastern Arabian sea: Evidence from planktonic foraminifera and Pteropods. Jour. Geol. Soc. Ind., v. 52, pp. 203-212.
- Singh G. and Murti, G. S. R. (1975) Charge distribution in Smectite developed from basalts in Madhys Pradesh. J. Ind. Soc. Soil Sci., v. 23, pp. 177-183.
- Singh, I. B. (1996) Geological evolution of Ganga Plain- An Overview. Jour. Palaeont. Soc. India v. 41, pp. 99-137.
- Singh, N., Pant, G. B., and Mulye, S. S. (1991) Distribution and long term features of the spatial variations of the moisture regions over India. Int. Jour. Climat., v. 11, pp. 413-427.
- Sinha, R., and Friend, P. F. (1999) Pedogenic alteration in the overbank sediments, North Bihar Plains, India. Jour. Geol. Soc. Ind., v. 53, pp. 163-171.

- Sirocko, F., Sarnthein, M., Lange, H and Eslenkenser, H. (1991) Atmospheric summer circulation and coastal upwelling in the Arabian Sea during the Holocene and last Glaciation. *Quat. Res.*, v.36, pp. 72-93.
- Sirocko, F., Sarnthein, M., Eslenkenser, H., Lange, H., Arnold, M. and Duplessy, J. C. (1993) Centuary-scale events in monsoonal climate over the past 24,000 years. *Nature*, v. 364, pp. 322-324.
- Sloss, L.L. (1991) The tectonic factor in sea level change: a countervailing view. *J. of Geophysical Research*, v. 96, pp. 6607-6617.
- Smith, D. G. (1983) Anastomosed fluvial deposits: modern examples from Western Canada. *Int. Assoc. Sedimentologists Spec. Publ.*, v. 6, pp.155-168.
- Smith, D. G. (1993) Vertebrate taphonomy of late Permian floodplain deposits in the southwestern Karoo basin of South Africa. *Palaios*, v. 8, pp. 45-67.
- Smith, G. A. (1986) Coarse grained non-marine volcanoclastic sediment: terminology and depositional process. *Geol. Soc. Amer. Bull.*, v. 97, pp. 1-10.
- Smith, R. H. M. (1990) Alluvial palaeosols and pedofacies sequences in the Permian lower Beaufort of the southwestern Karoo basin, South Africa. *J. Sed. Petrol.*, v. 60, pp. 258-276.
- Sridhar, V., Chamyal, L. S. and Merh, S. S. (1994) North Gujarat rivers: Remnants of a superfluvial system. *J. Geol. Soc. Ind.*, v. 44, pp. 427-434.
- Srivastava, P., Juyal, N., Singhvi, A. K., Wasson, R. J. and Bateman, M. D. (2001) Luminescence chronology of river adjustment and incision of Quaternary sediments in the alluvial plain of the Sabarmati river, north Gujarat, India. *Geomorphology*, v. 36, pp. 217-229.
- Steel, R. J., Machle, S., Nilsen, H., Roe, S. L. and Spinnagr, A. (1977) Coarsening-upward cycles in the alluvium of Hornelen basin (Devonian), Norway : Sedimentary response to tectonic events. *Geol. Soc. Am. Bull.*, v. 88, pp. 1124-1134.
- Subramanya, K. R. (1996) Active intraplate deformation in south India. *Tectonophysics*, v. 262, pp. 231-241.
- Tandon, S. K., Sareen, B. K., Someshwa Rao, M. and Singhvi, A. K. (1997) Aggradational history and luminescence chronology of Late Quaternary semi-arid sequences of the Sabarmati basin, Gujarat, Western India. *Paleogeog. Palaeoclimat. Palaeoecol.*, v. 128, pp. 339-357.
- Tiwari, H. C., Murty, A. S. N., Kumar, P. and Sridhar, A. R. (2001) A tectonic model of Narmada region. *Current Science*, v. 80, pp. 873-878.
- Todd, S. P. (1989) Steam-driven, high-density gravelly traction carpets: possible deposits in the Trabeg Conglomerate Formation, SW Ireland and some theoretical considerations of their origin. *Sedimentology*, v. 36, pp. 513-530.
- Valdiya, K. S. (2001) River response to continuing movements and the scarp development in central Sahyadri and adjoining coastal belts. *Jour. Geol. Soc. Ind.*, v. 57, pp. 13-30.

- Veldkamp A. and Van Den Berg, M. W. (1993) Three-dimensional modelling of Quaternary fluvial dynamics in clino-tectonic dependent system. A case study of Maas record (Maastricht, The Netherlands). *Global and Planetary change*, v. 8, pp. 203-218.
- Veldkamp, A and Vermeulen, S. E. J. W. (1989) River terrace formation, modelling and 3-D graphical simulation. *Earth Surface Processes and Landforms*, v. 14, pp. 641-654.
- Venkatesan, T. R., Pande, K. and Gopalan, K. (1986) $^{40}\text{Ar}/^{39}\text{Ar}$ dating of Deccan Basalts. *J. Geol. Soc. Ind.*, v. 27, pp. 102-109.
- Wainwright, G. J. (1964) The Pleistocene deposits of the Lower Narmada river and an early stone age industry from the River Chambal. *M.S. University Archaeology and Ancient History Series*, 7. Baroda.
- Walker, T. K., Waugh, B. and Crone, A. J. (1978) Diagenesis in first cycle desert alluvium of Cenozoic, southwestern United States and northwestern Mexico. *Bull. Geol. Soc. Am.*, v. 89, pp. 19-32.
- Wassan, R. J., Smith, G. I. and Agarwl, D. P. (1984) Late Quaternary sediments, minerals and inferred geochemical history of the Didwana lake, Thar desert, India. *Palaeogeography Palaeoclimatology Palaeoecology*, v. 4, pp. 345-372.
- Weaver, C. E. (1989) Clays, Muds and Shales. *Developments in Sedimentology* 44. Elsevier. pp. 819
- Wells, S. G. and Harvey, A. (1987) Sedimentologic and geomorphologic variations in storm-generated alluvial fans, Howgill Fells, northwest England. *Geol. Soc. Amer. Bull.* v. 98, pp. 182-198.
- Wells, S. G., Bullard, T. F., Menges, C. M., Drake, P. G., Karas, P. A. and Kelson, K. I. (1988) Regional variations in the tectonic geomorphology along a segmented convergent plate boundary, Pacific Coast of Costa Rica, *Geomorphology*, v. 1, pp. 239-265.
- West, W. D. (1962) A line of Narmada-Son valley, Mirzapur District. *Mem. Geol. Soc. Ind.*, v. 62, pp. 133-136.
- White, J. W. C., Ciais, P., Figge, R. A., Kenny, R. and Graf, V. (1994) High resolution record of atmospheric CO_2 content form carbon isotopes in peat. *Nature*, v. 367, pp. 153-156.
- Whiting, B. M., Karner, G. D. and Driscoll, N. W. (1994) Flexural and stratigraphic development of the west Indian continental margin. *J. Geophy. Res.*, v. 99, pp. 13791-13811.
- Widdowson, M. (1997) Tertiary palaeosurfaces of the SW Deccan western India: implications for passive margin uplift. In: Widdowson, M. (Ed.). *Recognition, Reconstruction and Palaeoenvironmental Interpretation*. *Geo. Soc. Lond. Spec. Publ.* No.120, pp. 221-248.
- Widdowson, M., and Mitchell, C. (1999) Large scale Stratigraphical, Structural and Geomorphological constraints for Earthquakes in the southern Deccan Traps, India: The case for denudationally-driven seismicity. *Mem. Geol. Soc. Ind.*, No.43, pp. 425-452.

- Wilding, L. P. and Tessier, D. (1988) Genesis and vertisols: Shrink – swell phenomena. In Vertisols: their distribution, properties, classification and management (Wilding, L. P and Puentes, R., Eds.) pp. 55-81. Tech. mono. No. 18. Texas A & M Printing centre, college station, TX.
- Williams, G. P. (1978) Bankfull discharge of rivers. Water Resource Research, v. 14, pp. 1141-1154.
- Williams, G. P. (1984) Palaeohydrologic equations for rivers. In: J. E. Costa and P.J. Fleisher, (Eds.) Developments and Applications of Geomorphology. Springer – Verlag, Berlin and New York, pp. 343-367.
- Williams, G. P. (1988) Palaeofluvial estimates from dimensions of former channels and meanders. In. V.R. Baker, R.C. Kochel and P.C. Patton, (Eds.) Flood Geomorphology, John Wiley and Sons. New York, pp. 321-334.
- Williams, M. A. J. and Clarke, M. F. (1984) Late Quaternary environments in north-central India. Nature, v. 308, pp. 633-635.
- Willis, B. J. and Behrensmeyer, A.K. (1994) Architecture of Miocene overbank deposits in Northern Pakistan. Jour. Sed. Petr., v. B64 (1), pp. 60-67.
- Woodroffe, C. D., Chappell, J., Thom, B. G, and Wallensky, E. (1989) Depositional model of a microtidal estuary and floodplain, South Alligator river, northern Australia, Sedimentology, v. 36, pp. 737-756.
- Wright, L. D., Coleman, J. M. and Thom, B. G. (1973) Processes of channel development in a high-tide-range environment: Cambridge Gulf-Ord River delta, Western Australia. J. of Geology, v. 81, pp. 15-41.
- Yan, Z. and Petit-Maire, N. (1994) The last 140 ka in the Afro-Asian arid/semi-arid transitional zone. Palaeogeog. Palaeoclimat. Palaeoecol., v. 110, pp. 217-233.
- Zeuner, F. E. (1950) Stone age and Pleistocene chronology in Gujarat. Deccan college Monograph series: 6, 48pp.
- Ziegler, P. (1983) Inverted basins in the Alpine foreland. In: Bally A.W. (ed), Seismic expression of structural styles-a picture and world atlas. Am. Assoc. Petrol. Geol., v. 15(3), pp. 3.3.3-3.3.12.
- Zieleinski, G. A., Mayewski, P. A., Meeker, L. D., Whitlow, S., Twickler, M. S., Morrison, M., Meese, D. A., Gow, A. J., and Alley, R. B. (1994) Records of volcanism since 7000 BC from the GISP2 Greenland core and implications for the volcano-climate system. Science, v. 264, pp. 948-952.

Papers published from the doctoral thesis

Bhandari S., Raj, R., Maurya, D. M. and Chamyal, L. S. (2001) Formation and Erosion of Holocene Alluvial fans along Narmada – Son Fault near Rajpipla in lower Narmada basin, western India. *Jour. Geol. Soc. Ind.*, v. 58, pp. 519-531.

Chamyal, L. S., Maurya, D. M., **Bhandari, S.**, and Raj, R. (2002) Late Quaternary geomorphic evolution of the Lower Narmada Valley, Western India: Implications for neotectonic activity along the Narmada-Son Fault. *Geomorphology*, v. 46 (3-4), pp. 177-202.

Raj R., **Bhandari, S.**, Maurya, D. M. and Chamyal, L. S. (2003) Geomorphic indicators of Active Tectonics in the Karjan River basin, Lower Narmada Valley, Western India. *J. Geol. Soc. of Ind.*, v. 62, pp. 739-752.

Bhandari, S., Maurya, D. M. and Chamyal, L. S. (2004) Late Pleistocene alluvial plain sedimentation in lower Narmada valley, Western India: Palaeoenvironmental implications. *Jour. of Asian Earth Sciences* (in press).

Other publications of the author

Maurya,D.M., **Bhandari, S.**, Thakkar, M.G. and Chamyal, L.S. (2003) Late Quaternary fluvial sequences of southern Mainland Kachchh, Western India. *Current Science*. v. 84, pp. 1056-1064.

Raj. R., Mulchandani, N., **Bhandari, S.**, Maurya, D. M. and Chamyal, L.S. (2003) Evidence of a Mid-Late Holocene seismic event from Dhadhar river basin, Gujarat alluvial plain, western India. *Current Science*. v. 85, pp. 812-815.

Juyal, N, Chamyal, L. S., **Bhandari, S.**, Maurya, D. M. and Singhvi, A. K. (In press) Environmental changes during Late Pleistocene in the Orsang River basin, Western India. *Mem. Geol. Soc. of India*.

Raj, R., Mulchandani, N., **Bhandari, S.**, Maurya D. M. and Chamyal, L. S. (In press) Channel shifting of a highly sinuous meandering river in alluvial plain, Vishwamitri River, Mainland Gujarat. *Current Science*.