

LIST OF FIGURES

	Page No.
Fig.1.1. A. Location map. B. Generalized geological map of the Narmada basin. Shaded area is lower Narmada basin	3
Fig.1.2. Map showing the major physiographic division, rivers and climatic zones of Gujarat. Shaded area is the lower Narmada basin (after Singh et al. 1991)	4
Fig.1.3. Isohyte map of Gujarat (after Singh et al. 1991)	4
Fig.1.4. Discharge in Narmada River at Garudeshwar gauged from 1991 to 2004 (Data for Garudeshwar Gauging site collected from Sardar Sarovar Nigam Ltd., Vadodara)	5
Fig.1. 5. Digital Elevation Model (DEM) of lower Narmada basin	5
Fig. 2.1. Geological map of lower Narmada basin (after GSI, 1975)	10
Fig 2.2. Geological map of lower Narmada basin (after Biswas, 1987)	11
Fig. 2.3. Map showing tectonic trends (after Biswas, 1987)	13
Fig. 2.4. A. Heat flow and hot spring (after Gupta et al. 1991) B. Seismic events in Western India (after Khattri et al. 1984)	14
Fig 2.5. DSS profile from Memdabad to Billimora showing basement configuration (after Kaila et al. 1981)	14
Fig. 2.6. A) Tectonic map of the lower Narmada basin (after Chamyal et al. 2002) Note the structural highs and reverse faults to the south of Narmada – Son Fault (structural data from Agarwal, 1986) B) Simplified cross-section across the NSF based on seismic reflection data showing the nature of the fault in the subsurface (after Roy, 1990)	16
Fig. 2.7. N-S cross section across Gujarat alluvial plains based on thickness data obtained from ONGC showing uneven Quaternary basement (after Maurya et al. 1995)	17
Fig 3.1. Drainage map of lower Narmada basin	20
Fig 3.2. IRS FCC image of the lower Narmada valley. 1. Alluvial Zone, 2. Tertiary Upland, 3. Trappean Upland	21
Fig. 3.3. Geomorphic map of the lower Narmada basin (after Chamyal et al. 2002). X-Y, X ₁ -Y ₁ and X ₂ -Y ₂ are the section lines of Fig. 3.4	22

Fig. 3.4. Topographic profiles showing the general physiography of the lower Narmada basin. Section lines are shown in Fig. 3.3	22
Fig. 3.5. Topographic profile along Karjan river basin showing the tilt blocks. Note the rugged topography and the subdued topography in the northern and southern parts respectively of each of each tilt blocks. Dashed line indicate the original position of the tilt block	23
Fig. 3.6. Field photograph showing southward tilting of trappean rocks at Khuta Amba	23
Fig.3.7. Geomorphic map of the Karjan River basin showing the various tectonic landforms. X-Y is the section line of Fig.3.1	24
Fig.3.8. Photomosaic of ENE-WSW trending ridge with intramontane valley in the foreground. At the base small colluvial fans have developed (1 km South of Ghantoli)	25
Fig.3.9. Colluvial Fan deposits of irregular shape and size (2 km north of Khamer)	25
Fig. 3.10. Intramontane valley developed at the base of parallel ridges near Ghantoli	26
Fig. 3.11. Photograph of a fluvial terrace in the proximal part of the Karjan river basin (Loc. Thava)	26
Fig. 3.12. Photograph of the Mohan Nadi cutting the basement rock besides the overlying terrace (Loc. Thava)	26
Fig. 3.13. Photograph showing cuestas in the background (Loc. Rajpipla)	27
Fig. 3.14. Upstream view of a deep gorge along a tributary stream of Karjan river in uplands (Loc. Near Moti Chikli)	27
Fig. 3.15. Maps showing ponding due to differential movements along faults (Raj et al. 2003) as seen in A- Karjan and Mohan Nadi, B- Karjan river and C- Tarav river	28
Fig. 3.16. Ponding in Karjan River at Thava. Note the ponding of the river and the flat topped incised terrace	28
Fig 3.17. Logitudinal profile of the Narmada, Karjan and Tarav Rivers. The places marked on the profile are the points of break or knickpoint	29
Fig. 3.18. Photograph of Waterfall in a tributary stream of Dhamankhadi (Loc. Rakaskundi)	29
Fig. 3.19. Ravine surface at Ranapur with Holocene terrace separated by palaeobank	30
Fig. 3.20. Photograph of the uplifted terrace at Gora. The upliftment is about 35 m	31
Fig. 3.21. Upstream view of the large entrenched meander in the alluvial zone near NSF (Loc. Tejpur)	31

Fig. 3.22. Flat topped Holocene estuarine terrace at Tavra	32
Fig. 3.23. Incised fluvial terrace exposing the S ₄ sediments (Loc. near Rajpardi)	32
Fig. 3.24. Estuarine terrace exposed near Bharuch	33
Fig. 4.1. A. Map showing locations of the sites studied. B. Vertical lithologs of the exposed sediments along the Narmada River	35
Fig. 4.2. Photograph of proximal gravel facies of alluvial fan 1 at Garudeshwar	36
Fig. 4.3. Photograph of gravelly sequence of Alluvial Fan 1 exposed at Rampura	36
Fig. 4.4. Incised cliff section showing the sediment succession of alluvial fan 1 at Tilakwada	37
Fig. 4.5. View of right bank cliff at Tilakwada comprising sediments of alluvial fan 1	37
Fig. 4.6. Close view of the cliff section exposed at Chandod	38
Fig. 4.7. Sediment succession at Nanderiya comprising basal clays overlain by alluvial fan facies and topped by alluvial plain facies	39
Fig. 4.8. Basal marine clays exposed at Nanderiya showing distinct vertic characteristics and extensive fracturing	39
Fig. 4.9. Downstream view of the right bank cliff at Maletha showing the coarse fan facies overlain by finer alluvial plain facies	40
Fig. 4.10. Photograph showing the upper part of the exposed Late Pleistocene sequence (loc. 1 km downstream of Ambali) 1. horizontally stratified overbank sands, 2. palaeosol, 3. thinly stratified sands and silts	40
Fig. 4.11. Sedimentary succession exposed at Patan revealing thick deposits of alluvial fan at the base overlain by sands and silts of alluvial plain environment on the top	41
Fig. 4.12. Field photograph showing the lithosection exposed at Kanjetha	41
Fig. 4.13. Photograph showing stratified silts and sands of alluvial plain facies exposed at Ranapur	42
Fig. 4.14. Field photograph of alluvial plain deposits exposed at Nikora. See the distinct pedogenised horizon overlying the massive sands	42
Fig. 4.15. Photograph of Late Pleistocene deposits exposed at Nava Tavra revealing clay deposits at the base overlaid by horizontally stratified sands	43

Fig. 4.16. A. Geomorphological map of the S ₃ surface (after Bhandari et al. 2001). B. N-S cross section along X-X' showing the inter-relationship between the various geomorphic surfaces around Rajpipla. Legend is same as in A	44
Fig. 4.17. A. Alluvial fans along the Narmada-Son Fault (NSF). Fans are numbered 1 to 5. B. Map showing detailed geomorphic characteristics of Fan3	45
Fig. 4.18. Lithologs and lithofacies assemblages of the sections measured in alluvial fans. Locations are shown in Fig. 4.17	46
Fig. 4.19. Field photograph of fan sediments exposed at Bhandra	47
Fig. 4.20. Close up of clast supported gravels exposed at Junawad	47
Fig. 4.21. Field photograph of Shamariya section (9m) exposing medial fan deposits	48
Fig. 4.22. Photograph of medial fan deposits at Phulwadi showing distinct layers of matrix supported gravels and horizontally stratified sand. (height of the person is 1.75 m)	48
Fig. 4.23. Photograph of Distal fan deposits exposed at Rawal Moti	49
Fig. 4.24. Field photograph of distal fan deposits exposed at Karotha. Note the finer nature of the sediments and the gravels deposited are well rounded	50
Fig. 4.25. Field photograph of Rajpipla section showing Late Pleistocene sediments capped by Early Holocene Fan deposits	51
Fig. 4.26. Lithologs of S ₄ surface exposed in the lower Narmada basin	52
Fig. 4.27. Field photograph of Holocene estuarine section at Uchediya (8.5 m) exposing alternate layers of organic clays and sands	52
Fig. 4.28. Field photograph of Holocene section exposed at Tavra (6 m). thick clay deposits with thin bands of intervening sand layers are exposed	53
Fig. 4.29. Holocene estuarine terrace section exposed at Bharuch	53
Fig. 4.30. Composite lithostratigraphic profile of the exposed Late Pleistocene sediments in the lower Narmada basin (after Bhandari et al. 2004)	54
Fig. 5.1. Thick fluvial deposits overlying the basal marine clays at Nanderiya	57
Fig. 5.2. Photograph showing clast supported gravels (Gc) of Alluvial Fan 2 at Maletha (length of hammer is 30cm)	58
Fig. 5.3. Alluvial Fan 2 sediments (Gms, Gp, Gh) overlain by alluvial plain facies (Sh) at Nanderiya (height of the person is 1.65m)	58
Fig. 5.4. Gp facies at Nanderiya overlain by Gms and Gh facies	59

Fig. 5.5. Gt facies at Nanderiya of Alluvial fan 2 sediments	59
Fig. 5.6. Planar cross stratified sand (Sp) at Ambali overlain by Sh facies (length of the hammer is 30cm)	60
Fig. 5.7. Photomosaic of laterally extensive cliff exposure at Chandod. All units show erosional bases. Height of the person is 1.78m. 1. distal alluvial fan facies, 2. large channel fill sands, 3. gaint epsilon cross bed strata, 4. horizontally stratified overbank strata, 5. palaeosol, 6. thinly stratified sands and silts. The sketch figure(not to scale) shows the vertical relation of the sediments at Chandod. Note the vertical stacking of the sediments and the two generations of channel fills	62
Fig. 5.8. Large channel fill structure exposed 1km downstream of Kanjetha. Note the prominent concave up geometry of the fill and the internal laminations. Tracing overlay brings out the major features of the channel fill	63
Fig.5.9. Partially eroded channel fill structure at the southern tip of the laterally extensive cliff exposure at Chandod. Tracing overlay of the photograph shows the major characteristics of the channel fill as seen in the outcrop	64
Fig.5.10. Field photograph of the Epsilon cross bedded strata 2 km downstream of Kanjetha which overlies channel fill structure. Tracing overlay brings out the Epsilon cross bedding	66
Fig. 5.11. Field photograph of the horizontally stratified overbank sands exposed near Ranapur. Height of the person in the photo is 1.65 m	67
Fig. 5.12. Field photograph of undulatory bedded sand and silt deposits at Kanjetha. Height of the person in the photo is 1.65 m	68
Fig. 5.13. Field photograph showing close view of the backswamp deposits at Nava Tavra. Note the abundant root structures in the form of rhizocretions	69
Fig. 5.14. Closeup view of the Gc facies at Mota Amba	72
Fig. 5.15. Close view of the Gms facies at Vavdi	72
Fig. 5.16. Sp facies overlain by Gp facies and underlain by Gh facies	73
Fig. 5.17. Four aggradational phases of Gms facies with intervening Sh facies at Phulwadi	73
Fig.5.18. Schematic longitudinal cross-section summarising geological setting and their facies associations (as demonstrated by Blair and McPherson, 1994)	75
Fig. 5.19. Schematic cross-section showing evolutionary stages and their relationship to dominant sedimentary processes (as demonstrated by Blair and McPherson, 1994)	76

Fig. 5.20. Depositional model for the alluvial fans near Rajpipla depicting various facies and morphological features	76
Fig. 5.21. Estuarine tidal valley fill terrace at Uchediya S ₄ surface	77
Fig. 6.1. Major faults and lineaments in the Karjan basin with rosette showing the dominant lineament trends	81
Fig. 6.2. Drainage map of Karjan basin with tributary sub-basins. Note the control of NNW-SSE trending lineaments on drainage	83
Fig. 6.3. Longitudinal Profile of Karjan River showing prominent breaks at Ambadi, Panuda, Moti Daberi and Bej	83
Fig. 6.4. Longitudinal Profile of Tarav River showing prominent breaks at Nimbapati, Mathavali, Khudali, Ghantoli and Bej	84
Fig. 6.5. Plot of Gradient Index of Karjan River	84
Fig. 6.6. Valley floor ratios of Karjan River	85
Fig. 6.7. Sinuosity values of Karjan River computed for segment indicated	86
Fig. 6.8. Map showing location of mountain fronts analysed and the values of Mountain Front Sinuosity (the tectonic activity class is after Bull & McFadden, 1977)	87
Fig. 6.9. Ravines of the lower Narmada basin with rosettes showing dominant ravine trends	88
Fig. 6.10. Intraformational open folds in the overbank deposits at Ambali (height of the person is 1.65m)	91
Fig. 6.11. Small-scale low angle thrust fault in the overbank deposits at Nava Tavra (length of the pencil is 12 cm.)	92
Fig. 6.12. Horizontal as well as vertical jointing seen in the Late Pleistocene sediments with deposition of calcretes along joints. See the minor deflection in the joints	92
Fig. 6.13. Syndepositional slump structure in the alluvial plain sediments at Patan (height of the person is 1.65m)	93
Fig. 6.14. Topographic profiles showing postdepositional tilting of the S ₁ surface consisting of Late Pleistocene sediments	93
Fig. 6.15. Load structures formed due to seismic loading in a fluvial Holocene terrace in the Orsang valley at Anklawadi	95

Fig. 6.16. Photograph showing soft sediment deformation structures seen in sand and clay in Amrawati River at Dhadal	95
Fig. 7.1. A. Litholog showing the exposed sedimentary sequence at Nanderiya. B. Detailed log of the basal clays showing the sample locations for the clay mineralogy and palynological analysis. C to F. are the percentage of Smectite, Illite, Kaolinite and Chlorite	97
Fig.7.2. Cross section across the Narmada basin with active and inactive floodplains and the palaeobanks	100
Table 7.1. Bankfull discharge calculated from the present channel and the early Holocene palaeobanks. The calculations have been done based on the methodology described by Williams (1978)	101
Fig. 7.3. Pollen diagram of Bharuch section. Also showing the environment during its deposition	105
Fig. 8.1. Physical stratigraphy of the exposed Late Pleistocene sediments (S_1 and S_2 surfaces) in the lower Narmada valley and the adjacent Mahi and Sabarmati valleys (after Chamyal et al. 2002). Note the correlatability of major aggradational phases. Dates are after Chamyal et al. (2002), Juyal et al. (2000), Tandon et al. (1997) and Allchin et al. (1978)	108