

BIBLIOGRAPHY

BIBLIOGRAPHY

- Adler, J.H. and A.S. Thomas. 1983. Occurrence of C-24 alkylated Δ^7 -sterols and Δ^5 sterols in the leaves of *Beta vulgaris*. Lipids, 18(3): 229-232.
- Airy Shaw. 1948. Flora Malesiana 4 : 27.
- Aguirre, S., Bruhn, J.G., Lundstrom, J. and Svensson, U. 1971. Cactaceae alkaloid A. Alkaloids of *Trichocereus* species and some other cacti lloydia, 34(2) : 183-37.
- Asarasingham, A.D., Bisset, N.C., Willard, A.H. and L.C. Wood. 1964. Phytochemical Survey of Malaya, Part III. Alkaloids and Saponins. Eco. Bot., 18(3) : 270.
- Andrews, R.S. and J.B. Pridham, 1966. Phytochemistry, 6 : 15-18.
- Annaev, Ch.M., Isamukhamedova A. and M.A. Abubakirov. 1994. Triterpene glycoside from *Salsola micranthera*. Thin. Layer. Sci. Technol., 0(1) : 727-32.
- Arakawa, Y., Y.Z. Asada, H. Ishida, M. Hideyukihihi and M. Isawa. 1982. Structure of new 2 isoflavones and 1 flavone from glass wort (*Salicornia europaea*). J. Fac. Agri. Hokkaido Univ., 61(1) : 1-12.
- Arisawa, Nobukominabe, M.R., Tokutaro, T. Takakuwa and T. Nakaoki. 1971. The components of flavonoids in *Chenopodium* genus plants I. Flavonoids of

- Chenopodium ambrosioides L. Yakugaku Zasshi, 91(5) : 522-24.
- Banerji, A. and M.S. Chadha. 1970. Insect moulting hormone from Achyranthus aspera. Phytochemistry, 9 : 1671.
- Banerji, A. and C.J. Chintalwar. 1971. Phytoecdysones and a new flavonols glycoside from Sesuvium portulacastrum Linn. Indian J. Chem., 9 : 1029-1030.
- Bate Smith, E.C., Harborne, J.B. and S.M. Devenport. 1966. Occurrence of azaleatin and caryatin in Luxyrrhia Nature, 212(5066) : 1065-66.
- Bate Smith, E.C. and T. Swain. 1966. The asperulosides and the aucubins - In Comparative Phytochemistry, Swain . T (ed) 159-174. Academic Press, London.
- Bate Smith, E.C. 1972. Chemistry and Phylogeny of the angiosperms. Nature, 236 : 353-354.
- Bathory, M., Toth, I., Swendrei, K., Rattai, M., Minker, L. and G. Blazso. 1984. Determination and isolation of ecdysteroids in native goosefoot species. Herba Hung, 23(1/2) : 131-42.
- Bech, T.D. 1968. On the flavonoids of red root (Amaranthus retroflexus). Farm. Zh., 20(5) : 33-41.
- Bech, T.D., Lastovska, E.A., Zubrytska, H.T., Artemenko, V.I. and M.M. Yuzvishyn. 1977. Flavonoids from plants of genus Amaranthus. Farm Zh. (Kiev), 32(4) : 90-91.

- Behnke, H.D. 1971. Sieve tube plastids of Magnoliidae and Ranunculidae in relation to Systematics. Taxon, 20: 723-730.
- Behnke, H.D. and Turner, B.L. 1971. Taxon, 20: 731.
- Behnke, H.D. 1972. Sieve-tube plastids in relation to angiosperm Systematics - An attempt towards a classification by ultrastructural analysis. Bot. Rev., 38 : 155-157.
- Behnke, H.D. 1974. Elektronenmikroskopische Untersuchungen and Siebröhren-plastiden und ihre Aussage über die Systematische Stellung Von Lophiocarpus. Bot. Jahrb. Syst., 94 : 114-119.
- Behnke, H.D. 1975. Die Siebelement-Plastiden der Caryophyllaceae. eine weitere spezifische Form der F-Type Plastiden bei Centrospermen. Bot. Jahrb. Syst., 95 : 327-333.
- Behnke, H.D. 1976. Ultrastructure of sieve element plastids in Caryophyllales (Centrospermae), Evidence for the Delimitation and classification of the order. Plant. Syst. Evol., 126: 31-54.
- Below, R.E., Leung, A.Y., McLaughlin, J.L. and A.G. Paul. 1968. Cactus alkaloids IV macromerene from Coryphantha runyonii. J. Pharm. Sci., 57(3) : 515-16.
- Bentham, G. and J.D. Hooker, 1865. Genera Plantarum Vol. 2, London.
- Bentham, G. and J.D. Hooker, 1860. Genera Plantarum, 3. London.
- Bessey, C., 1915. The phylogenetic taxonomy of flowering plants. Ann. Missouri Bot. Gard., 2 : 109-164.

- Behere B. and S.N. Patnaik. 1972. Cytotaxonomic studies in the family Amaranthaceae. Cytologia, 32 : 121-31.
- Blackwell, H. 1977. The subfamilies of Chenopodiaceae. Taxon, 26 : 395-97.
- Bogacheva, M.G., Leonids, ^{N.} ~~E.~~ Cogan, and N.A. Libzinov. 1972. Triterpenoid glycosides of Chenopodium ambrosioides. Khim. Prirod. Soedin (Tashk)., 8(3) : 395.
- Borkowski, A.A., M. Probst. 1966. Alkaloids of Salicornia herbaceae Linn. Pharmazie, 20 : 390-93.
- Braun, A. 1864. Übersicht des natürlichen Systems. In Ascherson, P. : Flora der Provinz Brandenburg, der Altmark und des Herzogthums Magdeburg, 12-67, Berlin : August Hirschwald
- Britton and Rose. 1873. In Willis . A Dictionary of the flowering plants and ferns (8th ed. 1973 revised by H.K. Airy Shaw) University press, Cambridge.
- Bruha, J. ^{J. &} Agurell G.A. and J.E. Lindgren. 1975. Cactaceae alkaloids XI. Phenethylamine alkaloids of Coryphantha species. Acta Pharm. Toxic. 12(2):199-204.
- Brutko, L.I., Massagetov P.S. and L.I. Utkin, 1968. Content of isomers of anabasine in different specimens of Anabasis aphylla. Russ. med. bot., 4(3): 354-36.
- Bukhariv, V.G. and S.P. Shcherbak. 1970. Triterpene glycosides of Herniaria glabra L. Khim. Prirod. Soedin (Tashk)., 6(3) : 307-11.

- C.
 Buschi, Carlos, A., and Alicia G. Comillo. 1982. Isorhamnetin
 5-O-robinobioside from Gomohrena martiana. J. Nat. Prod.
(Lloydia), 46(5) : 557-59.
- Buxbaum, R. 1961. Vorläufige Untersuchungen über Umlang,
 Systematische Stellung und Gliederung der Caryo-
 phyllales (Centrospermae). Botan. Biol. Pflanzen, 36:
 1-56.
- Chang, C.F. and Mabry, T.J. 1973. The constitution of the
 order Centrospermae : rRNA-DNA Hybridization studies
 among Betalain and Anthocyanin producing families.
Biochem. Syst. 1 : 185-190.
- Chang, C. Kizler, L. and T.J. Mabry, 1974. Biogenesis of
 betalamic acid. Phytochemistry, 13 : 2771-2775.
- Chumbalov, T.K. and V.B. Omurkamzinova, 1970. Flavonoids
 from Polygonum coriarium leaves. Khim. i. rir. Soedin
(Tashk), 6(3) : 366.
- Chumbalov, T.K., Mukhamed Yarova, R.M., I.S. Chanyshova and
 R.M. L'yasova, 1971. Flavonoids of Atraphaxis
pyrifolia and A. spinosa. II. Khim. i. rir. Soedin
(Tashk), 4 : 525-26.
- Chumbalov, T.K., Mukhamed Yarova, R.M. V.B. Omurkamzinova
 and I.S. Chanyshova 1975. 3-Rhamnoside of 7-O-methyl-
 gossypetin from Atraphaxis pyrifolia, Khim. i. rir.
Soedin (Tashk), 2 : 136-40.
- Chumbalov, T.K., Mukhamed Yarova, R.M., Omurkamzinova, V.B. and
 I.S. Chanyshova 1975. 3-Rhamnoside of 7-O-
 methylgossypetin from Atraphaxis pyrifolia. Khim.
i. rir. Soedin (Tashk) 2: 136-40.

- Clark, W.D., Brown, G.K. and R.L. Mays. 1980. Flower
Flavonoids of Opuntia subgenus cylindropuntia.
Phytochemistry, 19(9) : 2042-43.
- Crawford, L.J. 1975. Systematic relationships in the narrow-
leaved species of Chenopodium of western united
states. Brittonia, 27(3) : 279-88.
- Crawford, L.J. and A. Julian 1976. Seed protein in the
narrow leaved species of Chenopodium of western
United States : Taxonomic value and comparison with
distribution of flavonoid compounds. Am.J.Bot.,
63(3) : 302-308.
- Cronquist, A. 1965. The status of the general system of
classification of flowering plants. Ann.Missouri
Bot.Gard., 52 : 281-303.
- Cronquist, A. 1968. The evolution and classification of
flowering plants. Thomas Nelson (Printers) Ltd.
- Cronquist, A. 1981. An integrated system of classification
of flowering plants. Columbia University, press.
New York.
- Cronquist, A. 1987. A botanical critique of Cladism. Bot.
Rev., 53 : 1-52.
- Cordell, G.A. Lyon, R.L., Fong, H.H.S. and P.S. Benoit.
1977. Biological and phytochemical investigation
of Dianthus barbatus c.v ("Chinadall") Lloydia
(Clunc1)., 40(A) : 361-63.

- Csupor, L. 1970. Deoxy-rhaponticin, a new natural stilbene derivative in rhizome of Rheum rhaponticum. Arch. Pharm. Ber. Devt Pharm. Ges., 303(8) : 681-87.
- Dahlgren, R.H.T., 1980. A revised system of classification of the Angiosperms. J.Linn.Soc.Botany, 80 : 90-124.
- Dammer, 1895. Polygonaceae, In Willis - Dictionary of the flowering plants and Ferns (8th ed 1973) revised by H.R. Ainy Shaw) ^(H. Raven and J. E. Small) Cambridge.
- Daniel, M. and S.D. Sabnis. 1977. Chemotaxonomy of Loganiaceae. Curr.Sci., 48 : 383-385.
- Daniel, M. and S.D. Sabnis. 1986. Chemotaxonomic studies on some Indian members of Aizoaceae and Colluginaceae. Biome, 1(2) : 65-69.
- Davis, G.L. 1966. Systematic embryology of the Angiosperms. New York, London, Sydney : John Wiley and Sons.
- De Buhr L.E. 1975. Phylogenetic relationships of the Sarraceniaceae. Taxon, 24 : 297-306.
- De Eds, F. 1968. Title In Comprehensive Biochemistry Ed. M.Florkin and E.R. Stoltz. Elsevier publishing Co., Amsterdam.
- De Pascual-T.S., Govzalez, M.S., Vincente, S. and I.B. Jellido. 1981. Flavonoids from Chenopodium botryts. Planta Med., 41(4) : 389-91.
- De Pascual, A.E., Silvestri, G.E. and G.Tumino. 1980. Pharmacognostic research in Rheum compactum. Planta Med., 0(Suplt): 120-81.

- Dominquez, X.A., Jorgemarro, O. and M.M.Gutierrez. 1975.
Triterpene acetate and D-(+)- pinitol from
Drymaria drummondii. Phytochemistry (Oxf), 14(3):
815-16.
- Doskotch, R.W., Ray, A.D., Kubelka, W., Fairchild, E.H.,
Hufford, C.D. and J.S. Deal. 1974. Codonocarpus
alkaloids. II. The structure of Codonocarpine.
Tetrahedron, 30(8) : 3229-36.
- Doyle, J.J. 1983. Flavenoid races of Claytonia virginica
(Portulacaceae). Amer.J.Bot., 70: 1085-1091.
- Dubois, R.A., Zolc, A., Bouillant, M.L. and P. Delaveau. 1982.
Di-C-glycosyl flavones from Cerastium arvense L.
arvense new for Caryophyllaceae. Planta Med. 46(1) :
56-57.
- Dubois, M., Aleth, M.R. and J. Chopin. 1985. Isomollupentin
C-glucoside from Cerastium arvense. Phytochemistry,
24(5) : 1077-80.
- Eckardt, T. 1964. Centrospermae, in Melchior H. (ed.)
A. Engler's Syllabus der Pflanzenfamilien 1st ed.
Vol. III, 1-16. Berlin-Hirolasse.
- Ehrendorfer, F. 1976. Chromosome numbers and differentiation
of Centrospermae families. Pl.Syst.Evol., 126 :
27-30.
- Eichenberger, W. and W. Henke, 1966. Sterols in the leaves and
Chloroplast. Z.Naturforsch., 21b(a) : 650-65.

- Lichler, A.E. 1876. Syllabus der Vorlesungen über Phanerogamenkunde. Kiel : Schwesche-Buchhandlung.
- Lichler, A.E. 1878. Bluthendiagramme. Zweiter. Leipzig : W. Engelmann.
- Ellis, B.D. 1976. DGA ring cleavage in the biogenesis of stilobic acid in Cucurbita & Deeringiana. Phytochemistry, 15 : 489-91.
- Engler, G. 1964. Syllabus der Pflanzenfamilien. 12 Aufl. II. Band : Angiosperms (Hrsg. H. Melchior) Berlin-Mikolassee: Gebr. Borntraeger.
- Fukuyama, Y., Tsunoo, S., Asakawa, Y. and T. Takemoto. 1983. A potent cytotoxic carbonyl and related drimane-type sesquiterpenoids from Polygonum hydropiper. Phytochemistry, 21 (5) : 2895-99.
- Fukuyama, Y., Toshiyasu, F., Tsunoo, S., Iwamiya, H. and Y. Asakawa. 1985. Drimane-type sesquiterpenoid and nov. sesquiterpenoids from Polygonum hydropiper. Phytochemistry, 24 (7) : 1521-24.
- Garg, S.P., Bhushan, R. and R.C. Kapoor. 1980. Aervone, a new flavone from Aerva persica. Phytochemistry, 19: 1265-67.
- Geissman, T.A. (ed.). (1962). "Chemistry of the Flavonoid compounds". Pergamon Press Oxford.
- Gibbs, R.D. 1974. Chemotaxonomy of flowering plants McGill Queens, Univ. Press, Montreal.

- Grant, W.F. 1954. A cytological study of Gelosis argentea var. Cristata and their hybrids. Bot. Gaz., 115 : 324-35.
- Grinevich, L.I. and I.A. Ivanova, 1970. Pyropyrum emarginatum: A new potential source of rutin. Karmatsiya 19(1): 32-37.
- Gontar, L.K. and G.L. Vysockina, 1980. Anthraquinones in Humex rechingianum Subterranean organs of different colours. Rastit. Resur 16(1): 101-4.
- Gonzalez, A.G. Braton J.L., Castaneda, J.L., Praga B.A. and A. Morales. 1972. Triterpene from Phytolacca rivinoides Ph. An. Quim., 68(7/8): 1357-59.
- Habis, A.A.B., Hassen A.A.A and E.J. Suhtaci 1974. A new source of Anabazine, J. Pharm. Pharmacol., 26(10): 857-58.
- Hallier, H. 1912. L'origine et le systeme phyletique des Angiospermes expose. a l'aide de leur arbre genealogique. Arch. Neerl. Sci. Exact. Nat. Ser., 28: 1: 146-234.
- Harborne, J.B. 1967. Comparative Biochemistry of Flavonoids Academic press, London.
- Harborne, J.B. and C.A. Williams, 1971. O-hydroxy luteolin and coumestrol as phyletic markers in higher plants. Phytochemistry, 10: 367-373.
- Harborne, J.B. 1977. Flavonoids and the evolution of the angiosperms. Biochem. Syst. Ecol., 5 : 7-22.
- Harborne, J.B. 1984. Phytochemical methods, 2nd edn. Chapman and Hall, London.

- Hegnauer, R. 1966. Aucubinartige Glucoside. Über die Verbreitung und Bedeutung als systematisches Merkmal. Pharm. Acta Helv., 41 : 557-567.
- Hegnauer, R. 1969. Chemical evidence for the classification of some plant taxa - In Salain, T. and Harborne J.B. (eds.) Perspectives in Phytochemistry 1, 121-138.
- Hegnauer, R. 1971. Pflanzenstoffe und Pflanzensystematik Naturwiss., 58: 535-598.
- Heimerl, A., 1889. Phytolaccaceae, in Engler, A. and Prantl, R. Die natürlichen Pflanzenfamilien 1st ed. Vol. III, 1-16.
- Henry, A.J., Brion, C. and J.L. Guignard 1982. The saponins of Saponaria officinalis. Planta Med. Phytother., 15(4) : 192-200.
- Mikino, H.Y., Mikino, H., Komoto, T. and Takemoto, T. 1968. Cyasterone, an insect metamorphosing substance from Cyathula capitata: Structure. Tetrahedron, 24(13): 4855-4866.
- Mikino, H., Mikino, Y. and T. Takemoto. 1968. Synthesis of rubrosterone, a metabolite of insect moulting substance from Achyranthus rubrofusca. Tetrahedron Lett. 40 : 4255-56.
- Mikino, H., Komoto T., and T. Takemoto. 1970. Senego sterone, an insect metamorphosing substance from Cyathula capitata: Structure. Tetrahedron 26(3): 887-90.

- Mikino, H. and T. Takemoto. 1972. Arthropod moulting hormone from plants, Achyranthus and Cyathula. Naturwissenschaften, 59(3): 91-98.
- Rofmann U. 1973. Centrospermen-Studien 6. Morphologie und Systematik der Gattung Argyderma H.L.Br. (Desembryanthemaceae Tenzi). Diss. Hamburg.
- Holzschuh, L. Brigitte, K. and W. Hubelka. 1982. Physcion-3-O-β-D-gentiobioside, a new anthraquinone glycoside from Hubarb (Rheum palmatum) root. Planta Med. 46(3): 159-61.
- Hungund, J.L. and C.H. Nathak, 1971, U.S.D.A. Forest Service Research Paper RM-201.
- Hutchinson, J. 1948. British flowering plants, Academic Press, London 374.
- Hutchinson, J. 1959. Families of flowering plants. Dicotyledons Vol.1, Oxford University Press, London.
- Hutchinson, J. 1964. The genera of flowering plants. Dicotyledons Vol.1, Clarendon Press, Oxford.
- Hutchinson, J. 1967. The genera of flowering plants. Dicotyledons. Vol.2, Clarendon Press, Oxford.
- Hutchinson, J. 1973. The families of Flowering plants. (3rd edition) Clarendon Press, Oxford.
- Ibrahim, K.I. and G.H.M. Towers 1960. Arch. Biochem. and Biophys., 87: 125.
- Iljin, H. 1936. Chenopodiaceae. (V.L. Komarov and B.K. Schischkin Ed.) Flora Ussk. 6, 2-354.

- Iwashina, T., Ohniji, C. and Nakayashi. 1964.
 Neochilenin a new glycoside of 3-O-methyl quercetin
 and other flavonols in the tepals of Neochilenia,
Neoporteria and Porocia. Bot. Mag. Tokyo, 94(1045):
 23-30.
- Jensen, U. 1965. Serologische Untersuchungen zur Frage der
 systematischen Einordnung der Didiereaceae. Bot.
Jahrb. Syst. 84 : 273-53.
- Jensen, U.R., Nielsen, B.B. and A.L. Lohlgren. 1975. Iridoid
 compounds, their occurrence and systematic importance
 in angiosperms. Bot. Notiser, 128: 149-150.
- Johnson, N. and Y. Shimizu, 1974. Phytolaccinic acid, a new
 triterpene from Phytolacca americana. Tetrahedron,
30(4) : 2033-35.
- Joshi, R.C., Prakash, S. and P. Singh. 1973. Quinones and other
 constituents from Tabebuia rosea. Phytochemistry, 12:
 942.
- Kapadia, G.J., K.J. Shah and S. Kalucky, 1968. Peyote alkaloids II.
 Anhalotine, lophjotine and peyotine the quaternary
 alkaloids of Lophophora williamsii. J. Pharm. Sci.
57(2): 254-62.
- Kapadia, G.J. and R.T. Highet. 1968. Peyote alkaloids IV.
 Structure of peyyonine novel phenethylpyrrole from
Lophora williamsii. J. Pharm. Sci. 57(1): 191-92.
- Kartnig, I. 1972a. Determination of saponin in medicinal
 herbs. Planta Med., 21(1): 25-34.

- Kartnig, T. 1972 b. Studies on the preparation and stability of saponin drugs. Planta Med., 22(2) : 127-35.
- Kartnig, T. and C. Wegshalter, 1972. On the knowledge of the saponin from Hernairia glabra. Planta Med., 21 (2) : 144-49.
- Kimura, Y., Kozawa, M., Baba, K. and H. Kiyoshi. 1933. New constituents of roots of Polygonum cuspidatum. Planta Med., 48(3) : 164-68.
- Kirchner, H.W. and H.L. Bird. 1962. 3 6 Dihydroxy sterols in organpipe cactus Leptocarpus thurberi. Phytochemistry, 21(7) : 1705-10.
- Knight, J.C. and G.H. Etit. 1965. Arizona flora : The sterols of Leptocarpus argali. Phytochemistry, 5(2) : 47-82.
- Kopcewicz, J. 1972. Estrogen in the short day plants Perilla pectinifera and Chenopodium rubrum grown under inductive and non inductive light condition. Z. Pflanzen Physiol., 67(4) : 373-76.
- Kowal, T. 1954. Morphological and anatomical features of the seeds of the genus Anemone and key for their identification. Monogr-Botany, 2 : 163-65.
- Kretsu, L.G. and G.V. Lazur'evskii. 1970. Triterpene glycosides of Seta vulgaris. Khim. i fiz. kordin (Tashk), 6(2) : 272-73.
- Kringstad, R., Olsvik, A., G. Lingass and C.H. Gunn. 1980. 2-O-Methyl aldotetronic acid a new lactone forming acid present in plants. Phytochemistry, 19(4) : 543-48.

- Kryvenchuk, I.Ye., Lytvynenko, V.I. and V.M. Darnohral.
1968. Flavonoids from *Gypsophilla* leaves. Farm. Zh.
23(4) : 62-67.
- Kubitzki, R. 1969. Chemosystematische Betrachtungen zur
Grossgliederung der Dicotylen. Taxon, 18 : 360-
368.
- Kuroyanagi, H., Yamamoto, Y., Fukushima, S., Akira, U.,
Tadatakanoko, H. and T. Miyase. 1982. Chemical
studies on the constituents of *Polygonum nodosum*.
Chem. Pharm. Bull. (Tokyo), 30(5) : 1602-8.
- Lawrence, G.H.H. 1951. Taxonomy of vascular plants.
Clarendon press, Oxford.
- Leuck, E. and J.M. Miller 1982. Pollination biology and
Chemotaxonomy of the *Lechinocereus viridiflorus*
complex. Am. J. Bot., 69(10) : 1669-72.
- Livingstone, D., Marie Tomlinson, D.A., Friedman, G. and
R. Brooms. 1974. Stellate pore ornamentation in
pollen grains of the Amaranthaceae. Pollen et
Spores, 15(3)-4 : 345-51.
- Mabry, T.J., Taylor, A. and G.L. Turner, 1963. The
betacyanins and their distribution. Phytochemistry,
2 : 61-64.
- Mabry, T.J., Markham, K.R. and M.B. Thomas, 1970. The
Systematic identification of flavonoids. Springer-
Verlag, Berlin.

- Mabry, T.J. 1974. Is the order Centrospermae monophyletic.
In "Chemistry in Botanical Classification", G.Bendz
and J.Santesson, ed. Nobel Symposium 25.
- Mabry, T.J., Behnke, H.D. and I.J. Eifert. 1976. Betalains
and P-type sieve-element plastids in Gisekia L.
(Centrospermae) Taxon, 25(1) : 112-114.
- Mabry, T.J. 1977. The order Centrospermae. Ann.Missouri
Bot.Gard., 64 : 210-20.
- Maheshwari, P. 1964. Embryology in relation to taxonomy.
In Vistas in Botany, Turrill, W.B. Ed., Pergamon
Press, London. Vol. 4. pp. 55-97.
- Manske, R.H.F. 1944. The alkaloids. Ann.Rev.Biochem., 13 :
533-548.
- Mansour, R.M.A. and A.M. Saleh. 1981. 5,6,7-Trisubstituted-
flavones from Gomphrena maritima. Phytochemistry,
20(5) : 1173-79.
- Markham, K.R. 1982. Tehniques of Flavonoid Identification.
Academic Press, London.
- Mauritzon, J. 1939. Die Bedeutung der embryologischen
Forschung fur das naturliche System der pflanzen.
Lunds Univ.Arsskr. N.F.Adv., 2. 35(Nr.15): 1-70.
- Mc Neill, J. 1974. Synopsis of a revised classification of
the Portulacaceae. Taxon, 23(5/6) : 725-728.
- Meeuse, A.D.J. 1970. The descent of the flowering plants in
the light of new evidence from Phytochemistry and
from other sources. I and II Acta.Bot.Neerl., 19 :
61-72.

- Meyers, F.H., Jawetz, E. and Goldfien. 1972. Review of Medical Pharmacology 3rd edn. Lange Medical Publications, Los Altos, California.
- Meyer, B.N., Helfrich, J.S., Nichols, D.E., McLaughlin, J.L. Davis, D.V. and R.G. Cooks. 1983. Cactus alkaloid 53. Coryphanthine and O-methyl candicine, 2 new quaternary alkaloids from Coryphantha green woodii. J.Nat.Prod., 46(5) : 688-93.
- Miller, John, M. and B.A. Bohm. 1982. Flavonols and dihydroflavonol glycoside of Echinocereus triglochidiatus var. gurneyi. Phytochemistry, 24(4) : 951-52.
- Moquin-Tandon, A. 1849. Phytolaccaceae. In Prodromus systematis naturalis regni Vegetabilis, De Candolle, A., 13(2) : 2-40, Paris.
- Mukhamed' Yarova, M.M. 1968. Flavonoids of Polygonum persicaria. Khim Prior Soedin (Tashkent), 4(2) : 131.
- Muller, K. 1908. Beitrage zur Systematik der Aizoaccen. Bot.Jahrb., 42(Beibl. 97) : 54-94.
- Nakai, T. 1942. Notulae and plantas Asiae orientalis XVIII. J.Jap.Bot., 18 : 91-120.
- Neal, J.M., Sato, P.T., Howald, W.N. and J.L. McLaughlin. 1972. Peyote alkaloids : Identification in the mexican cactus Pelecypore aselliformis Ehrenberg. Science(Wash), 176(4036) : 1131-33.

- Niemann, G.J., Koerselmankooy, J.W., Steijns, J.M.
and K.Janvanbrederode. 1983. Flavone distribution
in leaves of different genotype of Silene pratensis.
Z.Pflanzen Physiol., 109(2) : 105-112.
- Nonaka, G.I., Nishihoka, I., Nagasawa, T. and H.Oura. 1981.
Tannin and related compounds in Rhubarb. Chem. Pharm.
Bull.(Tokyo), 29(10) : 2868-70
- Nonaka, G.I., Niwa, N. and I. Nishioka. 1982. Stilbene
glycoside gallates and proanthocyanins from Polygonum
multiflorum. Phytochemistry, 21(1) : 429-32.
- Nonaka, G.I., and I.Nishioka. 1983. Tannins and related
compounds. 10. Rhubarb : 2. Isolation and structure
of a glycerol gallide, gallic acid glycoside
gallates, galloyl glycoses and isolindleyin.
Chem.Pharm.Bull(Tokyo), 31(5) : 1652-58.
- Nonaka, G.I., Ezaki, E., Hayash, K. and I. Nishioka, 1983.
Flavonol glycosides from rhubarb and Rhaphirolepis
umbellata. Phytochemistry, 22(7) : 1659-62.
- Nowicke, J.W. 1975. Pollen morphology in the order
Centrospermae. Grana, 15 : 51-77.
- Nowicke, J.W. and J.J. Skvarla, 1977. Pollen morphology and
the relationship of the Plumbaginaceae, Polygonaceae
and Primulaceae to the order Centrospermae.
Smithsonian contributions to Botany, No. 37.

Smithsonian Institution Press, City of
Washington.

- Oshima, Y., Takatomi, O. and H. Hiroshihikino. 1984.
Structure of dianosides, C, D, E and F. Triterpenoid
saponins of Dianthus superbus Var. longicalycinum
herb. Planta Med. 0(1) : 43-47.
- Paech, K. and M.V. Tracey. 1965. Modern methods of plant
analysis, Vol. II. Springer-Verlag.
- Parkhurst, R.M. 1973. Molluscicidal saponins of Phytolacca
dodecandra : Oleanoglycotoxin-A. Phytochemistry,
12(6) : 1437-42.
- Pax, F. 1927. Zur Phylogenie der Caryophyllaceae Engl.
Bot. Jahrb., 61 : 223-41.
- Pax, F. and K. Hoffmann. 1934. In Engler, A. and Prantl :
Die natürlichen pflanzenfamilien. 2nd ed. Vol. 16c :
174-233.
- Pax, F. and K. Hoffmann. 1934. Alzooaceae, In Engler, A. and
K. Prantl. Die natürlichen Pflanzen familien 2nd ed.
Vol. 16c : 174-233.
- Petrescu, A.D. 1974. The chromatographic analysis of the
coumarins of Polygonum convolvulus L. Plant Med.
Phytother., 8(3) : 224-27.
- Plattelli, M., Minale and G. Prota, 1964. Isolation,
Structure and absolute configuration of indicaxanthin.
Tetrahedron, 20 : 2325.
- Price, J.R. 1963. The distribution of alkaloids in

- Rubiaceae. In Chemical Plant Taxonomy Swain T. (ed.) pp.429-452. Academic press, London.
- Pummangura, S., Mohamed, Y.A.H., Chang, C.J. and J.L. McLaughlin. 1982. 2. Simple tetrahydroxy isoquinoline alkaloid N-oxides from cacti. Phytochemistry, 21(9) : 2375-78.
- Putieva, Zh., Kondratenko, M.M.S. and N.K. Abubakirov. 1972. Triterpene glycosides of Acanthophyllum paniculata. Khim.Prir.Soedin(Tashk), 9(5) : 680-81.
- Rachel, M. and J.L. McLaughlin. 1980. Cactus alkaloid 42. 3,4-Dimethoxy phenethylamine and heliamine from the mexican ceroid, Backebergia militaris. J.Pharm.Sci. 69(1) : 94-95.
- Rada, K., Verma, H., Helena, S. and V. Brazdova. 1974. A study of Anthraquinone derivatives in Rumex species. Acta Fac.Pharm.Univ.Comenianae(Bratial), 25 : 153-76.
- Raghaven, T.S. and V.K. Srinivasan, 1940. Studies in the Indian Aizoaceae. Ann.Bot.(New Series), 4 : 651-62.
- Rai, J. and K.A. Thakar, 1970. Chemical investigation of Rumex vesicarium L., Indian J. Chem., 8(11) : 1046-47.
- Razdan, T.K., Harkar, S., Kachroo, V. and G.L.Koul. 1982. Phytolaccanol and epi acetylauritolic acid, 2 triterpenoids from Phytolacca acinosa. Phytochemistry, 21(9) : 2339-42.

- Razdan, R.N., Harkar, S., Kachroo, V., Koul, G.L. and
L.S. Nigam. 1983. Triterpenoids from Phytolacca
acutosa, 3 Oleanolic derivatives. Phytochemistry,
22(8): 1797-1800.
- Sengle, A.D. 1938. "The Classification of Flowering Plants"
2nd ed. (1975). Vol.11 (Dicotyledons).
- Richardson, M., 1978. Flavonols and C-Glycosylflavonoids of
the Caryophyllales. Biochem.Syst.Evol., 6 :283-286.
- Richardson, M.P. 1981. Flavonoids of some controversial
members of the Caryophyllales (Centrospermae). Pl.Syst.
Evol., 138: 227-33.
- Rochen, Th. 1927. Zur Embryologie der Centrospermen. Diss.
Uppsala.
- Rohwedder, C., 1965b. Centrospermen. Studien. 2. Entwicklung und
morphologische Deutung des Gynociums bei Phytolacca.
Bot. Jahrb.Syst., 84: 505-26.
- Rohwedder, C., 1965a. Centrospermen-Studien. 1. Der Blütenbau
bei Uebelinia kiwuensis T.C.L. Fries (Caryophyllaceae).
Bot. Jahrb. Syst., 83: 406-18.
- Romussi, G. and Ciarallo, 1974. 5,7 Dihydroxy chromone from
Polygonum persicaria seeds. Phytochemistry (Oxf),
13(12): 2890-91.
- Saatonov, Z., Usmanov, Z., and M.M. Abubakirov. 1979. Phytoecdysones
of Silene praemixta. I. Silenosterone. Khim.Prir.Soezin
(Tashk), 0(6): 793-97.

- Saatov, L., Gorovitis, M.B., Abdullaev, B.D., Usmanov, B.Z.
and N.K. Abubakirov. 1981. Phytoecdysteroids of
plant of the genus Silene. 3. A new glycoside
ecdysteroid from Silene brachuca. Khim.Frir.Boedin
(Tashk), 0(6): 738-44.
- Saatov, L., Abdullaev, Gorovits, M.B. and N.K. Abubakirov.
1985. Phytoecdysteroids of plants of the genus
Silene: 7, Sileneoide D. ecdysterone-3-O- -D galacto
pyranoside from Silene brachuca. Khim.Frir.Boedin.,
0(6) : 741-44.
- Sadykov, A.B., Aslanov, C.A. and J.K. Kusmuradov. 1966.
Quinoline alkaloids. Abhandl. Deut. Akad. Wiss. Berlin
(3) : 279-86.
- Sadykov, Yu.L., Khodzhimatov, N., and Yu.L. Komarov. 1978.
Alkaloids of fruit of Chenopodium botryts. Kastit.
Resur., 14(3): 385-87.
- Sandberg, F., Karl-Heinz K., Berntstaik, and T. Nelson, 1967.
Screening of Plants of the family Chenopodiaceae for
alkaloids. Acta Pharm. Suecica, 4(1): 51-58.
- Schinz, H., 1934. Amaranthaceae. Engler and Prantl. Die
naturlichen Pflanzenfamilien, ed. 2, Bd. 16c, 7-85.
- Schnarf, K., 1933. Die Bedeutung der embryologischen
Forschung für das natürliche System der pflanzen.
Biol. Gen., 9: 271-288.
- Sen, A.B. and Y.M. Shukla 1963. Chemical examination of
Hylocerus undatus. J. Indian Chem. Soc., 45(8): 745.

- Sharma, H.P. 1968. Morphological studies in the order Centrospermales. Bull. Bot. Soc. Univ. Saugar, 15(1/2): 27-35.
- Skvarla, J.J. and J.W. Novicke, 1976. Ultrastructure of Pollen exine in Centrospermous families. Plant Syst. Evol., 126, 55-78.
- Srinivasan, Lucas, S.T., Surrowes, C.B., Wanderman, H.A., Redner, A., Bernstein, S. and P.N. Sawyer, 1971. Effects of some flavonoids on the surface charge characteristics of the vascular system and their antithrombogenic characteristics. European Conf. Microcirculation, 6, 394.
- Soo, H. 1967. Die modern Systems der Angiospermen. Acta Bot. Acad. Sci. Hung., 13: 201-223.
- Souto, J. 1967. Triterpenoid Saponin in ripe fruits of *Aochia scoparia*. Rev. Inst. Agropecuar Ser. Z. Biol. Proves, 3(24): 367-83.
- Subramanyam, K. 1950. Proc. Indian Acad. Sci. B 31: 60-65.
- Takemoto, T.S., Ogawa, T., Nishimoto, K. and H. Hoffmeister. 1967. Steroids with moulting hormone activity from animals and plants. Z. Natur. Forsch., 22b(6): 681-82.
- Takemoto, T., Shuntaro, O. and K. Nishimoto. 1967. Studies on the constituents of Achyranthus radix. II. Isolation of the insect moulting hormone. Yakugaku Zasshi, 87(12) : 1462-73.

- Takemoto, T. 1968. Studies on the constituents of Achyranthus radix VII. The insect moulting substances in Achyranthus and Caythula genera: supliment. Yaku Gaku Zasshi, 88(10): 1293-97.
- Takemoto, T., Shuntaro, C. and N. Wshimoto. 1968. Studies on the constituents of Achyranthus radix. Yaku Gaku Zasshi, 87(12) : 1463-68.
- Takhtajan, A. 1959. De Evolution der Angiospermen. Jena : web G. Fischer.
- Takhtajan, A. 1966. Systems et Phylogenia Magnoliophytorum (in Russian) Moskan : Nauka.
- Takhtajan, A. 1960. Outline of the classification of flowering plants (Magnoliophyta). Bot. Rev., 46 : 225-359.
- Takashashi, K. and A. Brossi. 1962. Synthesis of weberene and nor weberene and their 1. Methyl congeners Retrocycles (Sendai), 19(4) : 691-96.
- Tandon, J.S. and K.P. Agarwal, 1966. Chemical Examination of Kocnia tricophylla. Indian J. Chem., 4(12) : 140-45.
- Thorne, R.F. 1968. Synopsis of a putatively phylogenetic classification of the flowering plants. Aliso, 6 : 57-66.
- Toth, I., Bathory, K., Swendrei, K., Minker, E. and G. Blazo. 1981. Ecdysteroids in Chenopodiaceae. Fitoterapia, 82(2) : 77-80.

- Trim, A.R. and R. Hill. 1951. The preparation and properties of acubin, Asperuloside and some related glycosides. Biochem.J., 50 : 310.
- Tschesche, R., Hubert, R. and G.wulf. 1969. Ueber Triterpe XXVIII Uber die saponinedes Spinats (Spinacia oleraceae) (Triterpenes XXVIII. The saponins of Spinach. Justus Liebigs Ann.Chem., 726 : 125-35.
- Ulbrich, E. 1934. Chenopodiaceae. In A.Engler and H.Haras, Die Naturlichen Pflanzen familien, 16c : 375-584.
- Usmanghani, K., Nazir, T. and V.U.Ahamed. 1982. Chemical constituents of Aerva javanica. Fitoterapia, 53(3) : 75-78.
- Van Rompuy, R., Ludo, L.L. and J.A.P. Zeevaart. 1976. Steroidal estrogens natural Plant Constituents. Phytochemistry (Oxf), 18(5) : 863-66.
- Varnaite, R.N., Pipinis, I.A. and R.S. Zaprometov, 1972. Phenolic compounds of alpine joint weed. Prikl. Biokhim.Mirobiol., 8(1) : 57-61.
- Verdejo vivos, G. 1974. Cactus alkaloids. Arch Inst. Acim Almeria Esp., 19 : 21-56.
- Vishnu-Mitre, N. 1963. Pollen morphology of Indian Amaranthaceae J.Indian Bot.Soc., XLII (1) : 186-101.
- Vyechar, A.S., Ruznyatsova, Z.P. and I.I.Chakalinskaya 1980. Phenol Carboxylic acid in species of the

- genus Polygonum. Vyestsi Akad Navuk ssr Syer Bival
Navuk, 0(6) : 71-74.
- Wagner, W.H. Jr. 1980. Origin and Philosophy of ground
plan-divergence methods of cladistics. Syst.Bot.,
5 : 173-193.
- Weiffening, J.H. 1966. Phytochemistry, 5 : 1055.
- West, L.G., Jeffry, L., McLaughlin, L. and W.H. Larle, 1975.
Filocrenine from Lophocereus schottii Monastrosus
and Miecklyanus. Phytochemistry, 14(1) : 291-92.
- Weirmann, R., Eckhard, W. and C.kehse, 1981. Yellow
flavonoids as component of pollen pigmentation.
Z.Naturforsch Sect C.Biosci., 36(3/4) : 204-6.
- Williams, J.T. and R.V. Ford-Lloyd. 1974. The Systematics
of Chenopodiaceae. Taxon, 23(2/3) : 353-54.
- Woo, W.S. 1973. The structure of paligonic acid, a new
triterpene from Phytolacca esculenta. Lloydia
(Cinci), 36(3) : 326-32.
- Woo, W.S. 1974. Steroids and pentacyclic tritenpenoid
from Phytolacca americana. Phytochemistry, 13(12) :
2887-89.
- Woodard, R., Cymermancaril, W.J. and J.B.Bruhn. 1978. The
absolute configuration of the cactus alkaloid (-).
calipamine. Acta Chem.Scand Ser B Org. Chem.Biochem.
32(8) : 619-20.
- *Wylar, H., Mabry, T.J. and A.S. Dreiding. 1963. Helv.Chim.
Acta, 46 : 1745.

* Wyler, W., Rosler, H., Mercier, E. and A.S. Dreiding.

1967. Helv.Chim.Acta, 50 : 545.

Yasukawa, K., Yamanouchi, S. and K. Tarido. 1981.

Constituents of water extracts of crude drugs. 3.

Roots of Stellaria dichotama var. lanceolata.

Yakugaku Zasshi, 101(1) : 64-66.

Yukhananov, D.Kh., Yatsyno, A.I., Turova, A.D. and V.S.

Kyvayev. 1972. The saponin content in the representative of Caryophyllaceae and their hemolytic activity.

Rastit Resur, 8(3) : 574-77.

Yukhananov, D.Kh. and L.A. Krasnov. 1973. Distribution of

triterpenoid glycosides and flavonoids of the family

Caryophyllaceae (1) Qualitative evaluation of plants

of the family Caryophyllaceae for the presence of

biologically active substances. Rastit Resur, 9(4) :

540-47.

Zykova, N.Ya., Madio, M. and M. Kovu, 1976. Flavonoids of

the white Campion. Farm Zh (Kiev), 31(4) : 61-63.

* Not referred to the original article.