




BIBLIOGRAPHY



Abdul-Ghani M, Nawaf G, Nawaf F, Itzhak B, Minuchin O, Vardi P. Increased prevalence of microvascular complications in type 2 diabetes patients with the metabolic syndrome. IMAJ;2006; 8: 378–382.

Adler AI, Stevens RJ, Manley SE, Bilous RW, Cull CA, Holman RR; UKPDS GROUP. Development and progression of nephropathy in type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS 64). *Kidney Int*; 2003; 63: 225–32.

Agrawal P, Rai V, Singh RB, Azad Khan AK, Akhtar S, Mahtab H. Randomized placebo-controlled single-blind trial of holy basil leaves in patients with non insulin-dependent diabetes mellitus: *Coccinia indica* in the treatment of patients with diabetes mellitus. *Int J Clin Pharmacol Ther*; 1996; 34: 406–409.

Agarwal SK, Dash SC. Spectrum of renal disease in Indian adults. *JAPI*; 2000; 48: 594 - 600.

Ahuja MMS. A decade of epidemiology of diabetes. *Intl. J. Diab. Dev. Countries*; 1991; 11: 2-4.

Ajay VS, Prabhakaran D, Jeemon P, Thankappan KR, Mohan V, Ramakrishnan L et al. Prevalence and determinants of diabetes mellitus in the Indian industrial population. *Diabet Med*; 2008; 25: 1187-94.

Alexander C, Landsman PB, Teutsch SM, Haffner SM. NCEP III defined metabolic syndrome, diabetes and prevalence of coronary heart disease among NHANES III participants age 50 years and older. *Diabetes*; 2003; 52: 1210–14.

Alwan A, MacLean DR. A review of non-communicable disease in low- and middle-income countries. *International Health*; 2009; 1: 3-9.

American Association of Clinical Endocrinologists/ American College of Endocrinologists (AACE/ ACE). AACE/ACE position statements on the prevention, diagnosis and treatment of obesity. *Endocrine Practise*; 1998; 4: 279- 230.

American Diabetes Association. Nephropathy in diabetes. Position statement. *Diabetes Care*; 2004; 27:S79–S83.

American Diabetes Association. Treatment of hypertension in adults with diabetes. *Diabetes Care*; 2003; 26: S80-S82.

American Diabetes Association. Standards of Medical Care in Diabetes-2009 (Position Statement). *Diabetes Care*; 2009; 32: S13-S61.

American Diabetes Association, National Institute of Diabetes and Digestive and Kidney Diseases. Prevention or delay of type 2 diabetes. *Diabetes Care*; 2004; 27: S47–S54.

American Diabetes Association. Economic consequences of diabetes mellitus in the U.S. in 2002. *Diabetes Care*; 2003; 26: 917–932.

American Diabetes Association. Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*; 2009; 32: S62-S67.

American Diabetes Association. Nutrition recommendations and interventions for diabetes. *Diabetes Care*; 2008; 31: S61-S78.

Amos AF, McCarty DJ, Zimmet P. The rising global burden of diabetes and its complications: estimates and projections to the year 2010. *Diabetic Medicine*; 1997; 14: S7 - S85.

Anderson JW, Johnstone BM, Cook-Newell ME. Metaanalysis of the effects of soybean protein intake on serum lipids. *N Engl J Med*; 1995; 333: 276-282.

Anderson JW, Zeigler JA, Deakins DA, Floore TL, Dillon DW, Wood CL et al. Metabolic effects of high carbohydrate, high fiber diet, for IDDM individuals. *Am J Clin Nutr*; 1991; 54: 936- 941.

Araki S, Haneda M, Koya D, Hidaka H, Sugimoto T, Isono M et al. Reduction in microalbuminuria as an integrated indicator for renal and cardiovascular risk reduction in patients with type 2 diabetes. *Diabetes*; 2007; 56: 1727-1730.

Aronoff SL, Berkowitz K, Shreiner B, Want L. Glucose metabolism and regulation: beyond insulin and glucagons. *Diabetes Spectrum*; 2004; 17: 183-190.

Arya D, Nandave M, Ojha S, Kumari S, Joshi S, Mohanty I. Myocardial scavenging effects of *Ocimum Sanctum* in experimental model of myocardial necrosis: a hemodynamic, biochemical and histoarchitectural assessment. *Current Science*; 2006; 91: 265-269.

Ascherio A, Katan M, Zock P, Stampfer M, Willet W. Trans fatty acids and coronary heart disease. *N Eng J Med*; 1999; 340: 1994-1998.

Assmann G, Schulte H. Relation of highdensity lipoprotein cholesterol and triglycerides to incidence of atherosclerotic coronary artery disease (the PROCAM experience): Prospective Cardiovascular Munster study. *Am J Cardiol*; 1992; 70: 733– 737.

Bahl VK, Prabhakaran D, Karthikeyan G. Coronary artery disease in Indians. *Indian Heart J*; 2001; 53: 707-13.

Bailey CJ, Day C. Traditional plant medicines as treatments for diabetes. *Diabetes Care*; 1989; 12: 553–564. .

Bantle J, Swanson J, Thomas W, Laine D. Metabolic effects of dietary sucrose in type II diabetic subjects. *Diabetes Care*; 1993; 16: 1301-1305.

Basi S, Mimran A, Fesler P, Lewis JB. Microalbuminuria in type 2 diabetes and hypertension-a marker, treatment target, or innocent bystander? *Diabetes Care*; 2008; 31: S194-S201.

Baskaran K, Ahamath BK, Shanmugasundaram KR, Shanmugasundaram ERB. Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin dependent diabetes mellitus patients. *J Ethnopharmacology*; 1990; 30: 295–305.

Baynes JW, Thorpe SR. Perspectives in diabetes. Role of oxidative stress in diabetic complications. A new perspective on an old paradigm. *Diabetes*; 1999; 48: 1–9.

Bazzano LA. Dietary intake of fruit and vegetables and risk of diabetes mellitus and cardiovascular disease. World Health Organization, 2005. Available at http://www.who.int/dietphysicalactivity/publications/f&v_cvd_diabetes.pdf

Benedet JA, Umeda H, Shibamoto T. Antioxidant activity of flavonoids isolated from young green barley leaves toward biological lipid samples. *J Agric Food Chem*; 2007; 55: 5499–5504.

Best MM, Duncan CH, Van Loon EJ, Wathen JD. Lowering of serum cholesterol by the administration of a plant sterol. *Circulation*; 1954; 10: 201–206.

Bharadwaj PK, Dasgupta DK, Parashar SS, Kaushal SS. Effective reduction of LDL cholesterol by indigenous plant products. *J. Ind. Med. Assoc*; 1994; 92: 80–81.

Bhatia V, Arya V, Dabadghao P, Balasubramaniam K, Sharma K, Verghese N et al. Etiology and outcome of childhood and adolescent diabetes mellitus in north India. *J Pediatr Endocrinol Metab*; 2004; 17: 993–999.

Bonora E, Kiechl S, Willeit J, Oberhollenzer F, Egger G, Targher G et al. Prevalence of insulin resistance in metabolic disorders: the Bruneck Study. *Diabetes*; 1998; 47: 1643–1649.

Bonora E, Targher G, Formentini G, Calcaterra F, Lombardi S, Marini F et al. The Metabolic Syndrome is an independent predictor of cardiovascular disease in Type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study. *Diabet Med*; 2004; 21: 52–58.

Boyko EJ, Fujimoto WY, Leonetti DL, Newell-Morris L. Visceral adiposity and risk of type 2 diabetes: a prospective study among Japanese Americans. *Diabetes Care*; 2000; 23: 465–471.

Brantsma AH, Atthobari J, Bakker SJL, de Zeeuw D, de Jong PE, Gansevoort RT for the PREVEND Study Group. What predicts progression and regression of urinary albumin excretion in the nondiabetic population? *J Am Soc Nephrol*; 2007; 18: 637–645.

Brook RD, Bard RL, Kehrner C, Bodary PF, Eitzman DL. Effect of the short term weight loss on the metabolic syndrome and conduit vascular endothelial function. *Am. J. Cardiol*; 2004; 93: 1012–1016.

Burden AC, McNally PG, Feehally J, Walls J. Increased incidence of end-stage renal failure secondary to diabetes mellitus in Asian ethnic groups in the United Kingdom. *Diabet Med*; 1992; 9: 641–645.

Burton WN, Conti DJ, Chen CY, Schultz AB, Edington DW. The role of health risk factors and disease on worker productivity. *Journal of Occupational & Environmental Medicine*; 1999; 41: 863–877.

Campos KE, Diniz YS, Cataneo AC, Faine LA, Alves MJQF, Novelli ELB. Hypoglycemic and antioxidant effect of onion *Allium cepa*: dietary onion addition, antioxidant activity and hypoglycemic effects on diabetic rats. *Int J Food Sci Nutr*; 2003; 54: 241–246.

Carnethon M, Whitsel LP, Franklin BA, Kris-Etherton P, Milani R, Pratt CA et al on behalf of the American Heart Association Advocacy Coordinating Committee; Council

on Epidemiology and Prevention; Council on the Kidney in Cardiovascular Disease; and Council on Nutrition, Physical Activity and Metabolism. Worksite wellness programs for cardiovascular disease prevention: a policy statement from the American Heart Association. *Circulation*; 2009; 120: 1725–1741.

Chandalia M, Garg A, Lutjohann D, von Bergmann K, Grundy SM, Brinkley LJ. Beneficial effects of high dietary fibre intake in patients with type 2 diabetes. *N Engl Journal of Medicine*; 2000; 342: 1392-1398.

Chandie SPK, Baboe F, van Es LA, van der Vijver JC, van de Ree MA, de Jonge N et al. South-Asian type 2 diabetic patients have higher incidence and faster progression of renal disease compared with Dutch-European diabetic patients. *Diabetes Care*; 2006; 29: 1383-1385.

Chandie SPK, Vandenbrouke JP, Tjandra YI, Rosendaal FR, Rosman JB, Geerlings W et al. Increased end-stage diabetic nephropathy in Indo-Asian immigrants living in the Netherlands. *Diabetologia*; 2002; 45: 337-341.

Chattopadhyay RR. Hypoglycemic effect of *Ocimum sanctum* leaf extract in normal and streptozotocin diabetic rats. *Ind J Exp Biol*; 1993; 31: 891–893.

Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA*; 2003; 289: 2560–2572.

Choi Y, Cho Y, Park C, Jang H, Park K, Kim S et al. Rapidly increasing diabetes-related mortality with socio-environmental changes in South Korea during the last two decades. *Diabetes Res Clin Pract*; 2006; 74: 295–300.

Chowdhry TA, Lasker SS, Mahfuz R. Ethnic differences in control of cardiovascular risk factors in patients with type 2 diabetes mellitus attending inner circle diabetic clinic. *Post grad Med J*; 2006; 82: 211-215.

Colditz GA, Willett WC, Stampfer MJ, Manson JE, Hennekens CH, Arky RA et al. Weight as a risk factor for clinical diabetes in women. *Am J Epidemiol*; 1990; 132: 501–513.

Cook C, Simmons G, Swinburn B, Stewart J. Changing risk behaviours for non-communicable disease in New Zealand working men - Is workplace intervention effective? *New Zealand Medical Journal*; 2001; 114: 175-178.

Costa LA, Canani LH, Lisboa HRK, Tres GS, Gross JL. Aggregation of features of the metabolic syndrome is associated with increased prevalence of chronic complications in Type 2 diabetes. *Diabetic Medicine*; 2004; 21: 252-255.

Cremer L, Herold A, Avram D, Szegli G. A purified green barley extract with modulatory properties upon TNF alpha and ROS released by human specialised cells isolated from RA patients. *Roum Arch Microbiol Immunol*. 1998 Jul-Dec;57(3-4):231-42.

Damsgaard EM, Mogensen CE. Microalbuminuria in elderly hyperglycaemic patients and controls. *Diabet Med*; 1986; 3: 430–35.

Daniel M, Marion SA, Shepa SB, Gamble D. Variation by BMI and age in waist to hip associations with glycemic status in an aboriginal population at risk for type 2 diabetes in British Columbia, Canada. *Am J Clin Nutr*; 1999; 69: 455-460 .

DECODE – DECODA study group on behalf of the European Diabetes Epidemiology group and the International Diabetes Epidemiology group. Age, BMI, and type 2 diabetes associations modified by ethnicity *Diabetologia*; 2003; 46: 1063-1070.

DeFronzo RA. Pathogenesis of type 2 diabetes: metabolic and molecular implications for identifying diabetes genes. *Diabetes Rev*; 1997; 5: 177–269.

DeFronzo RA, Bonadonna RC, Feranninni. Pathogenesis of NIDDM, A balanced view. *Diabetes Care*; 1992; 15: 618- 624.

DeFronzo RA, Ferrannini E. Insulin resistance: a multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia and atherosclerotic cardiovascular disease. *Diabetes Care*; 1991; 14: 173–94.

Deepa R, Sandeep S, Mohan V. Abdominal Adiposity, Visceral fat and type 2 diabetes-“Asian Indian phenotype” In: *Type 2 Diabetes Mellitus in South East Asians*. Ed. Mohan V, Rao G; 2007; pp, 138- 163.

Dengel DR, Kelly AS, Thomas PO, Kaiser DR, Dengel JL, Bank AJ. Effects of weight loss on insulin sensitivity and arterial stiffness in overweight adults. *Met Clin Exptl*; 2006; 55: 907:911.

Desai S, Mani U. Risk factor analysis in the development of chronic degenerative diseases in an industrial setup. PhD. Dissertation thesis, Dept. of foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda. Baroda, 2003

Despres JP, Lemieux I, Prud'homme D. Treatment of obesity: need to focus on high risk abdominally obese patients. *BMJ*; 2001; 322: 716-720.

Dixon AN, Raymond NT, Mughal S, Rahim A, O'Hare JP, Kumar S et al. Prevalence of microalbuminuria and hypertension in South Asians and white Europeans with type 2 diabetes: a report from the United Kingdom Asian Diabetes Study (UKADS). *Diabetes Vasc Dis Res*; 2006; 3: 22–5.

Djoussé L, Arnett DK, Coon H, Province MA, Moore LL, Ellison RC. Fruit and vegetable consumption and LDL cholesterol: the National Heart, Lung and Blood Institute Family Heart Study. *Am J Clin Nutr*; 2004; 79: 213-217.

Ducimetiere P, Richard J, Cambien, F. The pattern of subcutaneous fat distribution in middle-aged men and the risk of coronary heart disease: the Paris Prospective Study. *Int J Obes*; 1986; 10: 229–240.

The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med*; 1993; 329: 977–86.

Eddouks M, Maghrani M, Zeggwagh NA, Michel JB. Study of the hypoglycaemic activity of *Lepidium sativum* L. aqueous extract in normal and diabetic rats. *J. Ethnopharmacol*; 2005; 97: 391-395.

Edwards K, Kwaw I, Matud J, Kurtz I. Effect of pistachio nuts on serum lipid levels in patients with moderate hypercholesterolemia. *J Am Coll Nutr*; 1999; 18: 229 – 232.

Eely E, Stratton I, Haddon D, Turner R, Holman R. On behalf of the U.K. Prospective Diabetes Study Group: UKPDS 18: estimated dietary intake in type 2 patients with diabetes randomly allocated to diet, sulphonylurea or insulin therapy. *Diabet Med*; 1996; 13: 656–662.

Elbein SC. The genetics of human noninsulin-dependent (type 2) diabetes mellitus. *J Nutr*; 1997; 127: 1891S-1896S.

Engbers LH, van Poppel MN, Paw MJ, van Mechelen W. Worksite health promotion programs with environmental changes: a systematic review. *Am J Prev Med*; 2005; 29: 61–70.

Eriksson KF, Lindgarde F. Prevention of type 2 (non-insulin-dependent) diabetes mellitus by diet and physical exercise The 6-year Malmö feasibility study. *Diabetologia*; 1991; 34: 891-898.

Fagot C, Pettitt D, Engelgau M. Type 2 diabetes among North American children and adolescents: an epidemiologic review and a public health perspective *J Pediatric*; 2000; 136: 664-672.

Ferreres F, Krsková Z, Gonçalves RF, Valentão P, Pereira JA, Dusek J et al. Free water-soluble phenolics profiling in barley (*Hordeum vulgare* L.). *J Agric Food Chem*; 2009; 57: 2405-9.

Feskens EJ, Virtanen SM, Räsänen L, Tuomilehto J, Stengård J, Pekkanen J et al. Dietary factors determining diabetes and impaired glucose tolerance. A 20 year follow up of the Finnish and Dutch cohorts of the seven countries study. *Diabetes Care*; 1995; 18: 1104-1112.

Fischbacher C, Bhopal R, Rutter M, Unwin N, Marshall S, White M, Alberti K. Microalbuminuria is more frequent in South Asian than in European origin populations: a comparative study in Newcastle, UK. *Diabet Med*; 2003; 20: 31–36.

Fontbonne A, Eschwege E, Cambien F, Richard J, Ducimetiere P, Thibault N et al. Hypertriglyceridemia as a risk factor of coronary heart disease mortality in subjects with impaired glucose tolerance or diabetes: results from the 11 year follow up of the Paris Prospective Study. *Diabetologia*; 1989; 32: 300-4.

Ford ES, Imperatore G, Li C, Cook S. Age, sex, and ethnic variations, in serum insulin concentrations among US youth: findings from the National Health and Nutrition Examination Survey 1999-2002. *Diabetes Care*; 2006; 29: 2605-2611.

Friedberg C, Janssen M, Heine R, Grobbee D. Fish oil and glycemic control in diabetes: a meta-analysis. *Diabetes Care*; 1998; 21: 494–500.

Fujimoto W, Abbate S, Kahn S, Hokanson J, Brunzell J. The visceral adiposity syndrome in Japanese-American men. *Obes Res*; 1994; 2: 364–371.

Fuster V, Badimon L, Badimon J, Chesebro J. The pathogenesis of coronary heart disease and the acute coronary syndromes *N Eng J Med*; 1992; 326: 242-50.

Gaede P, Tarnow L, Vedel P, Parving HH, Pedersen O. Remission to normoalbuminuria during multifactorial treatment preserves kidney function in patients with type 2 diabetes and microalbuminuria. *Nephrol Dial Transplant*; 2004; 19: 2784–2788.

Garg A. High-monounsaturated fat diets for patients with diabetes mellitus: a meta-analysis. *Am J Clin Nutr*; 1998; 67: 577S–582S.

Garg A, Bantle J, Henry R, Coulston A, Griver K, Raatz S et al. Effects of varying carbohydrate content of diet in patients with non-insulin dependent diabetes mellitus. *JAMA*; 1994; 271: 1421–1428.

Gerich JE. The genetic basis of type 2 diabetes mellitus: impaired insulin secretion versus impaired insulin sensitivity. *Endocrine. Rev*; 1998; 19: 491-503.

Gerstein HC, Mann JF, Yi Q, Zinman B, Dinneen SF, Hoogwerf B et al. Albuminuria and risk of cardiovascular events, death, and heart failure in diabetic and nondiabetic individuals. *JAMA*; 2001; 286: 421-426.

Giacco R, Parillo M, Rivellese AA, Lasorella G, Giacco A, D'Episcopo L et al. Long-term dietary treatment with increased amounts of fiber-rich low-glycemic index natural foods improves blood glucose control and reduces the number of hypoglycemic events in type 1 diabetic patients. *Diabetes Care*; 2000; 23: 1461–1466.

Goenka S, Prabhakaran D, Ajay VS, Reddy KS. Preventing cardiovascular disease in India –translating evidence to action. *Current Science*; 2009; 97: 367-377.

Gordon DJ, Probstfield JL, Garrison RJ, Neaton JD, Castelli WP, Knoke JD et al. High-density lipoprotein cholesterol and cardiovascular disease: four prospective American studies. *Circulation*; 1989; 79: 8-15.

Goyal A, Yusuf S. The burden of cardiovascular diseases in the Indian sub continent. *Indian J Med Res*; 2006; 124: 235-44.

Groop LC, Bonadonna RC, DelPrato S, Ratheiser K, Zyck K, Ferrannini E et al. Glucose and free fatty acid metabolism in non-insulin-dependent diabetes mellitus. Evidence for multiple sites of insulin resistance. *J. Clin. Invest*; 1989; 84: 205–213.

Grover JK, Gupta SR. Hypoglycaemic activity of seeds of *Momordica charantia*. *Eur. J. Pharmacol* ; 1990 ; 183: 1026-1027.

Grundy SM, Brewer HB, Cleeman JI, Smith SC, Lenfant C. for the Conference Participants. Definition of metabolic syndrome. Report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on Scientific Issues Related to Definition. *Circulation*; 2004; 109: 433–438.

Gu D, Reynolds K, Duan X, Xin X, Chen J, Wu X et al. Prevalence of diabetes and impaired fasting glucose in the Chinese adult population: International Collaborative Study of Cardiovascular Disease in Asia (InterASIA). *Diabetologia*; 2003; 46: 1190-8.

Gupta DK, Verma LK, Khosla PK, Dash SC. The prevalence of microalbuminuria in diabetes: a study from north India. *Diabetes Res Clin Pract*; 1991; 12: 125–128.

Gupta R, Johri S, Saxena AM. Diabetes mellitus: The Pandemic of 21st Century! Asian J. Exp. Sci; 2009; 23: 261-268.

Hadden DR, Montgomery DAD, Skelly RJ, Trimble ER, Weaver JA, Wilson EA. Maturity onset diabetes mellitus: response to intensive dietary management. BMJ; 1975; 3: 276-278.

Haffner SM. Obesity and the metabolic syndrome: the San Antonio Heart Study British Journal of Nutrition; 2000; 83: S67-S70.

Haffner SM, Lehto S, Ronnema T, Pyörälä K, Laakso M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. N Engl J Med; 1998; 339: 229-234.

Haffner SM, Mykkanen L, Festa A, Burke JP, Stern MP. Insulin-resistant prediabetic subjects have more atherogenic risk factors than insulin-sensitive prediabetic subjects: implications for preventing coronary heart disease during the prediabetic state. Circulation; 2000; 101: 975-980.

Haffner SM, Stern MP, Hazuda HP, Mitchell BD, Patterson JK. Cardiovascular risk factors in confirmed prediabetic individuals: does the clock for coronary heart disease start ticking before the onset of clinical diabetes? JAMA; 1990; 263: 2893-2898.

Hagiwara Y. Green barley essence. Keats Publishing. 1986

Hara Y, Honda M. The Inhibition of α -Amylase by Tea Polyphenols. Agric. Biochem; 1990; 54: 1939-1945.

Harris W, Connor W, Illingworth D, Rothrock D, Foster D. Effects of fish oil on VLDL triglyceride kinetics in humans. J Lipid Res; 1990; 31: 1549-1558.

Harris M, Flegal K, Cowie C, Eberhardt M, Goldstein D, Little R et al. Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults: The Third National Health and Nutrition Examination Survey, 1988-1994. Diabetes Care; 1998; 21: 518-24.

Hayek T, Fuhrman B, Vaya J, Rosenblat M, Belinky P, Coleman R, et al. Reduced progression of atherosclerosis in apoprotein deficient E mice following consumption of red wine and its polyphenols quercetin or catechin is associated with reduced susceptibility of LDL to oxidation and aggregation. Arterioscler Thromb Vasc Biol; 1997; 17: 2744-2752.

Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. N Engl J Med; 1991; 325: 147-152.

Heinemann T, Axtmann G, Von Bergmann K. Comparison of intestinal absorption of cholesterol with different plant sterols in man. European Journal of Clinical Investigation; 1993; 23: 827-831.

Heinemann T, Kullak-Ublick GA, Pietruck B, von Bergmann K. Mechanisms of action of plant sterols on inhibition of cholesterol absorption-Comparison of sitosterol and sitostanol. European Journal of Clinical Pharmacology; 1991; 40: S59-S63.

Herman WH, Shahinfar S, Carides GW, Dasbach EJ, Gerth WC, Alexander CM et al. Losartan reduces the costs associated with diabetic end-stage renal disease. *Diabetes care*; 2003; 26: 683 – 687.

Hertog MGL, Feskens EJM, Holman PCH, Katan MB, Kromhout D. Dietary antioxidant flavonoids and risk of CHD: The Zutphen Elderly Study. *Lancet*; 1993; 342: 1007-1012.

Hertog MGL, Kromhout D, Aravanis C, Blackburn H, Buzina R, Fidanza F. Flavonoid intake and long term risk of coronary heart disease and cancer in the Seven countries study. *Arch Intern Med*; 1995; 155: 381-386.

Hertz RP, Unger AN, McDonald M, Lustik MB, Biddulph-Krentar JBS. The Impact of Obesity on Work Limitations and Cardiovascular Risk Factors in the U.S. Workforce. *Journal of Occupational and Environmental Medicine*; 2004; 46: 1196-1203.

Hillege HL, Fidler V, Diercks GFH, van Gilst WH, de Zeeuw D, van Veldhuisen DJ et al for The Prevend Study Group. Urinary Albumin Excretion Predicts Cardiovascular and Noncardiovascular Mortality in General Population. *Circulation*; 2002; 106: 1777-1782.

Hillege HL, Janssen WMT, Bak AAA, Diercks GFH, Grobbee DE, Crijns HJGM et al for The Prevend Study Group. Microalbuminuria is common, also in a nondiabetic, nonhypertensive population, and an independent indicator of cardiovascular risk factors and cardiovascular morbidity *Journal of Internal Medicine*; 2001; 249: 519-526.

Hokanson JE, Austin MA. Plasma triglyceride level is a risk factor for cardiovascular disease independent of highdensity lipoprotein cholesterol level: a meta-analysis of population-based prospective studies. *J Cardiovasc Risk*; 1996; 3: 213–219.

Hu FB, Manson JE, Stampfer MJ, Willet CC. Diet, lifestyle and risk of type 2 diabetes mellitus in women. *NEJM*; 2001; 343: 790-797.

Hu FB, van Dam RM, Liu S. Diet and risk of type II diabetes: the role of types of fat and carbohydrate. *Diabetologia*; 2001; 44: 805 –817.

Hu G, Sarti C, Jousilahti P, Peltonen M, Qiao Q, Antikainen R et al. The impact of history of hypertension and type 2 diabetes at baseline on the incidence of stroke and stroke mortality. *Stroke*; 2005; 36: 2538-2543.

Huxley R, Barzi F, Woodward M. Excess risk of fatal coronary heart diseases associated with diabetes in men and women: meta analysis of 37 prospective cohort studies. *Br. Med. J*; 2006; 332:73-78.

The IDF Consensus worldwide definition on the Metabolic Syndrome.
http://www.idf.org/webdata/docs/IDF_Meta_def_final.pdf (p. 13)

IDF Diabetes Atlas, 4th edition; International Diabetes Federation (2009). Available from: <http://www.diabetesatlas.org>.

Ignacimuthu S, Amalraj T. Effect of leaf extract of *zizyphus Jujuba* on diabetic rats. *Indian J. Pharmacol*; 1998; 30: 107–108.

International Labour Organization

LABORSTA, Geneva, International Labour Organization, 2007 (available at: <http://laborsta.ilo.org/>).

Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nisse M, Taskinen M, Groop L. Cardiovascular Morbidity and Mortality Associated With the Metabolic Syndrome. *Diabetes Care*; 2001; 24: 683–689.

Isomaa B, Henricsson M, Almgren P, Tuomi T, Taskinen MR, Groop L. The metabolic syndrome influences the risk of chronic complications in patients with type II diabetes. *Diabetologia*; 2001; 44: 1148-1154.

Iyer U, Desai P. Risk factor analysis of diabetes mellitus and impact of panchratna juice supplementation in the management of diabetes mellitus. PhD Dissertation thesis, Dept. of Foods and Nutrition, Faculty of Family and Community Sciences, The Maharaja Sayajirao University of Baroda. Baroda, 2008

Iyer U, Dholakia P. Beverage consumption pattern in delhi and vadodara: nutrient analysis of commonly consumed beverages. M.Sc Dissertation thesis, Dept. of foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda. Baroda, 2002

Iyer U, Joshi A, Dhruv S. Impact of amla (*embilica officinalis*) supplementation on the glycemic and lipidemic status of type 2 diabetic subjects. *Journal of Herbal Medicine and Toxicology*; 2009; 3: 15-21.

Iyer UM, Mani UV. Studies on the effect of curry leaves supplementation on the lipid profile, glycated proteins and amino acids in NIDDM patients. *Plant Foods for Human Nutrition*; 1989; 40: 275-282.

Iyer U, Mathur G. Nutritional determinants of non communicable diseases in an industrial set up of Vadodara. M.Sc Dissertation thesis, Dept. of Foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda. Baroda, 2007

Iyer U, Pandit N. Assessment of nutritional status and impact of garden cress seeds (*Lepidium sativum*) incorporated khakhra in the management of type 2 diabetes subjects. M.Sc Dissertation thesis, Dept. of Foods and Nutrition, Faculty of Family and Community Sciences, The Maharaja Sayajirao University of Baroda. Baroda, 2007

Iyer U, Shah A. Fruit consumption pattern in Vadodara: Impact of ascorbic acid rich fruit supplementation on the lipid profile of hyperlipidemic subjects. M.Sc Dissertation thesis, Dept. of foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda. Baroda, 2003

Jacobson BH, Aldana SG. Relationship between frequency of aerobic activity and illness-related absenteeism in a large employee sample. *JOEM*; 2001; 43: 1019-1025.

Jahromi MAF, Ray AB. Antihyperlipidemic effect of flavonoids from *Pterocarpus marsupium*. *J Nat Prod*; 1993; 7: 989–994.

Jarald E, Joshi SB, Jain DC. Diabetes and Herbal Medicines. *IJPT*; 2008; 7: 97-106.

Jayaprakasam B, Vareed SK, Olson LK, Nair MG. Insulin secretion by bioactive anthocyanins and anthocyanidins present in fruits. *J Agric Food Chem*; 2005; 53: 28–31.

Jenkins DJA, Kendall CWC, Marchie A cols. Type 2 diabetes and vegetarian diet. *Am J Clin Nutr*; 2003; 78: 610–616.

Jenkins D, Leeds A, Gassull M, Cochet B, Alberti G. Decrease in postprandial insulin & glucose concentrations by guar & pectin. *Annals of Internal Medicine*; 1977; 86: 20–23

Jenkins DJA, Wolever TMS, Taylor RH, et al. Glycemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr*; 1981; 34: 362–6.

Jeppesen J, Hein HO, Suadicani P, Gyntelberg F. Relation of high TG-low HDL cholesterol and LDL cholesterol to the incidence of ischemic heart disease: an 8-year follow-up in the Copenhagen Male Study. *Arterioscler Thromb Vasc Biol*; 1997; 17: 1114–1120.

Johnson IM, Gee JM. Effect of gel-forming gums on the intestinal unstirred layer and sugar transport in vitro. *Gut*; 1981; 22: 398–403.

Johnston KL, Clifford MN, Morgan LM. Coffee acutely modifies gastrointestinal hormone secretion and glucose tolerance in humans: glycemic effects of chlorogenic acid and caffeine. *Am J Clin Nutr*; 2003; 78: 728–733.

Jones PJ, Jew S. Functional food development: Concept to reality. *Trends in Food Science & Technology*; 2007; 18: 387–390.

Joshiyura KJ, Ascherio A, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, Hennekens CH, Spiegelman D, Willett WC. Fruit and Vegetable intake in relation to risk of ischemic stroke. *JAMA*; 1999; 282: 1233 – 1239.

Judd J, Clevidence B, Muesing R, Wittes J, Sunkin M, Podczasy J. Dietary trans fatty acids: effects on plasma lipids and lipoproteins of healthy men and women. *Am J Clin Nutr*; 1994; 59: 861–868.

K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*; 2002; 39: S1–S266.

Kakuda T, Sakane I, Takihara T, Ozaki Y, Takeuchi H, Kuroyanagi M. Hypoglycemic effect of extracts from *Lagerstroemia speciosa* L. leaves in genetically diabetic KK-A^Y mice. *Biosci. Biotechnol. Biochem*; 1996 ; 60 : 204–208.

Kalache A. Ageing: a global perspective. *Community Eye Health*; 1999; 12: 1–4.

Kanakamani J, Ammini AC, Gupta N, Dwivedi SN. Prevalence of microalbuminuria among patients with type 2 diabetes mellitus—a hospital-based study from north India. *Diabetes Technology & Therapeutics*; 2010; 12: 161–166.

Karlsen A, Retterstøl L, Laake P, Paur I, Kjølrsrud-Bøhn S, Sandvik L, Blomhoff R. Anthocyanins inhibit nuclear factor-kappaB activation in monocytes and reduce plasma concentrations of pro-inflammatory mediators in healthy adults. *J Nutr*; 2007; 137: 1951–1954.

Kassie F, Rabot S, Uhl M, Huber W, Qin HM, Helma C et al. Chemoprotective effects of garden cress (*Lepidium sativum*) and its constituents towards 2-amino-3-methylimidazo [4,5-f] quinoline (IQ)-induced genotoxic effects and colonic preneoplastic lesions. *Carcinogenesis*; 2002; 23: 1155-1161.

Katan MB, Grundy SM, Jones P, Law M, Miettinen T, Paoletti R; Stresa Workshop Participants. Efficacy and safety of plant stanols and sterols in the management of blood cholesterol levels. *Mayo Clin Proc*; 2003; 78: 965-978.

Kaur P, Rao TV, Sankarasubbaiyan S, Narayanan AM, Ezhil R, Rao SR, Gupte MD. Prevalence and distribution of cardiovascular risk factors in an urban industrial population in south India: A Cross- Sectional Study. *JAPI*; 2007; 55: 771-776.

Keisewetter H, Jung F, Pindur G, Jung EM, Mrowietz C, Wenzel E. Effect of garlic on thrombocyte aggregation, microcirculation, and other risk factors. *Int J Clin Pharm Ther Tox*; 1991; 29: 151-155.

Kenny SJ, Aubert RE, Giess LS (1995)
Prevalence and incidences of non insulin dependent diabetes mellitus. In: *Diabetes in America*. Ed. National Diabetes Data Group, NIH, Bethesda pp: 47-68

Khan A, Safdar M. Role of diet, nutrients, spices and natural products in diabetes mellitus. *Pakistan Journal of Nutrition*; 2003; 2: 1-12.

Khaw KT, Bingham S, Welch A, Luben R, Wareham N, Oakes S, Day N. Relation between plasma ascorbic acid and mortality in men and women in EPIC-Norfolk prospective study: a prospective population study. *Lancet*; 2001; 357: 657-663.

Kim JI, Kim JC, Kang MJ, Lee MS, Kim JJ, Cha IJ. Effects of pinitol isolated from soybeans on glycaemic control and cardiovascular risk factors in Korean patients with type II diabetes mellitus: a randomized controlled study. *European Journal of Clinical Nutrition*; 2005; 59: 456-458.

Kim JS, Kwon CS, Son KH. Inhibition of Alpha-glucosidase and Amylase by Luteolin, a Flavonoid Biosci. *Biotechnol. Biochem*; 2000; 64: 2458-2461.

King H, Aubert RE, Herman WH. Global burden of diabetes 1995- 2025: Prevalence, numerical estimates, and projection. *Diabetes Care*; 1998; 21: 1414 – 1431.

Klein S, Sheard NF, Pi –Sunyer X, Daly A, Wylie-Rosset J, Kulkarni K, Clark G. Weight management through lifestyle modification for the prevention and management of type 2 diabetes: rationale and strategies. *Diabetes Care*; 2004; 27: 2067- 2073.

Knowler WC, Barret –Connor E, Fowler SE, Hamman RF, Nathan DF. Diabetes prevention program research group: reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *NEJM*; 2002; 346: 393- 403.

Kobayashi Y, Suzuki M, satsu H, Arai S, Hara Y, Suzuki K et al. Green Tea Polyphenols Inhibit the Sodium-Dependent Glucose Transporter of Intestinal Epithelial Cells by a Competitive Mechanism. *J. Agric. Food Chem*; 2000; 48: 5618-5623.

Kocyigit A, Koylu AA, Keles H. Effects of pistachio nuts consumption on plasma lipid profile and oxidative status in healthy volunteers. *Nutr Metab Cardiovasc Dis*; 2006; 16: 202 – 209.

Krauss RM. Lipids and Lipoproteins in Patients With Type 2 Diabetes. *Diabetes Care*; 2004; 27: 1496–1504.

Kulkarni P. Clinical Assessment of the Effect of Sookshma Triphala in Lipoma. *Ayurvedic Resesearch Papers II*, Pune; 1995; 66–71.

Kuppu R, Shrivastava A, Krishnaswami C, Bagavathiammal K, Janaki I, Vijaykumar G et al. Hypoglycemic and hypotriglyceridemic effects of methika churn. *Antiseptic*; 1998; 95: 78-79.

Kuriyama S, Hozawa A, Ohmori K, Suzuki Y, Nishino Y, Fujita K, Tsubono Y, Tsuji I. Joint impact of health risks on health care charges: 7-year follow-up of National Health Insurance beneficiaries in Japan (the Ohsaki Study). *Prev Med*; 2004; 39: 1194-1199.

Kuusisto J, Mykkanen L, Pyorala K, Laakso M. Hyperinsulinemic microalbuminuria: a new risk indicator for coronary artery disease. *Circulation*; 1995; 90: 831–837.

La Vecchia C, Decarli A, Pagano R. Vegetables consumption and risk of chronic disease *Epidemiology*; 1998; 9: 208-210.

Lamarche B, Depres JP, Moorjani S, Cantin B, Dagenais GR, Lupien PJ. Triglycerides and HDL-cholesterol as risk factors for ischemic heart disease: results from the Quebec Cardiovascular Study. *Atherosclerosis*; 1996; 119: 235–245.

Lapidus L, Bengtsson C, Larsson B, Pennert K, Rybo E, Sjostrom L. Distribution of adipose tissue and risk of cardiovascular disease and death: a 12 year follow up of participants in the population study of women in Gothenburg, Sweden. *Br Med J*; 1984; 289: 1257–1261.

Lardinois CK, Starich GH. Polyunsaturated fats enhance peripheral glucose utilization in rats. *J Am Coll Nutr*; 1991; 10: 340 –345.

Larsson B, Svardsudd K, Welin L, Wilhelmsen L, Bjorntorp P, Tibblin G. Abdominal adipose tissue distribution, obesity, and risk of cardiovascular disease and death: 13 year follow up of participants in the study of men born in 1913. *Br Med J*; 1984; 288: 1401–1404.

Lean MEJ, Han TS, Decuenberg P. Predicting body composition by densitometry from simple anthropometric measurements. *Am J Clin Nutr*; 1996; 63: 4 –14.

Lee ET, Lee VS, Lu M, Lee JS, Russel D, Yeh J. Incidence of renal failure in NIDDM: the Oklahoma Indian Diabetes Study. *Diabetes*; 1994; 4: 572-579.

Lele RD, Joshi SR, Gupte A. Association of adipocytokines (Leptin, Adiponectin TNF-alpha), insulin and proinsulin with diabetes – The Mumbai Obesity Project [MOP]. *JAPI*; 2006; 54: 689-696.

Liener IE. Implications of antinutritional components in soybean foods. *Crit Rev Food Sci Nutr*; 1994; 34: 31–67.

Liese A, Mayer Davis E, Haffner S. Development of the insulin resistance syndrome: an epidemiologic perspective. *Epidemiol Rev*; 1998; 20: 157-72.

Lindquist CH, Gower BA, Goran MI. Role of dietary factors in early risk of cardiovascular disease and type 2 diabetes *AJCN*; 2000; 71: 725-732.

Lindström J, Eriksson JG, Valle TT. Prevention of diabetes mellitus in subjects with impaired glucose tolerance in the Finnish Diabetes Prevention Study: results from a randomized clinical trial. *J Am Soc Nephrol*; 2003;14: S108–S113.

Lo GS, Goldberg AP, Lim A, Grundhauser JJ, Anderson C, Schonfeld G. Soy fiber improves lipid and carbohydrate metabolism in primary hyperlipidemic subjects. *Atherosclerosis*; 1986; 62: 239–48.

Look AHEAD Research Group. Reduction in weight and cardiovascular disease risk factors in individuals with type 2 diabetes: one-year results of the Look AHEAD trial. *DiabetesCare*; 2007; 30: 1374–1383.

Loreto Di C, Fanelli C, Lucidi P. Validation of a counselling strategy to promote the adoption and maintenance of physical activity by type 2 diabetic subjects *Diabetes Care*; 2003; 26: 404- 408.

Ludwig DS. Dietary glycaemic index and obesity. *Journal of Nutrition*; 2000; 130: 280S-283S.

Maghrani M, Zeggwagh NA, Michel JB, Eddonks M. Antihypertensive effect of *Lepidium sativum* L. in spontaneously hypertensive rats. *J. Ethnopharmacol*; 2005; 100: 193-197.

Mahan LK and Escott-Stump S. *Krause's Food and Nutrition Therapy*, 12th Edition; 2008. Saunders Elsevier.

Mahdi AA, Chandra A, Singh RK, Shukla S, Mishra LC, Ahmad S. Effect of herbal hypoglycemic agents on oxidative stress and antioxidant status in diabetic rats. *Ind. J. Clin. Biochem*; 2003; 18: 8-15.

Makino H, Haneda M, Babazono T et al. Prevention of transition from incipient to overt nephropathy with telmisartan in patients with type 2 diabetes. *Diabetes Care*; 2007; 30: 1577–8.

Malerbi D, Paiva E, Duarte A, Wajchenberg B. Metabolic effects of dietary sucrose and fructose in type 2 diabetic subjects. *Diabetes Care*; 1996; 19: 1249–1256.

Manach C, Mazur A, Scalbert A. Polyphenols and prevention of cardiovascular disease. *Curr Opin Lipidol*; 2005; 16: 77–84.

Mani U, Bhat S, Shah V, Mehta N. Glycemic index of conventionally used carbohydrate foods. *J Am Coll Nutr*; 1990; 9: 573-77.

Mani U, Gujrati D. Mapping the prevalence of chronic degenerative diseases among free living adult population of urban Vallabh Vidyanagar. M.Sc Dissertation thesis, Dept. of foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda, Baroda, 2002

Mani U, Mani I, Iyer U, Prakash B, Manivannam J, Campbell S et al. Glycemic and lipemic response to various regional meals and south Indian snacks. *Int J Diab Dev Countries*; 1997; 17: 75-79.

Mani U, Pradhan S, Thakkar D, Mehta N, Iyer U, Mani I. Glycemic index of conventional carbohydrate meals. *Br J Nutr*; 1993; 68: 445-50.

Mani U. Glycemic index of cereal pulse mix (diabetic mix) in NIDDM subjects. *Int J Food Sci Nutr*; 1994; 45: 141-45.

Manninen V, Tenkanen L, Koskinen P, Huttunen JK, Mänttäri M, Heinonen OP, Frick MH. Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary heart disease risk in the Helsinki Heart Study: implications for treatment. *Circulation*; 1992; 85: 37-45.

Manson JE, Nathan DM, Krolewski AS. A prospective study of exercise and incidence of diabetes among US male physicians. *JAMA*; 1992; 268: 63-67.

Marchesini G, Forlani G, Cerrelli F. WHO and ATP III proposals for the definition of the metabolic syndrome in patients with Type 2 diabetes. *Diabet Med*; 2004; 21: 383-7.

Marin R, Gorostidi M, Tranche S, Rodriguez-Mañas L, Abellan J. P-444: Hypertension prevalence and control in type 2 diabetes: A primary care-based study. *Am J Hypertens*; 2002; 15: 192A-192A.

Markham KR, Mitchell KA. The mis-identification of the major antioxidant flavonoids in young barley (*Hordeum vulgare*) leaves. *Z Naturforsch C*; 2003; 58: 53-56.

Marshall J, Bessesen D, Hamman R. High saturated fat and low starch and fibre are associated with hyperinsulinaemia in a non-diabetic population: The San Luis Valley Diabetes Study. *Diabetologia*; 1997; 40: 430-438.

Martineau LC, Couture A, Spoor D, Benhaddou-Andaloussi A, Harris C, Meddah B, Leduc C, Burt A, Vuong T, Mai Le P, Prentki M, Bennett SA, Arnason JT & Haddad PS. Anti-diabetic properties of the Canadian lowbush blueberry *Vaccinium angustifolium* Ait. *Phytomedicine*; 2006; 13: 612-623.

Mateos-Aparicio, Cuenca AR, Villanueva-Suárez MJ, Zapata-Revilla MA. Soybean, a promising health source. *Nutr Hosp*; 2008; 23: 305-312.

Mather HM, Chaturvedi N, Kehely AM. Comparison of prevalence and risk factors for microalbuminuria in South Asians and Europeans with type 2 diabetes mellitus. *Diabet Med*; 1998; 15: 672-677.

Mathur P. Risk factors for non-communicable diseases: Getting beyond data. *J Postgrad Med*; 2006; 52: 171-2.

Matos MFD, Souza e Silva NA, Pimenta AJM, Alves da Cunha AJL. Prevalence of Risk Factors for Cardiovascular Disease in Employees of the Research Center at Petrobras. *Arq Bras Cardiol*; 2004; 82: 5-8.

Matsumoto M, Ishigaki F, Ishigaki A, Iwashina H, Hara Y. Reduction of blood glucose levels by tea catechin. *Biosci. Biotechnol. Biochem*; 1993; 57: 525-527.

Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RL. Homeostasis model assessment: Insulin resistance and beta cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*; 1985; 28: 412–419.

McAuley KA, Williams SM, Mann JI, Goulding A, Chisholm A, Wilson N et al. Intensive lifestyle changes are necessary to improve insulin sensitivity. *Diabetes Care*; 2002; 25: 445-452.

McCarty D, Zimmet P. Diabetes 1994 to 2010: Global Estimates and Projections. International Diabetes Institute, Melbourne, 1994

Mehan MB, Kantharia NB, Surabhi S. Risk factor profile of noncommunicable diseases in an industrial productive (25-59 years) population of Baroda. *Int J Diab Dev Ctries*; 2007; 27: 116-121.

Mehan MB, Srivastava N, Pandya H. Profile of non communicable disease risk factors in an industrial setting. *J Postgrad Med*; 2006; 52: 167-71.

Menrad K. Market and marketing of functional food in Europe. *Journal of Food Engineering*; 2003; 56: 181–188.

Messina MJ. Legumes and soybeans: overview of their nutritional profiles and health effects. *Am J Clin Nutr*; 1999; 70: 439-450.

Meyer KA, Kushi LH, Jacobs DR Jr, Slavin J, Sellers TA, Folsom AR. Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr*; 2000; 71: 921–930.

Miller J. Glycemic index/Glycemic Load. *Am J Clin Nutr*; 2001; 76: 5-56.

Mills PR. The development of a new corporate specific health risk measurement instrument, and its use in investigating the relationship between health and well-being and employee productivity. *Environ Health*; 2005; 4:1.

Misra A, Vikram NK. Insulin resistance syndrome (metabolic syndrome) and obesity in Asian Indians: evidence and implications. *Nutrition*; 2004; 20: 482-491.

Miura K et al. Relation of vegetable, fruit and meat intake to 7-y blood pressure change in middle –aged men: the Chicago Western Electric Study. *Am. J. of Epidemiology*; 2004; 159: 572-580 .

Miyake T, Shibamoto T. Inhibition of Malonaldehyde and Acetaldehyde Formation from Blood Plasma Oxidation by Naturally Occurring Antioxidants. *J. Agric. Food Chem.*, 1998, 46 (9), pp 3694–3697

Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA. Indian Herbs and Herbal Drugs Used for the Treatment of Diabetes. *J Clin Biochem Nutr*; 2007; 40: 163–173.

Mohan V and Deepa R (2007). Prevention of diabetes in India. Type 2 Diabetes Mellitus in South East Asians. Ed. Mohan V, Rao G, pp, 331- 343

Mohan V, Deepa M, Deepa R, Shanthirani CS, Farooq S, Ganesan A, et al. Secular trends in the prevalence of diabetes and impaired glucose tolerance in urban South

India—the Chennai Urban Rural Epidemiology Study (CURES-17). *Diabetologia*; 2006; 49: 1175-8.

Mohan V, Deepa M, Farooq S, Prabhakaran D, Reddy KS. Surveillance for risk factors of cardiovascular disease among an industrial population in southern India. *Natl Med J India*; 2008; 21: 8-13.

Mohan V, Mathur P, Deepa R, Deepa M, Shukla DK, Menon GR, Anand K, Desai NG, Joshi PP, Mahanta J, Thankappan KR, Shah B. Urban rural differences in prevalence of self reported diabetes in India-The WHO- ICMR Indian NCD risk factor surveillance. *Diabetes Res Clin Pract*; 2008; 80: 159- 68.

Mohan V, Sandeep S, Deepa R, Shah B, Varghese C. Epidemiology of type 2 diabetes: Indian scenario. *Indian J Med Res*; 2007; 125: 217-230.

Mohan V, Shantirani CS, Deepa R. Glucose intolerance (diabetes and IGT) in selected south Indian population with special reference to family history, obesity and lifestyle factors- the Chennai Urban Population Study (CUPS-14). *J. Assoc. Physicians India*; 2003; 51: 771-777.

Mohan V, Sudha V, Radhika G, Radha V, Rema M, Deepa R. Gene environment interaction and the diabetes epidemic in India. *Forum Nutr*; 2007; 60: 118-26.
Mollet B, Rowland I. Functional foods: At the frontier between food and pharma. *Current Opinion in Biotechnology*; 2002; 13: 483-485.

Montori V, Farmer A, Wollan P, Dinneen S. Fish oil supplementation in type 2 diabetes: a quantitative systematic review. *Diabetes Care*; 2000; 23: 1407-1415.

Moonichoodapa C. Epidemiology and burden of type 2 diabetes mellitus. In: *Type 2 Diabetes: The Indian Scenario, 2002*. Ed. Jayaram BM. Micro Labs Limited., Bangalore, India. Pp, 13-23

Moore LL, VISIONI AJ, Wilson PW, D'Agostino RB, Finkle WD, Ellison RC. Can sustained weight loss in overweight individuals reduce the risk of diabetes mellitus? *Epidemiology*; 2000; 11: 269 -273.

Murakami N, Murakami T, Kodoya M, Matsuda H, Yamahara J, Yoshiyoka M. New hypoglycemic constituents in "gymnemic acid" from *Gymnema sylvestre*. *Chem. Pharm. Bull*; 1996; 44: 469-471.

Mutalik S, Chetana M, Sulochana B, Devi P, Udupa N. Effect of Dianex, a herbal formulation on experimentally induced diabetes mellitus. *Phytother*; 2005; 19: 409-415.

Mykkanen L, Haffner SM, Kuusisto J, Pyorala K, Laakso M. Microalbuminuria precedes the development of NIDDM. *Diabetes*; 1994; 43: 552-57.

Naik G, Priyadarsini K, Hari M. Free radical scavenging reactions and Phytochemical analysis of triphala, an ayurvedic formulation *Current Science*; 2006; 90: 115-119.

National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults 2001 Executive summary of The Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA*; 2001; 285: 2486 -2497.

Neeraja A, Rajyalakshmi P. Hypoglycemic effect of processed fenugreek seeds in humans. *JFST*; 1996; 33: 427-430.

Nelson RW. The role of fiber in managing diabetes mellitus. *Vet. Med*; 1989; 84: 1156-1160.

Niva M. 'All foods affect health': Understandings of functional foods and healthy eating among health-oriented Finns. *Appetite*; 2007; 48: 384-393.

Norata G, Marchesi P, Passamonti S. Anti-inflammatory and antiatherogenic effects of catechin, caffeic acid and trans-resveratrol in apoprotein E deficient mice. *Atherosclerosis*; 2007; 191: 265-271.

The National Kidney and Urologic Diseases Information Clearinghouse (NKUDIC). Kidney Disease of Diabetes Fact Sheet. NIH Publication No. 08-3925; 2008. Available at www.kidney.niddk.nih.gov.

Oberlinner C, Humpert PM, Nawroth PP, Zober A, Morcos M. Metabolic syndrome in a large chemical company: prevalence in a screened worksite sample. *Acta Diabetologica*; 2008; 45: 31-35.

Obrazanek E. Effects on blood lipids of a blood pressure lowering diet: the DASH trial. *AJCN*; 2001; 74: 80-89.

Ohtake H, Nonaka S, Sawada Y, Hagiwara Y, Hagiwara H, Kubota K. Studies on the Constituents of Green Juice from Young Barley Leaves. Effect on Dietarily Induced Hypercholesterolemia in Rats. *Yakugaku Zasshi*; 1985; 105: 1052-1057.

Okosun IS, Cooper RS, Rotimi CN, Osotimehin B, Forrester T. Association of waist circumference with risk of hypertension and type 2 diabetes in Nigerians, Jamaicans, and African - Americans. *Diabetes Care*; 1998; 21: 1836-1842.

Osakabe N, Baba S, Yasuda A, Iwamoto T, Kamiyama M, Takizawa T et al. Daily cocoa intake reduces the susceptibility of low-density lipoprotein to oxidation as demonstrated in healthy human volunteers. *Free Radic Res*; 2001; 34: 93-99.

Osganian SK, Stampfer MJ, Rimm E, Spiegelman D, Hu FB, Manson JE et al. Vitamin C and risk of coronary heart disease in women. *J Am Coll Cardiol*; 2003; 42: 246-252.

Ossman SS. Diabetic Nephropathy: Where We Have Been and Where We Are Going. *Diabetes Spectrum*; 2006; 19: 153-156.

Pan XR, Li GW, HU YH. Effect of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing, IGT and diabetes study *Diabetes Care*; 1997; 20: 537-544.

Pandya HB, Mehan M. Is the industrial city of Baroda prepared to tackle the rising trend of non communicable diseases in workplace setting? M.Sc Dissertation thesis, Dept. of Foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda. Baroda, 2004.

Paranjape A, Mehta A. A Study on Clinical Efficacy of *Lepidium sativum* Seeds in Treatment of Bronchial Asthma. *Iranian Journal of Pharmacology & Therapeutics*; 2006; 5: 55-59.

- Parikh P, Mani U, Iyer U. Role of spirulina in the control of glycemia and lipidemia in Type 2 diabetes mellitus. *Journal of Medicinal Food*; 2001; 4: 193-99.
- Parving HH, Lehnert H, Brochner-Mortensen J, Gomis R, Andersen S, Arner P. The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med*; 2001; 345: 870-878.
- Parving HH, Lewis JB, Ravid M, Remuzzi G, Hunsicker LG. Prevalence and risk factors for microalbuminuria in a referred cohort of type II diabetic patients: A global perspective. *Kidney International*; 2006; 69: 2057-2063.
- Patole AP. Effect of mucilaginous seeds on in vitro rate of starch hydrolysis and blood glucose levels of NIDDM subjects with special reference to garden cress seeds. *J. Med. Aro. Plant Sci*; 1998; 20: 1005-1008.
- Paulíčková I, Ehrenbergerová J, Fiedlerová V, Gabrovská D, Havlová P, Holasová M et al. Evaluation of Barley Grass as a Potential Source of Some Nutritional Substances. *Czech J. Food Sci*; 2007; 25: 65-72.
- Pelletier B, Boles M, Lynch W. Change in health risks and work productivity over time. *J Occup Environ Med*; 2004; 46: 746-754.
- Persaud SJ, Al-Majed H, Raman A, Jones PM. *Gymnema sylvestre* stimulates insulin release in vitro by increasing membrane permeability. *J Endocrinol*; 1999; 163: 207-212.
- Pettitt DJ, Saad MF, Bennett PH, Nelson RG, Knowler WC. Familial predisposition to renal disease in two generations of Pima Indians with type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia*; 1990; 33: 438-443.
- Phillipson B, Rothrock D, Connor W, Harris W, Illingworth D. Reduction of plasma lipids, lipoproteins, and apoproteins by dietary fish oils in patients with hypertriglyceridemia. *N Engl J Med*; 1985; 312: 1210-1216.
- Pickup JC. Inflammation and activated innate immunity in the pathogenesis of type 2 diabetes. *Diabetes Care*; 2004; 27: 813-823.
- Polidori MC, Mecocci P, Stahl W, Parente B, Cecchetti R, Cherubini A et al. Plasma levels of lipophilic antioxidants in very old patients with Type 2 diabetes. *Diab. Metab. Res. Rev*; 2000; 16: 15-19.
- Polidori MC, Stahl W, Eichler O. Profile of antioxidants in human plasma. *Free Radic Biol Med*; 2001; 30: 456-462.
- Polonsky KS, Sturis J, Bell GI. Non-insulin-dependent diabetes mellitus — a genetically programmed failure of the beta cell to compensate for insulin resistance. *N Engl J Med*; 1996; 334: 777-783.
- Prabhakaran D, Shah P, Chaturvedi V, Ramakrishnan L, Manhapra A, Reddy KS. Cardiovascular risk factor prevalence among men in a large industry of northern India. *The Natl Med J India*; 2005; 18: 59-65.
- Prentki M, Nolan CJ. Islet β cell failure in type 2 diabetes. *J. Clin. Invest*; 2006; 116: 1802-1812.

Price KD, Price CSC, Reynolds RD. Hyperglycemia-induced ascorbic acid deficiency promotes endothelial dysfunction and the development of atherosclerosis. *Atherosclerosis*; 2001; 158: 1–12.

Prujm MT, Madeleine G, Riesen WF, Burnier M, Bovet P. Prevalence of microalbuminuria in the general population of Seychelles and strong association with diabetes and hypertension independent of renal markers. *Journal of Hypertension*; 2008; 26: 871-877.

Quintiliani L, Sattelmair J, Sorensen G. The workplace as a setting for interventions to improve diet and promote physical activity. Background paper prepared for the WHO/WEF Joint Event on Preventing Noncommunicable Diseases in the Workplace, 2008

Quio Q, Hu G, Tuomilhto J. Age and sex specific prevalence of diabetes and impaired glucose regulation in impaired glucose regulation in 11 Asian cohorts *Diabetes Care*; 2003; 1770-1780.

Raal F, Kalk J, Lawson M, Esser J, Buys R, Fourie L et al. Effect of moderate dietary protein restriction on the progression of overt diabetic nephropathy, a 6 month prospective study. *Am J Clin Nutr*; 1994; 60: 579-85.

Rai V, Iyer U, Mani U. Effect of tulsi (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels, serum lipids and tissue lipids in diabetic rats. *Plant Foods for Human Nutrition*; 1997; 50: 9-16.

Rainor J. Diabetes 2001 Vital Statistics. Alexandria: American Diabetes Association; 2001:43–74.

Rajala U, Laakso M, Qiao Q, Keinanen-Kiukaanniemi S. Prevalence of retinopathy in people with diabetes, impaired glucose tolerance, and normal glucose tolerance. *Diabetes Care*; 1998; 21: 1664–69.

Ramachandran A, Snehalatha C, Baskar ADS, Mary S, Kumar C, Selvam S et al. Temporal Changes in Prevalence of Diabetes and Impaired Glucose Tolerance Associated With Life Style Transition Occurring in Rural Population in India. *Diabetologia*; 2004; 47: 860–65.

Ramachandran A, Snehalatha C, Vijay V. Low risk threshold for acquired diabetogenic factors in Asian Indians. *Diabetes Res Clin Pract*; 2004; 65: 189-195.

Ramachandran A, Snehalatha C, Vijay V. Temporal changes in prevalence of type 2 diabetes and impaired glucose tolerance in urban southern India. *Diabetes Res Clin Pract*; 2002; 58: 55-60.

Ramachandran A, Snehalatha C, Viswanathan V. Burden of type 2 Diabetes and its complications - The Indian Scenario. *Curr. Sci*; 2002; 83: 1471-1476.

Ramachandran A, Snehlata C, Vijay V. For the Diabetes Epidemiology Study Group in India (DESI). High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey. *Diabetologia*; 2001; 44:1091-1101.

Ramachandran A. Epidemiology of Diabetes in India. In: Type 2 Diabetes Mellitus in South East Asians, 2007. Ed. Mohan V, Rao G, pp, 20-39

Ramachandran A, Mary S, Yamuna A, Murugesan N, Snehlata C. High prevalence of diabetes and cardiovascular risk factors associated with urbanization in India. *Diabetes Care*; 2008; 31: 893-898.

Ramachandran A, Snehlata C, Kapur A. Temporal changes in prevalence of type 2 diabetes and impaired glucose tolerance in urban south India. *Diab. Clin. Res. Prac*; 2002; 58: 55-60.

Ramachandran A. Epidemiology of diabetes in India, three decades of research *Assoc Physic India*; 2005; 53: 34-8.

Rao BN. Bioactive phytochemicals in Indian foods and their potential in health promotion and disease prevention. *Asia Pacific J Clin Nutr*; 2003; 12: 9-22.

Rao T, Sakaguchi N, Juneja L, Wada E, Yokozawa T. Amla (*Emblica officinalis* Gaertn.) Extracts Reduce Oxidative Stress in Streptozotocin-Induced Diabetic Rats. *Journal of Medicinal Food*; 2005; 8: 362-68.

Ravid M, Brosh D, Ravid-Safran D, Levy Z, Rachmani R. Main risk factors for nephropathy in type 2 diabetes mellitus are plasma cholesterol levels, mean blood pressure, and hyperglycemia. *Arch Intern Med*; 1998; 158: 998-1004.

Ravikumar P, Anuradha C. Effect of fenugreek seeds on blood lipid peroxidation and antioxidants in diabetic rats. *Phytother Res*; 1999; 13: 197-201.

Razak F, Anand S, Vuksan V, Davis B, Jacobs R. Ethnic differences in the relationships between obesity and glucose metabolic abnormalities: a cross sectional population study. *Int.J. Obesity*; 2005; 29: 656-667.

Reaven GM. Role of insulin resistance in human disease. *Diabetes*; 1988; 37: 1595-607.

RDA (2009). ICMR, Nutrient requirements and recommended dietary allowances for Indians. Report of expert group: 1978, 1989 and current 2009. Nutrition Society of India. 41st National conference, Material from Pre conference workshop on "Assessment of Nutritional Status and Dissemination of the New RDA for Indians". Pg 12-13.

Reddy KS, Prabhakaran D, Chaturvedi V, Jeemon P, Thankappan KR, Ramakrishnan L et al. on behalf of the Sentinel Surveillance System for Indian Industrial Populations Study Group. Methods for establishing a surveillance system for cardiovascular diseases in Indian Industrial populations. *Bulletin of the World Health Organization*; 2006; 84: 461-469.

Reddy KS, Shah B, Varghese C, Ramadoss A. Responding to the threat of chronic diseases in India. *Lancet*; 2005; online publication. 50140-6736 (05): 67454-5

Renaud S, de Lorgeril M. Wine, alcohol, platelets, and the French paradox for coronary heart disease. *Lancet*; 1992; 339: 1523-1526.

Resnick HE, Valsania P, Halter JB, Lin X. Relation of weight gain and weight loss on subsequent diabetes risk in overweight adults. *J Epidemiol Community Health*; 2000; 54: 596-602.

Retnakaran R, Cull CA, Thorne KI, Adler AI, Holman RR. Risk factors for renal dysfunction in type 2 diabetes. U.K. Prospective Diabetes Study 74. *Diabetes*; 2006; 55: 1832–1839.

Ritz E, Orth SR. Nephropathy in patients with type 2 diabetes mellitus. *N Engl J Med*; 1999; 341: 1127–1133.

Roberfroid MB. An European consensus of scientific concepts of functional foods. *Nutrition*; 2000; 16: 689–691.

Rossi MCE, Nicolucci A, Pellegrini F, Comaschi M, Ceriello A, Cucinotta D et al. Identifying patients with type 2 diabetes at high risk of microalbuminuria: results of the DEMAND (Developing Education on Microalbuminuria for Awareness of reNal and cardiovascular risk in Diabetes) Study. *Nephrol Dial Transplant*; 2008; 23: 1278–1284.

Ruderman N, Chisholm D, Pi-Sunyer X, Schneider S. The metabolically obese, normal-weight individual revisited. *Diabetes*; 1998; 47: 699–713.

Sadikot SM, Nigam A, Das S et al. The burden of diabetes and impaired glucose tolerance in India using the WHO 1999 criteria: prevalence of diabetes in India study (PODIS). *Diabetes Res Clin Pract*; 2004; 66: 301–07.

Sadikot SM, Nigam A, Das S, Bajaj S, Zargar AH, Prasannakumar KM et al. Diabetes India. The burden of diabetes and impaired fasting glucose in India using the ADA 1997 criteria: prevalence of diabetes in India study (PODIS). *Diabetes Res Clin Pract*; 2004; 66: 293–300.

Sato Y, Nagasaki M, Nakai N, Fushimi T. Physical Exercise Improves Glucose Metabolism in Lifestyle-Related Diseases. *Exp. Biol. Med*; 2003; 228: 1208–1212.

Scalbert A, Williamson G. Dietary intake and Bioavailability of Polyphenols. *Journal of Nutrition*; 2000; 130: 2073S–2085S.

Schneider CA, Herold G, Gysan D, Albus C, Bjarnason-Wehrens B, Predel H. Prevalence of Cardiovascular Risk Factors in Middle-Aged Type 2 Diabetic Automobile Employees Assessed at the Workplace-The Praeford Study. *Journal of Occupational and Environmental Medicine*; 2007; 49: 481–483.

Serxner SA, Gold DB, Bultman KK. The impact of behavioral health risks on worker absenteeism. *J Occup Environ Med*; 2001; 43: 347–354.

Seth SD, Sharma B. Medicinal plants of India. *Indian J. Med. Res*; 2004; 120: 9–11.

Shah S, Saikia M, Barman N, Snehlata C, Ramchandran A. High prevalence of Type 2 diabetes in Urban population in North Eastern India. *International Journal of Diabetes in Developing Countries*; 1998; 18: 97–101.

Shamoon H. The relationship between hemoglobin A1c levels and the long term complications of diabetes. *Drug Ben Trends*; 1996; 10–17.

Shane-McWhorter L. Biological complementary therapies: a focus on botanical products in diabetes. *Diabetes Spectrum*; 2001; 14: 199–208.

Shanmugasundaram ERB, Rajeswari G, Baskaran K, Kumar BRR, Shanmugasundaram KR, Ahmath BK. Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus. *J Ethnopharmacology*; 1990; 30: 281–294.

Sharma K, Garg R. Effect of garden cress seeds (*Lepidium sativum*) and niger seeds (*Guizotia abyssinica*) chikki supplementation on hemoglobin levels of anemic young women (17-25 y)- implications for a food based approach to combat anemia. M.Sc Dissertation thesis, Dept. of foods and Nutrition, Faculty of HSc. The Maharaja Sayajirao University of Baroda. Baroda, 2002

Sharma R, Sarkar A, Hazra D, Mishra B, Singh J, Maheshwari B et al. Use of fenugreek seed powder in the management of non insulin dependent diabetes mellitus. *Nutr Res*; 1996; 16: 1331-39.

Sheela CG, Augusti KT. Antidiabetic effects of S-allyl cysteine sulphoxide isolated from garlic *Allium sativum* Linn. *Indian J Exp Biol*; 1992; 30: 523–526.

Sheridan MJ, Cooper JN, Erario M, Cheifetz CE. Pistachio Nut Consumption and Serum Lipid Levels. *Journal of the American College of Nutrition*; 2007; 26: 141-148.

Shichiri M, Kishikawa H, Ohkubo Y, Wake N. Long term results of the Kumamoto study on optimal diabetes control in type 2 diabetic patients. *Diabet Care*; 2000; 23: B21 – 29.

Shih-Te T, Shun-Jen C, Jung-Fu C, Kai-Jen T, Jeng-Yueh H, Hung-Chun C et al. Prevention of Diabetic Nephropathy by Tight Target Control in an Asian Population With Type 2 Diabetes Mellitus. A 4-Year Prospective Analysis. *Arch Intern Med*; 2010; 170: 155-161.

Sicree R, Shaw J, Zimmet P. Prevalence and projections. In *Diabetes Atlas*. 3rd ed. Brussels, Belgium, International Diabetes Federation; 2006; pg 16–104.

Singh P. Mycotoxin elaboration in triphala and its constituents. *Indian Phytopathol*; 2003; 56: 380–383.

Singh RB et al. Effect of fat –modified and fruit –and vegetable –enriched diets on blood lipids in the Indian Diet Heart Study. *Am. J. Cardio*; 1992; 70: 869-874.

Singh RB. Can guava fruit intake decrease blood pressure and blood lipids? *J. Hum. Hypertens*; 1993; 7: 33- 38.

Singhal S, Gupta R, Goyle A. A comparison of antioxidant efficacy of vitamin E, Vitamin C, Vitamin A and fruits in coronary heart disease: A controlled trial. *The Journal of the Association of Physicians of India*; 2001; 49: 327-31.

Sitprija S, Plengvidhya C, Kangkaya V, Bhuvapanich S, Tunkayoon M. Garlic and diabetes mellitus phase II clinical trial. *J Med Assoc Thai*; 1987; 70: 223–227.

Snijder MB, Dekker JM, Visser M, Bouter LM, Stehouwer CDA, Kostense PJ et al. Associations of hip and thigh circumferences independent of waist circumference with the incidence of type 2 diabetes: the Hoorn Study. *AJCN*; 2003; 77: 1192-1197.

Srivastava Y, Venkatakrishna-Bhatt H, Verma Y, Venkaiah K, Raval BH. Antidiabetic and adaptogenic properties of *Momordica charantia* extract: An experimental and clinical evaluation. *Phytother. Res*; 1993; 7: 285-289.

Stephens NG, Parsons A, Schofield PM. Randomised controlled trials of Vitamin E in patients with coronary disease CHAOS (Cambridge Heart Antioxidant Study). *Lancet*; 1996; 347: 781-786.

Stevens J, Ahn K, Juhaeri, Houston D, Steffan L, Couper D. Dietary fiber intake and glycemic index and incidence of diabetes in African-American and white adults: the ARIC study. *Diabetes Care*; 2002; 25: 1715-1721.

Storlien LH, Jenkins AB, Chisholm DJ, Pascoe WS, Khouri S, Kraegen EW. Influence of dietary fat composition on development of insulin resistance in rats. Relationship to muscle triglyceride and omega-3 fatty acids in muscle phospholipid. *Diabetes*; 1991; 40: 280-289.

Stratton IM, Adler AI, Neil HAW, Mathews DR, Manley SE, Gull CA et al. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS-35): prospective observational study. *BMJ*; 2000; 221: 405-412.

Stratton IM, Kohner EM, Aldington SJ, et al. UKPDS 50: risk factors for incidence and progression of retinopathy in type II diabetes over 6 years from diagnosis. *Diabetologia*; 2001; 44: 156-63.

Suraniti S, Bled F, Girault A, Fressinaud P, Marre M. Serum lipids and urinary albumin excretion in non-insulin-dependent diabetics. *Mol Cell Biochem*; 1992; 109: 197-200.

Suryanarayana P, Anil Kumar P, Saraswat M, Petrash JM, Reddy GB. Inhibition of aldose reductase by tannoid principles of *Emblica officinalis*: Implications for the prevention of sugar cataract. *Molecular Vision*; 2004; 10:148-54.

Swinburn BA. Effect of dietary lipid on insulin action. Clinical studies. *Ann N Y Acad Sci*; 1993; 683: 102-109.

Taskiran M, Iversen A, Klausen K, Jensen GB, Jensen JS. The association of microalbuminuria with mortality in patients with acute myocardial infarction. A ten-year follow-up study. *Heart International*; 2010; 5:

Taylor R, Ram P, Zimmet P. Physical activity and prevalence of diabetes in Melanesian and Indian men in Fiji. *Diabetologia*; 1984; 27: 578-582.

Tenkanen L, Sjöblom T, Härmä M. Joint effect of shift work and adverse life-style factors on the risk of coronary heart disease. *Scand J Work Environ Health*; 1998; 24: 351-357.

Thankappan KR, Shah B, Mathur P, Sarma PS, Srinivas G, Mini GK et al. Risk factor profile for chronic non-communicable diseases: Results of a community-based study in Kerala, India. *Indian J Med Res*; 2010; 131: 53-63.

Tiwari AK and Rao JM. Diabetes Mellitus and multiple therapeutic approaches of phytochemicals: present status and future prospects. *Current Science*; 2002; 83: 30-38.

Tiwari AK. Invigorated barley in diabetes. *Current Science*; 2008; 95: 25-29.

Tobe SW, McFarlane PA, Naimark DM. Microalbuminuria in diabetes mellitus. *CMAJ*; 2002; 167: 499–503.

Tomaru M, Takano H, Osakabe N, Yasuda A, Inoue K, Yanagisawa R et al. Dietary supplementation with cacao liquor proanthocyanidins prevents elevation of blood glucose levels in diabetic obese mice. *Nutrition*; 2007; 23: 351-5.

Track N, Lar V, Chur S. Guar by product improves carbohydrate tolerance in healthy human subjects. *Diabetes Res Clin Pract*; 1985; 115-119.

Tsai SP, Donnelly RP, Wendt JK. Obesity and Mortality in a Prospective Study of a Middle-Aged Industrial Population. *J Occup Environ Med*; 2006; 48: 22-27.

Tsai CE, Teng L, Lin Y. Effect of Young Barley Leaf Essence on Serum Lipids in Hamsters. *Nutr Sci J*; 2005; 30: 21~27.

Tsai SP, Wendt JK, Cardarelli KM, Fraser AE. A mortality and morbidity study of refinery and petrochemical employees in Louisiana. *J Occup Environ Med*; 2003; 60: 627–633.

Tsuda T, Horio F, Uchida K, Aoki H, Osawa T. Dietary cyanidin 3-O-beta-D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. *J Nutr*; 2003; 133: 2125–2130.

Tuomilhto J, Lindstorm J, Eriksson JJ (The Finnish Diabetes Prevention Study). Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *New Eng J Med*; 2002; 344: 1343-1350.

UK Prospective Diabetes Study Group. Intensive blood glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes : UKPDS 33. *Lancet*; 1998; 352: 837-53.

Unnikrishnan R, Rema M, Pradeepa R, Deepa M, Shanthirani CS, Deepa R et al. Prevalence and Risk Factors of Diabetic Nephropathy in an Urban South Indian Population. The Chennai Urban Rural Epidemiology Study (CURES 45). *Diabetes Care*; 2007; 30: 2019–2024.

Valabhji J, Mc Coll AJ, Richmond W, Schachter M, Rubens MB, Elkeles RS. Total antioxidant status and coronary artery calcification in type 1 diabetes. *Diabetes Care*; 2001; 24: 1608–1613.

Valsa AK, Sudheesh S, Vijayalakshmi NR. Effect of catechin on carbohydrate metabolism. *Indian J. Biochem. Biophys*; 1997; 34: 406–408.

Van den Heuvel SG, Boshuizen HC, Hildebrandt VH, Blatter BM, Ariëns GA, Bongers PM. Effect of sporting activity on absenteeism in a working population. *Br J Sports Med*; 2005; 39: e15.

van't Veer P, Jansen MCJF, Klerk M, Kok FJ. Fruits and vegetables in the prevention of cancer and cardiovascular disease. *Public Health Nutrition*; 2000; 3: 103–107.

Varghese A, Deepa R, Rema M, Mohan V. Prevalence of microalbuminuria in type 2 diabetes mellitus at a diabetic center in South India. *Post-Grad Med J*; 2001; 77: 399–402.

Vasan RS, Larson MG, Leip EP, et al. Impact of high-normal blood pressure on the risk of cardiovascular disease. *N Engl J Med*; 2001; 345: 1291–7.

Vedavanam K, Srijayanta S, O'Relly J, Raman A. Antioxidant action and potential antidiabetic properties of an isoflavonoid-containing soyabean phytochemical extract (SPE). *Phytother. Res*; 1999; 13: 601–608.

Viberti G, Wheeldon NM. Microalbuminuria reduction with valsartan in patients with type 2 diabetes mellitus: a blood pressure-independent effect. *Circulation*; 2002; 106: 672–678.

Vijay V, Snehalatha C, Shina K, Lalitha S, Ramachandran A. Familial aggregation of diabetic kidney disease in Type 2 diabetes in South India. *Diabetes Res Clin Pract*; 1999; 43: 167–171.

Vijayakumar G, Arun R, Kutty VR. High Prevalence of Type 2 Diabetes Mellitus and Other Metabolic Disorders in Rural Central Kerala. *JAPI*; 2009; 57: 563–567.

Vikram NK, Tandon N, Misra A, Srivastava MC, Pandey RM, Mithal A et al. Correlates of type 2 diabetes mellitus in children, adolescents and young adults in north India: a multisite collaborative case-control study. *Diabet Med*; 2005; 23:293–298.

Vinson AJ, Jang J. Invitro and invivo lipoprotein antioxidant effect of a citrus extract and ascorbic acid on normal and hypercholesterolemic human subjects. *J Med Food*; 2001; 4: 187–192.

Vinson AJ, Su X, Zabik L, Bose P. Phenol antioxidant quality and quantity in foods: Fruits. *J. Agric Food Chem*; 2001; 49: 5315–5321.

Vinson JA, Mandarano MA, Shuta DL, Bagchi M, Bagchi D. Beneficial effects of a novel IH636 grape seed proanthocyanidin extract and a niacin-bound chromium in a hamster atherosclerosis model. *Molecular and Cellular Biochemistry*; 2002; 240: 99–103.

Viswanathan V, Snehalatha C, Terin Mathai, Muthu Jayaraman, Ramachandran A. Cardiovascular morbidity in proteinuric South Indian NIDDM patients. *Diabetes Res Clin Pract*; 1998; 39: 63 - 67.

Viswanathan V. Prevention of diabetic nephropathy: A diabetologist's perspective. *Indian J Nephrol*; 2004; 14: 157–162.

Vivancos M, Moreno JJ. Effect of resveratrol, tyrosol and β -sitosterol on oxidised low-density lipoprotein-stimulated oxidative stress, arachidonic acid release and prostaglandin E₂ synthesis by RAW 264.7 macrophages. *British Journal of Nutrition*; 2008; 99: 1199–1207.

Vuksan V, Sievenpiper J, Owen R, Swilley J, Spadafora P, Jenkins D et al. Beneficial effects of viscous dietary fibre from konjac-mannan in subjects with the insulin resistance syndrome. *Diabetes Care*; 2000; 23: 9–14.

Waddington E, Puddey IB, Croft KD. Red vine polyphenolic compounds inhibit atherosclerosis in apoprotein E deficient mice independently of effects on lipid peroxidation. *Am J Clin Nutr*; 2004; 79: 54–61.

Wang Z, Hoy WE. Albuminuria as a marker of the risk of developing type 2 diabetes in non-diabetic Aboriginal Australians. *International Journal of Epidemiology*; 2006; 35: 1331–1335.

Watanabe J, Kawabata J, Kurihara H, Niki R. Isolation and identification of α -glucosidase inhibitors from tochu-cha. *Biosci. Biotech. Biochem*; 1997; 61: 177-178.

Welhinda J, Karunanayake E, Sheriff M, Jayasinghe K. Effect of *Momordica charantia* on glucose tolerance in maturity onset diabetes. *J Ethanopharmacology*; 1986; 17: 277-282.

Welsh CA, Lachance PA, Wasserman BP. Dietary Phenolic Compounds: Inhibition of Na^+ -Dependent D-Glucose Uptake in Rat Intestinal Brush Border Membrane Vesicles *J. Nutr*; 1989; 119: 1698–1704.

Weng X, Liu Y, Wang W, Yang G, Caballero B. Use of body mass index to identify obesity –related metabolic disorders in the Chinese population. *Eur. J. Clin. Nutr*; 2006 ; 60: 931-937.

Whincup PH, Gilg JA, Owen CG, Odoki K, Cook DG. British south Asians aged 13-16 years have higher fasting glucose and insulin levels than Europeans. *Diab Met*; 2005; 22: 1275- 1277.

Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 200 and projections for 2030. *Diabetes Care*; 2004; 27: 1047-53.

Willett S, Stampfer M, Hu F, Franz M, Sampson L, Hennekens C et al. A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *American Journal of Clinical Nutrition*; 2000; 71: 1455-61.

Willett W, Manson J, Simin L. Glycemic index, glycemic load and risk of type 2 diabetes. *American journal of Clinical Nutrition*; 2002; 76: 274S-80S.

Wilson PWF, D'Agostino RB, Parise H, Sullivan L, Meigs JB. Metabolic Syndrome as a Precursor of Cardiovascular Disease and Type 2 Diabetes Mellitus. *Circulation*; 2005; 112: 3066-3072.

Wirta OR, Pasternack AI, Mustoren JT, Koivula TA, Harmoinen A. Urinary albumin excretion rate and its determinants after six years in non-insulin-dependent diabetic patients. *Nephrol Dial Transplant*; 1996; 11: 449-456.

World Health Organization / World Economic Forum. Preventing Noncommunicable Diseases in the Workplace through Diet and Physical Activity. WHO/World Economic Forum Report of a Joint Event, 2008

World Health Organization. Noncommunicable Diseases. About NCDs. Available from: http://www.searo.who.int/en/Section1174/Section1459_7409.htm

World Health Organization. Global Strategy on Diet, Physical Activity and Health. Geneva, World Health Organization, 2004.

World Health Organization. Preventing chronic diseases: a vital investment, 2005

World Health Organization. The World Health Report, Reducing Risks, Promoting Healthy Life, 2002

Wright E, Scism-Bacon L, Glass C. Oxidative stress in type 2 diabetes: the role of fasting and postprandial glycaemia. *Int J Clin Pract*; 2006; 60: 308-314.

Wu YT, Kong NCT, de Leon FA, Pan CY, Tai TY, Yeung VTF, Yoo SJ, Rouillon A, Weir MR. An alarmingly high prevalence of diabetic nephropathy in Asian type 2 diabetic patients: the MicroAlbuminuria Prevalence (MAP) Study. *Diabetologia*; 2005; 48: 17-26.

Yeh GY, Eisenberg DM, Kaptchuk TJ, Phillips RS. Systematic Review of Herbs and Dietary Supplements for Glycemic Control in Diabetes. *Diabetes Care*; 2003; 26: 1277-1294.

Yoshikawa M, Murakami T, Harada E, Murakami N, Yamahara J, Matsuda H. Bioactive Saponins and Glycosides. VII. On the Hypoglycemic Principles from the Root Cortex of *Aralia elata* SEEM: Structure Related Hypoglycemic Activity of Oleanolic Acid Oligoglycoside. *Chemical & Pharmaceutical Bulletin* 1996, 44, 1923-1927.

Yoshikawa M, Murakami T, Kadoya M, Li Y, Murakami N, Yamahara J, Matsuda H. Medicinal Foodstuffs. IX. The inhibitors of glucose absorption from the leaves of *Gymnema Sylvestre* R. BR. (Asclepiadaceae): Structures of Gymnemosides a and b *Chemical & Pharmaceutical Bulletin*; 1997; 45: 1671-1676.

Yoshikawa M, Shimada H, Morikawa T, Yoshizumi S, Matsumura N, Murakami T et al. J. Medicinal Foodstuffs. VII. On the Saponin Constituents with Glucose and Alcohol Absorption-Inhibitory Activity from a Food Garnish "Tonburi", the Fruit of Japanese *Kochia scoparia* (L.) SCHRAD: Structures of Scoparianosides A, B, and C. *Chemical & Pharmaceutical Bulletin*; 1997; 45: 1300-1305.

Yoshikawa M, Shimada H, Nishida N, Li Y, Toguchida I, Yamahara J et al. Antidiabetic Principles of Natural Medicines. II. Aldose Reductase and α -Glucosidase Inhibitors from Brazilian Natural Medicine, the Leaves of *Myrcia multiflora* DC. (Myrtaceae) : Structures of Myrciacitrins I and II and Myrciaphenones A and B. *Chem. Pharm. Bull*; 1998; 46: 113-119.

Youdim KA, McDonald J, Kalt W, Joseph JA. Potential role of dietary flavonoids in reducing microvascular endothelium vulnerability to oxidative and inflammatory insults (small star, filled). *J Nutr Biochem*; 2002; 13: 282-288.

Yu YM, Chang WC, Chang CT, Hsieh CL, Tsai CE. Effects of young barley leaf extract and antioxidative vitamins on LDL oxidation and free radical scavenging activities in type 2 diabetes. *Diabetes Metab*; 2002; 28: 107-114.

Yu YM, Chang WC, Liu CS, Tsai CM. Effect of young barley leaf extract and adlay on plasma lipids and LDL oxidation in hyperlipidemic smokers. *Biol Pharm Bull*; 2004; 27: 802-805.

Yu YM, Tsai CE. LDL cholesterol and oxidation are significantly reduced in Type 2 diabetic patients receiving a barley leaf essence supplemented olive oil diet. *Food science and agricultural chemistry*; 2003; 5: 1-6.

Yu YM, Wu CH, Tseng YH, Tsai CE, Chang WC. Antioxidative and hypolipidemic effects of barley leaf essence in a rabbit model of atherosclerosis. *Jpn J Pharmacol*; 2002; 89: 142-148.

Yuan Z, He P, Cui J, Takeuchi H. Hypoglycemic Effect of Water-soluble Polysaccharide from *Auricularia auricula-judae* Quel. on Genetically Diabetic KK-A^y Mice. *Biosci. Biotechnol. Biochem*; 1998; 62: 1898–1903.

Zimmet PZ. Diabetes epidemiology as a tool to trigger diabetes research and care. *Diabetologia*; 1999; 42: 499-518.

<http://da3.diabetesatlas.org/indexd9bd.html>

www.drugs.com

www.theolife.com

http://www.idf.org/webdata/docs/Underlying_pathophysiology.ppt#436,1,Targeting the Underlying Pathophysiology of Type 2 Diabetes