

## List of Figures

	Description	Page No
<b>Chapter 1</b>		
<b>Figure 1</b>	Main nitrogen assimilation pathways in cyanobacteria	7
<b>Figure 2</b>	Nitrogen regulation of cyanobacteria: Interconnections between the regulatory proteins NtcA and PII	9
<b>Figure 3</b>	Heterocyst metabolism and nitrogen fixation.	15
<b>Figure 4</b>	The filamentous heterocystous cyanobacterium <i>Anabaena</i> PCC 7120 grown with (N+) or without (N-) combined nitrogen	15
<b>Figure 5</b>	Filaments of <i>Anabaena</i> PCC 7120 containing a plasmid carrying a <i>patS-gfp</i> reporter 27 hours after nitrogen step-down	19
<b>Chapter 2</b>		
	Schematic representation of the construction of N terminal of <i>xisA</i> gene	51
<b>Chapter 3</b>		
<b>Figure 1</b>	Mechanisms of sensing N status in <i>E. coli</i>	65
<b>Chapter 4</b>		
<b>Figure 1</b>	Restriction digestion pattern of pMX25 (a) and pMX32 (b) plasmids by EcoR1	78
<b>Figure 2a</b>	Restriction digestion pattern of <i>E. coli</i> JM101 containing pMX25 and pAM1323 plasmids by EcoR1	78
<b>Figure 2b</b>	Restriction digestion pattern of <i>E. coli</i> JM101 containing pMX25 and pAM1323 plasmids	79
<b>Figure 2c</b>	Restriction digestion pattern of <i>E. coli</i> JM101 containing pMX32 and pAM1323 plasmids	79
<b>Figure 3</b>	Excision of <i>nifD</i> element of <i>Anabaena</i> PCC 7120 in <i>E. coli</i> JM101 containing pMX25 and pAM1323 plasmids	80
<b>Figure 4</b>	Restriction digestion pattern of pKK1 and pAM1323 plasmids by EcoR1	80
<b>Figure 5</b>	$\beta$ -galactosidase activity of <i>PxisA-lacZ</i> reporter fusion in <i>E. coli</i> JM101 containing <i>ntcA</i> gene in the stationary phase of Luria broth and M9 minimal medium	81

<b>Figure 6</b>	Isolation of genomic DNA of <i>Anabaena</i> sp. strain PCC 7120	81
<b>Figure 7</b>	Amplification of <i>ntcA</i> gene of <i>Anabaena</i> sp. Strain PCC 7120	82
<b>Figure 8</b>	Map of plasmid (A) pACYC184 and (B) pTZ57R.	82
<b>Chapter 5</b>		
<b>Figure A</b>	Protein domain structure of $\lambda$ integrase	93
<b>Figure 1</b>	Amplification of the N terminal domain of the <i>xisA</i> gene	94
<b>Figure 2</b>	Restriction digestion of T vector containing N terminal of <i>xisA</i> gene	94
<b>Figure 3</b>	Restriction digestion of the pNTMCGm plasmid	94
<b>Figure 4</b>	Restriction digestion of <i>E. coli</i> DH5 $\alpha$ transformants	95
<b>Figure 5</b>	Amino acid sequence encoded by N terminal of <i>xisA</i> gene in pTZ57R and pNTMCGm plasmid	95
<b>Figure 6</b>	Expression of N-terminus Protein in <i>E. coli</i> DH5 $\alpha$ in LB	96
<b>Chapter 6</b>		
<b>Figure 1</b>	Amplification of <i>xisA</i> without promoter	100
<b>Figure 2</b>	Colony PCR of <i>xisA</i> without promoter	100
<b>Figure 3</b>	Digestion of pMalp2 and pMalp2-Xis with HindIII and ScaI	101
<b>Figure 4</b>	Protein profile of total cell extract of <i>E. coli</i> pMALp2-Xis grown in Luria Broth at 37°C	101
<b>Figure 5</b>	Protein profile of periplasmic extract of <i>E. coli</i> pMALp2-Xis and pMalp2 grown in Luria Broth at 37°C	102
<b>Figure 6</b>	Protein profile of total cell extract of <i>E. coli</i> pMALp2-Xis and pMalp2 grown in 100 ml Luria Broth at 25°C	103
<b>Figure 7</b>	Protein profile of periplasmic extract of <i>E. coli</i> pMALp2-Xis and pMalp2 grown in 100 ml Luria Broth at 25°C	103
<b>Figure 8</b>	Protein profile of TCA precipitated periplasmic extract of <i>E. coli</i> pMALp2-Xis and pMalp2 grown in 100 ml Luria Broth at 25°C	104
<b>Figure 9</b>	Protein profile of TCA precipitated Total cell extract of <i>E. coli</i> pMALp2-Xis and pMalp2 grown in 100 ml Luria Broth at 25°C	105

## Chapter 7

<b>Figure 1</b>	Cloning of <i>xisA</i> in (a) pET20b(+) and (b) pVN6	109
<b>Figure 2</b>	Amplification of the <i>xisA</i> gene	110
<b>Figure 3</b>	HindIII digestion of plasmids containing <i>xisA</i> gene	111
<b>Figure 4</b>	BamHI Digestion of pVN6 <i>xisA</i> plasmid	111
<b>Figure 5</b>	HindIII digestion of pVN6 <i>xisA</i>	111
<b>Figure 6</b>	HindIII digestion of pET20b(+) and pET <i>xisA</i> plasmids	112
<b>Figure 7</b>	SDS PAGE profile of expression of <i>xisA</i> gene using regulatory circuit	113
<b>Figure 8</b>	Induction of the <i>xisA</i> gene expression in dicistronic form with C protein of the Mu phage under regulatory circuits in <i>E. coli</i> BL26 pNC1 pVN6 <i>xisA</i>	113
<b>Figure 9</b>	Induction of the <i>xisA</i> gene expression in dicistronic form with C protein of the Mu phage under regulatory circuits in <i>E. coli</i> BL26 pNC1 pVN6 <i>xisA</i>	114
<b>Figure 10</b>	<i>xisA</i> gene expression form BL26 pNC1 & pVN6 <i>xisA</i>	115
<b>Figure 11</b>	Activity assay of XisA protein on substrate plasmid pBR322NG1 using low salt buffer of the NEB	115
<b>Figure 12</b>	Phosphocellulose column purification fractions of BL26 pNC1 pVN6 <i>xisA</i> .	116
<b>Figure 13</b>	XisA expression using different combination of plasmid in BL21 (DE3) using plasmid pLysE or without pLysE	117
<b>Figure 14</b>	Expression of XisA using BL21 harboring pLysE pVN6 <i>xisA</i> after 5 h of induction with 0.5 mM final concentration of IPTG	118
<b>Figure 15</b>	Protein expression profile of the sonicated supernatant and pellet sample of BL21 pLysE pET <i>xisA</i>	118
<b>Figure 16</b>	PEI precipitation in presence of different NaCl concentration BL21 pLysE pET <i>xisA</i>	119
<b>Figure 17</b>	Different ammonium sulphate precipitation of PEI cut samples of BL21 pLysE and pET <i>xisA</i>	120
<b>Figure 18</b>	Protein purification using affinity column chromatography where phosphocellulose as column material	121
<b>Figure 19</b>	Heparin column purified early fractions	121
<b>Figure 20</b>	Mono Q fractions loaded on the 7% SDS-PAGE	122