LIST OF TABLES

Table-1:	Recoveries of ciprofloxacin HCl from ciprofloxacin ophthalmic solution	72
Table-2:	Recoveries of ciprofloxacin HCl from ciprofloxacin ophthalmic gel	72
Table-3:	Recoveries of ciprofloxacin HCl from ciprofloxacin ophthalmic ointment	74
Table-4:	Recoveries of ciprofloxacin from Cip+Dexa suspension	79
Table-5:	Recoveries of ciprofloxacin from Cip+Dexa solution	79
Table-6:	Recoveries of dexamethasone from Cip+Dexa suspension	79
Table-7:	Recoveries of dexamethasone from Cip+Dexa solution	80
Table-8:	Results of the recovery studies of benzalkonium chloride from various formulations	81
Table-9:	Results of the standardization of ciprofloxacin HC1	84
Table-10:	Results of the standardization of dexamethasone	87
Table-11:	Results of the standardization of benzalkonium chloride	.89
Table-12:	Results of the standardization of other additives	91
Table-13:	Recovery studies of ciprofloxacin in the presence of various additives and the pH of the resulting solutions	95
Table-14:	Results of the recovery studies of ciprofloxacin in the presence of various polymers and the pH of the resulting solutions	97
Table-15:	Results of the evaluation of sterilizing CipHCl by exposure to EtO gas	103
Table-16:	Results of the evaluation of sterilizing dexamethasone at 140°C <i>in vacuo</i> and in air	107

.

,

Table-17:	Results of the assay and measurement of pH of CipHCl solution, before and after sterilization	15
Table-18:	Results of the liquefaction and gelling behaviour of gels prepared with various concentrations of poloxamer	28
Table-19:	Results of the initial drug content of the various preparations	36
Table-20:	Results of the initial assay (as % of the label claim) of BKC from various preparations	37
Table-21:	Results of the particle-size distribution of CipHCl powder, dexamethasone powder as well as Dexamethasone in suspension	41
Table-22:	Results of pH measurements of the preparation, on the day of their manufacture	42
Table-23:	Viscosity measurements of Cip. Gel at 25°C, at different spindle speeds	43
Table-24:	T ₉₀ values obtained from the release profile drug from of the gel at different polymer concentrations	46
Table-25:	Accelerated stability data: Content of ciprofloxacin in Cip. solution 1	49
Table-26:	Accelerated stability data: Content of ciprofloxacin in Ciplox ^R	50
Table-27:	Accelerated stability data: Content of ciprofloxacin in Cip. gel (in vial) 1	51
Table-28:	Accelerated stability data: Content of ciprofloxacin in Cip. Gel (in ointment tube) 1	52
Table-29:	Accelerated stability data: Content of ciprofloxacin in Cip. Dintment 1	53
Table-30:	Accelerated stability data: Content of ciprofloxacin in Cip+Dexa solution 1	54
Table-31:	Accelerated stability data: Content of dexamethasone in Cip+Dexa solution 1	55
Table-32:	Accelerated stability data: Content of ciprofloxacin in Cip+Dexa suspension. 1	56

.

~

Table-33:	Accelerated stability data: Content of dexamethasone in Cip+Dexa suspension.	156
Table-34:	Accelerated stability data: Assay of benzalkonium chloride (% of initial) from various preparations, at the end of 6 months of accelerated stability studies	157
Table-35:	Accelerated stability data: pH of the preparations at different times during accelerated stability studies	158
Table-36:	Accelerated stability data: Data obtained from the <i>in-vitro</i> release profile of ciprofloxacin from the gel: initial (zero day) values and those obtained after storage for 6 months at different temperatures	159
Table-37:	Accelerated stability data: Viscosity of the gel (in vial) after the completion of 6 months of accelerated stability studies	161
Table-38:	Accelerated stability data: Particle size distribution of dexamethasone, initially and after storage at 5°C and 25°C for a period of 6 months	162
Table-39:	Results of the preservative effectiveness test	167
Table-40:	Treatment protocol for efficacy study of Cip+Dexa formulations	174
Table-41:	Dosing and sampling protocol for determining the aqueous humour drug concentration - time profile	178
Table-42:	Dosing and sampling protocol for determination of the aqueous humour drug concentration - time profile following application of multiple doses of Cip.solution and single dose of Cip. gel	180
Table-43:	Dosing and sampling protocol for determination of the aqueous humour drug concentration - time profile obtained after application of a single dose of Cip. ointment and multiple doses of Cip. solution	181
Table-44:	Results of the pharmacokinetic values obtained for different dosage forms	186