

B
I
B
L
I
O
B I B L I O G R A P H Y
R
A
P
H
Y

BIBLIOGRAPHY

Adibi, S.A. (1971).

Alteration in the urinary excretion rate of aminoacids and nitrogen by dietary means in obese and normal human subjects. J. Lab. Clin. Med., 77, 278-289.

Aksu, A., Morrow, G. III, and Barness, L.A. (1974).

Urinary excretion of non-nitrogenous organic acids by healthy infants and children. Clin. Chem., 20, 603-605.

Albanese, A.A. and Irby, V. (1944).

Determination of urinary amino nitrogen by the copper method. J. Biol. Chem., 153, 583-588.

Albanese, A.A. and Orto, L.A. (1967).

Urinary excretion of amino acids. In 'Newer Methods of Nutritional Biochemistry' (A.A. Albanese, Ed.), Academic Press, New York, Vol. III, p. 22.

Albanese, A.A. and Wangerin, D.M. (1944).

The creatine and creatinine excretion of normal adult males. Science, 100, 58.

Alleyne, G.A.O. (1968).

Studies on total body potassium in infantile malnutrition. The relation to body fluid spaces and urinary creatinine. Clin. Sci., 34, 199-209.

Alleyne, G.A.O., Viteri, F.E. and Alvarado, J. (1970).

Indices of body composition in infantile malnutrition : total body potassium and urinary creatinine. Am. J. Clin. Nutr., 23, 875-878.

Altman, P.L. and Dittmer, D.S. (1974).

Biology Data Book, Federation of American Societies for
Experimental Biology, Bethesda, Maryland, Vol. III.

Anasuya, A. and Rao, B.S.N. (1966a).

Urinary hydroxyproline in kwashiorkor. Ind. J. Med. Res.,
54, 849-856.

Anasuya, A. and Rao, B.S.N. (1966b).

Urinary hydroxyproline in kwashiorkor. Lancet, 1, 94-95.

Anon, (1971).

Creatinine height index in malnourished children.

Nutr. Revs., 29, 134-137.

Annual Report of National Institute of Nutrition (1973).

Hyderabad, India.

Applegarth, D.A. and Ross, P.M. (1975).

The unsuitability of creatinine excretion as a basis for
assessing the excretion of other metabolites by infants
and children. Clin. Chim. Acta., 64, 83-85.

Araya, J., Araya, H. and Tagle, M.A. (1974).

Excretion of endogenous-N in faeces by the rat.

Arch. Latinoam. Nutri., 24, 97-106.

cf : Nutr. Abst. Revs., 45, 4202 (1975).

Arima, M., Okazaki, R. and Kitaharo, T. (1977).

The cycle AMP/creatinine ratio in random urine samples
of children. Igaku. No Ayumi, 100, 677-678.

cf : Chem. Abstr., 86, 168638 (1977), p. 334.

Arroyave, G. (1961).

Biochemical evaluation of nutritional status in man.

Fed. Proc., 20, (Suppl. 7), 39-49.

Arroyave, G. (1962).

Biochemical signs of mild-moderate forms of protein-calorie malnutrition. In 'Mild-moderate forms of protein-calorie Malnutrition'. Symp. Swedish Nutr. Found. I, Bastad, Uppsala, Almqvist and Wiksell, p. 32.

Arroyave, G. (1962).

The estimation of relative nutrient intake and nutritional status by biochemical methods : proteins.

Am. J. Clin. Nutr., 11, 447.

Arroyave, G. (1966).

Biochemical characteristics of malnourished infants and children. Prof. Western Hemisphere Nutrition Congress, Nov., 1965, A.M.A., 535 N. Dearbon St. Chicago, 60610, p.30.

Arroyave, G. and De Funes, C. (1967).

Razon nitrogeno ureico/creatinina comoindicador del nivel de ingesta proteica, II. Arch. Latinoam. Nutr., 17, 49.

Arroyave, G., Janson, A.A.J. and Torrico, M. (1966).

Razon nitrogeno ureico/creatinina comoindicador del nivel de ingesta proteica. Arch. Latinoam. Nutr., 16, 203.

Arroyave, G., Sandstead, H. and Schumacher, B. (1958).

Relation of urinary creatinine and vitamins to nutritional status in pre-school children. Fed. Proc., 17, 469.

Arroyave, G. and Wilson, D. (1961).

Urinary excretion of creatinine of children under different nutritional conditions. *Am. J. Clin. Nutr.*, 9, 170-175.

Arroyave, G., Wilson, D., Behar, M. and Scrimshaw, N.S. (1961).

Serum and urinary creatinine in children with severe protein malnutrition. *Am. J. Clin. Nutr.*, 9, 176-179.

Ashworth, A. (1969).

Metabolic rates during recovery from protein-calorie malnutrition-the need for a new concept of specific dynamic action. *Nature*, 223, 407-409.

Ashworth, A. and Harrower, A.D.B. (1967).

Protein requirements in tropical countries : Nitrogen losses in sweat and their relation to nitrogen balance. *Brit. J. Nutr.*, 21, 833.

Assessment of protein Nutritional status (1970).

A Committee Report. *Am. J. Clin. Nutr.*, 23, 807-819.

Atwater, W.C. and Bryant, A.P. (1899).

cf : Munro, H.N. (1964b).

Balfour, W.M., Habin, P.F., Bale, W.F., Pommerenke, W.T. and

Whipple, G.H. (1942). *J. Exptl. Med.*, 76, 15.

cf : Beaton, G.H. (1960).

Banerjee, S. (1962).

Studies on energy metabolism. *Ind. Counc. Med. Res.*, Spl. Rept. Ser. No. 43, ICMR, New Delhi.

Banerjee, S. (1972).

Some observations on energy intake and expenditure in different occupational groups. In 'Perspectives in Nutrition' (R.Rajalakshmi, Ed.), Department of Biochemistry, M.S. University, Baroda, 6-13.

Barnard, W.P. and Logan, R.W. (1971). @

Clin. Chim. Acta., 34, 377.

cf : Willocks, J. (1971).

Barrett, E. and Addis, T. (1947).

The serum creatinine concentration of normal individuals. J. Clin. Invest., 26, 875.

Bauer, J.D., Ackermann, G. and Toro, G. (1968).

Bray's Clinical Laboratory Methods. C.V. Mosby Company, p. 313.

Beach, C.F., Munks, B. and Robinson, A. (1943).

The amino acid composition of animal tissue protein. J. Biol. Chem., 148, 431-439.

Beard, H.H. (1943).

Creatine-creatinine Metabolism. Chemical Publ. Co., Brooklyn, New York.

Beaton, G.H. (1960).

Nutrition in maternal and infant feeding.

Proc. V Int. Cong. Nutr., Panel IV, Washington, D.C.

Beaton, G.H. (1961).

Nutritional and physiological adaptations in pregnancy. Fed. Proc., 20, 196-201.

Beaton, G.H., Arroyave, G. and Flores, M. (1964).

Alterations in serum proteins during pregnancy and
lactation in urban and rural populations in Guatemala.
Am. J. Clin. Nutr., 14, 269-279.

Benedict, F.G. (1907).

The influence of inanition on metabolism. Carnegie
Inst. Publ. No. 77.

cf : Keys et al (1950).

Benedict, P.G. (1915).

A study of prolonged fasting. Washington Carnegie
Inst. Publ. No. 203, p. 416.

Benedict, U.D., Kalinsky, H.J., Scarrone, L.A., Wertheim, A.R.
and Stetten, D. Jr. (1955).

The origin of urinary creatine in progressive muscular
dystrophy. J. Clin. Invest., 34, 141-145.

Bent Friis-Hansen (1965).

Hydrometry of growth and aging. In 'Human Body
Composition' (J. Brozek, Ed.). Pergamon Press, Oxford,
Vol. II, p. 197.

Bernhardt, W. (1974).

Die Wirkung Mehrtagiger Kreatinbelastung auf eine erhöhte
Kreatinausscheidung. J. Neurol., 207, 205.

cf : Talafant et al (1977).

Best, W.R. (1953).

Physiological factors in urinary creatinine excretion.
Army Med. Nutr. Lab. Report No. 118.

Best, W.R., Kuhl, W.J. Jr. and Consolazio, C.F. (1953).

Relation of creatinine coefficient to leanness-fatness
in man. J. Lab. Clin. Med., 42, 14, p. 184.

Best, C.H. and Taylor, N.B. (1967).

The Physiological Basis of Medical Practice. 8th edtn.
Scientific Book Agency, Calcutta.

Beydoun, S.N., Cuenca, V.G., Evans, L.P. and Aubry, R.H. (1972)

Maternal nutrition. The urinary urea-N/total-N ratio as
as index of protein nutrition. Am. J. Obst. Gynec.,
114, 198-203.

Bjornesjo, K.B., Belew, M. and Zaar, B. (1965).

Biochemical study of advanced protein malnutrition in
Ethiopia. Scand. J. Clin. Lab. Invest., 18, 591.

Blackburn, M.L. and Calloway, D.H. (1974).

Energy expenditure of pregnant adolescents.
J. Am. Diet. Assoc., 65, 24-30.

Bleiler, R.E. and Schedl, H.P. (1962).

Creatinine excretion : variability and relationships to
diet and body size. J. Lab. Clin. Med., 59, 945-955.

Bloch, A.S., Miles, H.M. and Shils, E.M. (1973).

Appendix. In 'Modern Nutrition in Health and Disease'.
(R.S. Goodhart and E.M. Shils, Eds.) Lea and Febiger,
Philadelphia, p. 1057.

Bloch, K. and Schoenheimer, R. (1939).

Studies in protein metabolism XI. The metabolic relation
to creatine and creatinine studied with isotopic nitrogen.
J. Biol. Chem., 131, 111-119.

Bohdal, M. and Simmons, W.K. (1969).

A comparison of the nutritional indices in healthy African, Asian and European children. Bull. W.H.O., 40, 166.

Bohdal, M., Gibbs, N.E. and Simmons, W.K. (1968).

Nutrition Survey and Campaign Against Malnutrition in Kenya, 1964-1968. Ministry of Health, Nairobi, Kenya.

Booher, L.E., Behan, I. and Mc Means, E. (1951).

Biologic utilization of unmodified food starch. J. Nutr., 45, 75-95.

Boothby, W.M. and Sandiford, R.B. (1921).

Nomogram for calculations of surface area from height and weight. Boston Med. Surg.J., 185, 337.

cf : West, et al, (1966).

Boothby, W.M., Berkson, J. and Dunn, H.L. (1936).

Am. J. Physiol., 116, 468.

cf : West et al (1966).

Borsook, H. and Dubnoff, J.W. (1947).

The hydroxylysis of phosphocreatine and origin of urinary creatinine. J. Biol. Chem., 168, 493-510.

Boyne, A.W., Chalmers, M.I. and Cuthbertson, D.P. (1953).

Z. Physiol. Chem., 295, 424.

cf : Blaxter, K.L. (1964).

Pregnancy and Lactation. In 'Mammalian Protein Metabolism' (H.N. Munro and J.B. Allison, Eds.), Academic Press, New York, Vol. II, Chapter, 15, p. 195.

- Bray, G., Schwartz, M., Rozin, R. and Lister, J. (1970).
Relationship between oxygen consumption and body
composition of obese patients. *Metabol. Clin. Exptl.*,
19, 418-429.
- Brewer, W., Porter, T., Ingalls, R. and Ohlson, M.A. (1946).
The urinary excretion of riboflavin by college women.
J. Nutr., 32, 583-596.
- Bricker, M.L. and Smith, J.M. (1951).
A study of endogenous nitrogen output of college women,
with particular reference to use of the creatinine output
in calculation of the biological values of the protein of
egg and sunflower seed flour. *J. Nutr.*, 44, 553-573.
- Brin, M., Dibble, M.V., Peel, A., McMullen, E., Bourquin, A.,
and Chen, N. (1965).
Some preliminary findings on the nutritional status of
aged in Onondaga country, New York. *Am. J. Clin. Nutr.*,
17, 240-258.
- Brody, S. (1945).
Bio-energetics and Growth, Reinhold, New York.
cf : Hegsted, D.M. (1964).
- Brooke, O.G. and Ashworth, A. (1972).
The influence of malnutrition on the postprandial
metabolic rate and respiratory quotient.
Brit. J. Nutr., 27, 407-415.
- Brown, R.A. and Sturtevant, M. (1949).
Vitamin requirements of the growing rats.
Vitamins and Hormones, 7, 171-199.

Burger, M. (1919).

Z. Ges. Exptl. Med., 9, 262.

cf : Waterlow, J.C. (1969).

Buskirk, E.R., Iampietro, P.F. and Welch, B.E. (1957).

Metabolism, 6, 144-

Cabacungan, N.B., Miles, C.W., Abernathy, R.P. and Ritchey, S.J.

(1973). Hydroxyproline excretion and nutritional status of children. Am. J. Clin. Nutr., 26, 173-176.

Calloway, D.H. (1975).

Nitrogen balance of men with marginal intakes of protein and energy. J. Nutr., 105, 914-923.

Calloway, D.H. and Margen, S. (1971).

Variation in endogenous nitrogen excretion and dietary nitrogen utilization as determinants of human protein requirements. J. Nutr., 101, 205-216.

Camara, A.A., Arn, K.D., Reimer, A. and Neuburgh, L.H. (1951).

The 24 hourly endogenous creatinine clearance as a measure of functional state of kidneys. J. Lab. Clin. Med., 37, 743.

Campbell, J.A. (1919).

Nitrogen partition in the urine of races in Singapore. Biochem. J., 13, 239-247.

Canham, J.E., Baker, E.M., Harding, R.S., Sauberlich, H.E. and Plough, I.C. (1969).

Dietary protein - its relationship to vitamin B₆ requirements and function. Ann. N.Y. Acad. Sci., 166, 16-29.

Cathcart, E.P. (1907).

Über die Zusammensetzung des Hungerharns.

Biochem. Z., 6, 109-148.

cf : Keys et al (1950).

Chambers, W.H. and Milhorat, A.T. (1928).

Muscular exercise and nitrogen metabolism of dogs.

J. Biol. Chem., 77, 603-618.

Chanutin, A. (1926).

The fate of creatine when administered to man.

J. Biol. Chem., 67, 29-41.

Chattaway, F.W., Hullin, R.P. and Odds, F.C. (1969).

The variability of creatinine excretion in normal subjects, mental patients and pregnant women.

Clin. Chim. Acta, 26, 567.

Chaudhuri, S.K. (1970).

Relationship of protein-calorie malnutrition with toxemia of pregnancy. Am. J. Obst. Gynec., 107, 33-37.

Cheek, D.B. (1968).

In 'Human Growth'. Lea and Febiger, Philadelphia.

Cheek, D.B., Hill, D.E., Cordano, A. and Graham, G.G. (1970).

Malnutrition in infancy : changes in muscle and adipose tissue before and after rehabilitation.

Pediat. Res., 4, 135.

Chetal, U., Mehra, U.R., Nath, K. and Ranjhan, S.K. (1975).

On the variation of urinary creatinine in buffalo calves and the effect of dietary protein intake on urinary

creatinine, creatine-nitrogen ratio and creatinine coefficient. J. Agri. Sci., 84, 1-5.
cf : Nutr. Abstr. Revs., 46, 139 (1976).

Chinn, K.S.K. (1966).

Potassium and creatinine as indexes of muscle and non-muscle protein in rats. J. Nutr., 90, 323-330.

Chinn, K.S.K. (1967).

Prediction of muscle and remaining tissue protein in man. J. Appl. Physiol., 23, 713-715.

Clark, H.E., Malzer, J.L., Onderka, H.M., Howe, J.M. and Moon, W.H. (1973).

Nitrogen balances of adult human subjects fed combinations of wheat, beans, corn, milk and rice. Am. J. Clin. Nutr., 26, 702-706.

Clark, H.E., Moon, W.H., Malzer, J.L., Birt, D.F. and Pang, R.L. (1974).

Nitrogen retention of adult human subjects fed varying quantities of tryptophan. J. Nutr., 104, 1121-1126.

Clark, H.E., Moon, W.H., Malzer, J.L. and Pang, R.L. (1974).

Nitrogen retention of young men who consumed between sixteen and eight grams of nitrogen from a combination of rice, wheat, chicken and milk. Am. J. Clin. Nutr., 27, 1059-1064.

Clark, L.C., Jr., Thompson, H., Beck, E.I. and Jacobson, W. (1951).

Excretion of creatine and creatinine by children. Am. J. Dis. Child., 81, 774-783.

Clarke, R.P., Cosgrove, L.D. and Morse, E.H. (1966).

Vitamin to creatinine ratios. Variability in separate voidings of urine of adolescents during a 24 hr period. Am. J. Clin. Nutr., 19, 335-341.

Cohenour, S.H. and Calloway, D.H. (1972).

Blood, urine and dietary pantothenic acid levels of pregnant teenagers. Am. J. Clin. Nutr., 25, 512-517.

Cohn, M., Simmonds, S., Chandler, J.P. and Du Vigneaud, V. (1946).

The effect of dietary level of methionine on the rate of transmethylation reactions in vivo. J. Biol. Chem., 162, 343-351.

Cole, S.W. (1933).

Practical physiological chemistry, Cambridge.

Concepcion, I. (1918).

Analysis of normal Filipino urine. Philip. J. Sci., 13, 347-359.

Consolazio, C.F., Johnson, H.L., Nelson, R.A., Dramise, J.G. and Skala, J.H. (1975).

Protein metabolism during intensive physical training in the young adult. Am. J. Clin. Nutr., 28, 29-35.

Consolazio, C.F., Nelson, R.A., Matoush, C.O., Harding, R.S. and Canham, J.E. (1963).

Nitrogen excretion of sweat and its relation to nitrogen balance requirements. J. Nutr., 79, 399-406.

Correa, O. (1944).

The creatinine standard in the prediction of basal metabolism. Rev. Chilena. Pediat. (Santiago, Chile), 15, 360-371. cf : Chem. Abst., 38, 6361-6362 (1944).

Corsa, L., Olney, J.M., Steenburg, R.W., Ball, M.R. and Moore, F.D. (1950).

The measurement of exchangeable potassium in man by isotope dilution. *J. Clin. Invest.*, 29, 1280-1295.

Cramer, K., Cramer, H. and Selander, S. (1967).

A comparative analysis between variation in 24 hr. urinary creatinine output and 24 hr. urinary volume. *Clin. Chim. Acta*, 15, 331-335.

Crim, M.C., Calloway, D.H. and Margen, S. (1972).

Effect of creatine feeding on creatine pool size in men. *Fed. Proc.*, 31, 727 (abstr.)

Crim, M.C., Calloway, D.H. and Margen, S. (1975).

Creatine metabolism in men. Urinary creatine and creatinine excretion with creatine feeding. *J. Nutr.*, 105, 428-438.

Crim, M.C., Calloway, D.H. and Margen, S. (1976).

Creatine metabolism in men : creatine pool size and turnover in relation to creatine intake. *J. Nutr.*, 106, 371-381.

Crowne, R.S., Wharton, B.A. and McCance, R.A. (1969).

Hydroxyproline indices and hydroxyproline/creatinine ratios in older children. *Lancet*, 1, 395-396.

Curtis, G. and Fogel, M. (1970).

Creatinine excretion : diurnal variation and variability of whole and part-day measures. *Physiosom. Med.*, 32, 337.

Dakshayani, R. and Ramanamurthy, P.S.V. (1964).

Basal metabolism and creatinine excretion in different physiological groups. Ind. J. Med. Res., 52, 1159-1164.

Daly, C. and Dill, D.B. (1937).

Salt economy in humid heat. Am. J. Physiol., 118, 285.

Darby, W.J., et al (1953).

The vanderbilt cooperative study of maternal and infant nutrition, IV. Dietary, laboratory and physical findings in 2, 129 delivered pregnancies. J. Nutr., 51, 565-597.

Das, T.K. and Waterlow, J.C. (1974).

The rate of adaptation of urea cycle enzymes, amino transferases and glutamic dehydrogenase to changes in dietary protein intake. Brit. J. Nutr., 32, 353-373.

Dave, J.R. (1976).

Studies on the composition of the bone in relation to age and nutritional status. Doctoral thesis, M.S.University, Baroda.

Davis, M.V., Oldham, H.G. and Roberts, L.J. (1946).

Riboflavin excretion of young women on diets containing varying levels of the B-vitamins. J. Nutr., 32, 143-161.

Dean, R.F.A. (1961).

Nitrogenous constituents of urine in kwashiorkor.

Fed. Proc., 20, 202-204.

Doolan, C.D.E., Alpen, E.L. and Theil, G.B. (1962).

A clinical appraisal of the plasma concentration and endogenous clearance of creatinine. Am. J. Med., 32, 65-79.

Donald, E.A., McBean, L.D., Simpson, M.H.W., Sun, M.F. and
Aly, H.E. (1971).

Vitamin B₆ requirement of young adult women.
Am. J. Clin. Nutr., 24, 1028-1041.

Drabikowski, W. (1957).

Modifikacja dwuacetylowej metody ozanaczania Kreatyny
i fosfokreatyny. Acta Bio-chim. Polon., 4, 41
cf : Talafant et al (1977).

Dugdale, A.E. and Edkins, E. (1964).

Urinary urea/creatinine ratio in healthy and
malnourished children. Lancet, 1, 1062-1064.

Du Plessis, J.P. (1967).

An evaluation of biochemical criteria for use in
Nutrition status surveys, National Nutrition Research
Institute, Council of Scientific and Industrial Research
Report No. 261, Pretoria, South Africa,

Plessis, J.P., De Lange, D.J. and Fellingham, S.A. (1966).

Biochemical investigations of the nutrition status of
urban school children aged 12-15 years : protein status.
S. Afr. Med. J., 27, 509.

Durnin, J.V.G.A. (1967).

Proc. VII Intern. Congr. Nutr., Hamburg, 1966, Pergamon
Press, Oxford, Vol. IV, p- 185.

Duque, E., Bolanos, O., Lotero, H. and Mayoral, L.G. (1975a).

Enteropathy in adult protein malnutrition light
microscopic findings. Am. J. Clin. Nutr., 28, 901-913.

Duque, E., Lotero, H., Bolanos, O. and Mayoral, L.G. (1975b).

Enteropathy in adult protein malnutrition : Ultra-
structural findings. Am. J. Clin. Nutr., 28, 914-924.

Edozien, J.C. and Phillips, E.J. (1961).

Partition of urine nitrogen in kwashiorkor.

Nature, 191, 47-49.

Edwards, O.M., Bayliss, R.I.S. and Miller, S. (1969).

Urinary creatinine excretion as an index of the
completeness of 24 hr. urine collection. Lancet, 2, 1165.

Edwards, K.D.G. and Whyte, H.M. (1959).

Creatinine excretion and body composition.

Clin. Sci., 18, 361-366.

Efremov, V.V. and Sakaeva, E.A. (1974).

Urinary excretion of creatine and creatinine by
experimental animals depending on physical work and
different amounts in the diet. Voprosy Pitaniya, 6, 19-22.
cf : Nutr. Abstr. Revs., 45, # 8813, (1975).

Ellestad-sayed, J., Hildes, J.A., Schaefer, O. and Lobban,
M.C. (1975).

Twenty four hour urinary excretion of vitamins, minerals
and nitrogen by Eskimos. Am. J. Clin. Nutr., 28, 1402-1407.

Elsom, K.O.S., Reinhold, J.G., Nicholson, J.T.L. and
Chornock, C. (1942).

Studies of the B vitamins in human subjects. V. The
normal requirement for thiamin some factors influencing
its utilization and excretion. Am. J. Med. Sci., 203,
569-577.

Everett, G.A. and Apgar, J. (1977).

Urinary taurine, sulfate and urea as possible indicators of zinc status in the rat. Nutr. Repts. International., 16, 391-396.

FAO (1965).

Protein requirements, FAO Nutr. Meet. Rept. Ser. No. 30, Food and Agriculture Organization of the United Nations, Rome.

FAO (1967).

Requirements of vitamin A, thiamine, riboflavine and niacin. Report of a joint FAO/WHO expert group, FAO Nutr. Meet. Repts. Ser. No. 41, Food and Agriculture Organization of the United Nations, Rome.

FAO (1973).

Energy and protein requirements. FAO Nutr. Rept. Ser. No.52 Food and Agriculture Organization of the United Nations, Rome.

Fenner, A., Lange, G-U., Monkemier, D. and Ohlenroth, G. (1974).

Creatinine levels in first urine of male preterm and term infants. J. Perinatol. Med., 2, 185.

Fisher, H. (1965).

Variations in the urinary creatinine excretion of rats fed diets with different protein and aminoacids content. J. Nutr., 85, 181-186.

Fitch, C.D., Lucy, D.D., Bornhofen, J.H. and Dalrymple, G.V. (1968).

Creatine metabolism in skeletal muscle. II. Creatine kinetics in man. Neurology, 18, 32-42.

Fitch, F.D., Oates, J.D. and Dinning, J.S. (1961).

The metabolism of creatine-1-C¹⁴ by mice with hereditary muscular dystrophy. J. Clin. Invest., 40, 850-856.

Fitch, C.D. and Sinton, D.W. (1964).

A study of creatine metabolism in diseases causing muscle wasting. J. Clin. Invest., 43, 444-452.

Folin, O. (1905).

Laws governing the chemical composition of urine.

Am. J. Physiol., 13, 66.

Frazier, E.I., Prather, M.E. and Hoene, E. (1955).

Nicotonic acid metabolism in humans. I. The urinary excretion of nicotonic acid and its metabolic derivatives on four levels of dietary intake. J. Nutr., 56, 501-516.

Friedemann, T.E., Kinney, V.M., Berryman, G.H., Henderson, C.R. and Youmans, J.B. (1948).

The effect of dietary restriction of B-complex vitamins and protein on the excretion of creatinine by human subjects. J. Nutr., 35, 117-136.

Fry, P.C., Fox, H.M. and Fry, E.I. (1968).

Effect of diet on skinfold measurements and creatinine and urea excretion of Hongkong chinese children.

Am. J. Clin. Nutr., 21, 1197-1201.

Fuwa, H. (1964).

J. Japan Soc. Food Nutr., 17, 240.

Galal, O.M. and Salem, S. (1977).

The role of nutrition on the process of virilization in boys during puberty. Nutr. Repts. International, 16, 103-113.

Galal, O.M., Salem, S., Mostafa, S., and Morcos, S.R. (1976).

The nutritional status and steroid excretion pattern during puberty. Nutr. Repts. International, 13, 413-424.

Garn, S.M. (1975).

Book Review of skeletal maturity of children 6-11 years, United States, by A.F. Rosche, J. Roberts and P.U.V. Hammill. From vital and health statistics, Series 11, no. 140, DHEW Publication No. 75-1622, p. 62, 1974. Am. J. Clin. Nutr., 28, p. 1337.

Garn, S.M., Smith, N.J. and Clark, D.C. (1975).

The magnitude and the implications of apparent race differences in hemoglobin values. Am. J. Clin. Nutr., 28, 563-565.

Garrow, J.S., Fletcher, K. and Halliday, D. (1965).

Body composition in severe infantile malnutrition. J. Clin. Invest., 44, 417-425.

Garrow, J.S. and Hawes, S.F. (1972).

The role of amino acid oxidation in causing 'Specific dynamic action' in man. Brit. J. Nutr., 27, 211-219.

Gault, M.H. and Cockcroft, D.W. (1975).

Creatine clearance and age. Lancet, 11, 612.

Ghadimi, H. (1975).

Protein economy in new born and infants. In 'Total prenatal nutrition' (H. Ghadimi, Ed.), A Wiley Bio-medical Health Publication, John Wiley and Sons, New York.

Glickman, N., Keeton, R.W., Cole, W.H., Calloway, N.O., Mitchell, H.H., Sapienza, A.R., Dyniewicz, J. and Howes, D. (1948).

Early ambulation in surgical convalescence.

Arch. Phys. Med., 29, 770-783.

Glickman, N., Mitchell, H.H., Lambert, E.H. and Keeton, R.W.

(1948). The total specific dynamic action of high protein and high-carbohydrate diets on human subjects.

J. Nutr., 36, 40-57.

Gokhale, S.K. (1963).

Nitrogen partition in urine. A study based on the examination of urine of 251 healthy Indian men.

Ind. J. Med. Res., 51, 91-102.

Goldsmith, G.A. (1950).

Relationships between nutrition and pregnancy as observed in recent surveys, in Newfoundland.

Am. J. Publ. Hlth., 40, 953-959.

Goldsmith, G.A. (1964).

The B vitamins : thiamine, riboflavin, e niacin. In 'Nutrition : A comprehensive treatise', (G.H.Beaton, and E.W. McHenry, Eds.) Academic Press, New York, Vol. II.

Goldsmith, G.A., Sarett, H.P., Register, U.D. and Gibbens, J.

(1952). Studies on niacin requirement in man. I. Experimental pellagra in subjects on corn diets low in niacin and tryptophan. J. Clin. Invest., 31, 533-542.

Goldsmith, G.A., Rosenthal, H.L., Gibbens, J. and Unglaug, W.G. (1955).

Studies on niacin requirement in man. II. Requirement on wheat and corn diets low in tryptophan. J. Nutr., 56, 371-386.

Gontzea, I., Sutzescu, P. and Dumitrache, S. (1974).

The influence of muscular activity on nitrogen balance and on the need of man for proteins. Nutr. Repts. International, 10, 35-43.

Gontzea, I., Sutzescu, R. and Dumitrache, S. (1975).

The influence of adaptation to physical effort on nitrogen balance in man. Nutr. Repts. International, 11, 231-236.

Gopalan, C. and Narasingrao, B.S. (1966).

Effect of protein depletion on urinary nitrogen excretion in undernourished subjects. J. Nutr., 90, 213-218.

Grande, F., Anderson, J.T. and Keys, A. (1955).

Creatinine excretion of men in negative and positive nitrogen balance. Fed. Proc., 14, ~~195~~ 195, p. 61.

Graystone, J.E. (1968).

In 'Human Growth' (D.B. Cheek, Ed.), Lea and Febiger, Philadelphia, p. 182.

Gutman, A.B. (1965).

Harvey Lectures Ser. 60, 35.

- Hahn, P.F., Carothers, E.L., Darby, W.J., Martin, M.,
Sheppard, C.W., Cannon, R.O., Beam, A.S., Densen, P.M.,
Peterson, J.C. and McClellan, G.S. (1951).
Iron metabolism in human pregnancy as studied with the
radioactive isotope Fe^{59} . Am. J. Obst. Gynec., 61, 477-486.
- Halliday, D. (1967).
Chemical composition of the whole body and individual
tissues of two Jamaican Children whose death resulted
from malnutrition. Clin. Sci., 33, 365-370.
- Hansen, J.M., Kampmann, J. and Laursen, H. (1970).
Renal excretion of drugs in the elderly.
Lancet, i, 1170.
- Hawk, P.B., Oser, B.L. and Summerson, W.H. (1954).
Practical physiological chemistry, Blachiston Company,
New York, 13th edn..
- Hawley, E.E., Murlin, J.R., Nasset, E.S. and Szymanski, T.A.
(1948). Biological values of six partially-purified
proteins. J. Nutr., 36, 153-169.
- Hegsted, D.M. (1964).
Protein requirements. In 'Mammalian Protein Metabolism'
(H.N. Munro and J.B. Allison, Eds.) Academic Press,
New York, Vol. II, Chapter 14.
- Hegsted, D.M., Gershoff, S.N., Trulson, M.F. and Jolly, D.H.
(1956). Variation in riboflavin excretion.
J. Nutr., 60, 581-597.

- Henshaw, J.L., Noakes, G., Morris, S.O., Bennion, M. and Gubler, C.J. (1970).
Method for evaluating thiamine adequacy in college women.
J. Am. Diet. Assoc., 57, 436.
- Hill, A.V. and Hill, A.M. (1913).
J. Physiol., 46, p. 95.
- Hoberman, H.D., Sims, E.A.H. and Engstrom, W.W. (1948).
The effect of methyl testosterone on the rate of synthesis of creatine. J. Biol. Chem., 173, 111-116.
- Hoberman, H.D., Sims, E.A.H. and Peters, J.H. (1948).
Creatine and creatinine metabolism in the normal male adult studied with the aid of isotopic nitrogen.
J. Biol. Chem., 172, 45-58.
- Horwitt, M.K. (1972).
Riboflavin. XII. Requirements and factors influencing them. In 'The Vitamins' (W.H. Sebrell and R.S. Harris, Eds.), Academic Press, New York, Vol. V, p. 88.
- Horwitt, M.K., Liebert, E., Kreisler, O. and Wittmann, P. (1948).
Investigation of human requirements for B-complex vitamins. National Research Council Bull. No. 116, National Research Council, Washington, D.C.
- Horwitt, M.K., Harvey, C.C., Hills, O.W. and Liebert, E. (1950)
Correlation of urinary excretion of riboflavin with dietary intake and symptoms of ariboflavinosis.
J. Nutr., 41, 247-264.

- Horwitt, M.K., Harvey, C.C., Rothwell, W.S., Cutler, J.L.
and Haffron, D. (1956).
Tryptophan-niacin relationships in man.
J. Nutr., 60, 1-43.
- Howat, P.M., Korslund, M.K., Abernathy, E.P. and Ritchey, S.J.
(1975). Sweat nitrogen losses and nitrogen balances of
preadolescent girls consuming three levels of dietary
protein. Am. J. Clin. Nutr., 28, 879-882.
- Howell, G.R., Wharton, B.A. and McCance, R.A. (1967).
Value of hydroxyproline indices in malnutrition.
Lancet, 1, 1082-1083.
- Hsu, J.M. and Antony, W.L. (1975).
Effect of zinc deficiency on urinary excretion of
nitrogen compounds and liver amino acid catabolizing
enzymes in rats. J. Nutr., 105, 26-31.
- Huang, P.C., Chong, H.E. and Rand, W.M. (1972).
Obligatory urinary and fecal nitrogen losses in young
chinese men. J. Nutr., 102, 1605-1614.
- Huang, P.C., Lo, C.C. and Ho, W.T. (1975).
Protein requirements of men in hot climate; decreased
urinary nitrogen losses concomitant with increased sweat
nitrogen losses during exposure to high environmental
temperature. Am. J. Clin. Nutr., 28, 494-501.
- Hughes, T.A., Malik, K.S. and Sahai, P.N. (1931).
A note on the output and distribution of urinary nitrogen
in the normal Punjabi. Ind. Med. Gaz, 66, 433-435.

Hyde, E. (1942).

Creatine feeding and creatine-creatinine excretion in males and females of different age groups.

J. Biol. Chem., 143, 301-310.

Inoue, G., Fujita, Y., Kishi, K., Yamamoto, S. and Niiyama, Y. (1974).

Nutritive values of egg protein and wheat gluten in young men. Nutr. Repts. International, 10, 201-207.

James, W.H. (1960).

Symposium on metabolic patterns in preadolescents children: Nitrogen balance. Fed. Proc., 19, 1009.

Jayalakshmi, V.T., Venkatachalam, P.S. and Gopalan, C. (1959).

Nitrogen balance studies in pregnant women in South India. Ind. J. Med. Res., 47, 86-92.

Jeffay, H. and Winzler, R.J. (1958).

The metabolism of serum proteins: The effect of dietary protein on the turnover of rat serum proteins.

J. Biol. Chem., 231, 111-116.

Jones, C.R., Bergman, M.W., Kittner, P.J. and Pigman, W.W. (1964).

Urinary hydroxyproline excretion in normal children and adolescents. Proc. Soc. Exp. Biol. Med., 115, 85-87.

Jourdan, M., Margen, S. and Bradfield, R.B. (1974).

Protein-sparing effect in obese women fed low calorie diets. Am. J. Clin. Nutr., 27, 3-12.

Junkersdorf, P. and Lisenfeld, F. (1926).

Stoffwechselversuche an Zwei 'Hunger Kunstlern' mit langdauernder Hungerzeit. Pflüger's Arch. Ges. Physiol., 214, 250-273. cf : Keys et al (1950).

Karambelkar, P.V., Patwardhan, V.N. and Srinivasan, A. (1952).

Studies in protein metabolism: Creatinine excretion in urine in relation to source of dietary protein.

Ind. J. Med. Res., 40, 89-93.

Katz, St. I. (1970).

The amino acid ratio and hydroxyproline creatinine index in marginal protein-calorie malnutrition.

Trop. Geogr. Med., 22, 389.

Kehar, N.D., Mukherjee, R., Murty, V.V.S. and Sen, K.C. (1943).

Studies on protein metabolism. II. The effect of muscular work on endogenous protein catabolism in cattle.

Ind. J. Vet. Sci., 13, 263-266.

Kennedy, W.P. (1961).

Creatine, creatinine, and total body potassium in relation to muscle mass in children. Arch. Dis. Child., 36, 325-327.

Kenney, R.A. (1954).

Renal clearance of creatinine and urea in the West Africans. Brit. Med. J., 1, 1130.

Kenny, F.M., Richards, C. and Taylor, F.H. (1970).

Reference standards for cortisol production and 17-hydroxy-corticosteroid excretion during growth, variation in the pattern of excretion of radio labeled cortisol metabolites. Metabol. Clin. Exptl., 19, 280-290.

Keys, A. (1955).

'Weight Control' (E.S. Eppright, P. Swanson, and G.A. Iverson, Eds.) Iowa State College, Press.

Keys, A., Brozek, J., Henschel, A., Michelsen, O. and Taylor, H.L. (1950).

The Biology of Human Starvation, Minneapolis, University of Minnesota Press.

Kiriyama, S. (1970).

Biological quality of dietary protein and urinary nitrogen metabolites. In 'Newer Methods of Nutritional Biochemistry' (A.A. Albanese, Ed.), Academic Press, New York, Vol. IV, Chapter 2.

Kiriyama, S. and Ashida, K. (1964).

Effect of quality of dietary protein on nitrogen compounds in the urine of rats. J. Nutr., 82, 127-134.

Kiriyama, S., Suzuki, T. and Iwao, H. (1963).

Ann. Rept. Natl. Inst. Nutr. Japan, p. 34(abstr)

cf : Kiriyama, S. (1970).

Kiriyama, S., Yagishita, Y., Suzuki, T. and Iwao, H. (1967).

Agr. Biol. Chem., 31, 743.

cf : Kiriyama, S. (1970).

Klein, L.R. and Yen, S.S.C. (1970).

Urinary peptide hydroxyproline before and during postpartum involution of human uterus. Metabol. Clin. Exptl., 19, 19-23.

Kocher, R.A. (1914).

Über die Grosse des Eiweisszerfalles und bei Fieber und bei Arbeitsleistung Untersuchungen mittele des Stickstoffminimums. Deut. Arch. Klin. Med., 115, 82-123.
cf : Mitchell, H.H. (1962).

Kopple, J.D. and Coburn, J.W. (1973).

Metabolic studies of low protein diets in ureamia.

1. Nitrogen and potassium. *Medicine*, 52, 583-595.

Krehl, W.A. and Hodges, R.E. (1965).

The interpretation of nutrition survey data.

Am. J. Clin. Nutr., 17, 191-199.

Kreisberg, R.A., Bowdoin, B. and Meador, C.K. (1970).

Measurement of muscle mass in humans by isotopic dilution of Creatine-C¹⁴. *J. Appl. Physiol.*, 28, 264-267.

Krzywicki, H.J., Consolazio, C.F., Matoush, L.O. and

Johnson, H.L. (1968).

Metabolic aspects of acute starvation, body composition changes. *Am. J. Clin. Nutr.*, 21, 87-97.

Kulkarni, M. and Kilgore, L. (1973).

Diurnal variations of hydroxyproline and creatinine excretion in children. *Am. J. Clin. Nutr.*, 26, 1069-1072.

Kumar, I., Land, D.G. and Boyne, A.W. (1959).

The determination of body composition of living animals.

The daily endogenous creatinine excretion as a measure of body composition in rats. *Brit. J. Nutr.*, 13, 320-329.

Lamden, M.P. (1972).

Thiamine, IX. Biochemical detection of deficiency.

In 'The Vitamins' (W.H. Sebrell and R.S. Barris, Eds.)

Academic press, New York, Vol. V, p. 134.

Lee, M. and Lucia, S.P. (1963).

Some relationships between caloric restriction and body weight in rat. IV. Fecal nitrogen, fat, energy and ash and urinary creatinine. J. Nutr., 81, 117-122.

LeRoy, E.C. (1967).

The technique and significance of hydroxyproline measurement in man. Adv. Clin. Chem., 10, 213-240, Academic Press, New York.

Lewis, J.S., Bunker, M.L., Getts, S.S. and Essien, R. (1975).

Variability of creatinine excretion of normal, phenylketonuric and galactosemic children and children treated with anticonvulsant drugs. Am. J. Clin. Nutr., 28, 310-315.

Linkswiler, H. (1967).

Biochemical and physiological changes in vitamin B₆ deficiency. Am. J. Clin. Nutr., 20, 547-557.

Lis, A.S., Mc Laughlin, D.I., Mc Laughlin, R.K. and

Hackbeil, K.F. (1972).

The function of creatinine: nocturnal-diurnal variation.

Physiol. Chem. Physics, 40, 70.

Lojkin, M.E., Wertz, A.W. and Dietz, C.G. (1952).

Metabolism of nicotinic acid in pregnancy.

J. Nutr., 46, 335-352.

Lowry, O.H. (1952).

Biochemical evidence of nutritional status.

Physiol. Rev., 32, 431-448.

Lusk, G. (1931).

The specific dynamic action. J. Nutr., 3, 519-530.

Luyken, R., Dubois, B., Leegwater, D.C., Pikkar, N.A. and
Van Staveren, W. (1970).

Correlation of protein intake and nitrogen and urea in
urine. Proc. Nutr. Soc., 29, 53A.

Macy, I.G. (1942a).

Chemical aspects of digestion and excretion. In 'Nutrition
and Chemical Growth in Childhood'. Charles C. Thomas,
Springfield, Illinois, Vol. I, Chapter VI. 113-131.

Macy, I.G. (1942b).

Energy metabolism in childhood. In 'Nutrition and Chemical
Growth in Childhood'. Charles C. Thomas, Springfield,
Illinois, Vol. I, Chapter VII, 132-150.

Magendie, F. (1783-1855).

cf : Munro, H.N. (1964b).

Malhotra, M.S., Chandra, U., Rai, R.M., Venkataswamy, Y., and
Sridharan, K. (1976).

Food intake and energy expenditure of Indian troops in
training. Brit. J. Nutr., 35, 229-244.

Manniello, J., Harmuth, E. and Fryer, J.H. (1960).

Body composition changes in men during weight loss on
a protein deficient diet. Fed. Proc., 19, 328.

Manual of Nutrition Surveys, (1963).

Interdepartmental Committee on Nutrition for National
Defence, Superintendent of Documents, U.S. Government
Printing Office, Washington, D.C., 2nd edn.

Margaria, R. and Foa, P. (1939).

Der Einfluss von Muskelarbeit auf den Stickstoffwechsel
die Kreatin-und Saureausscheidung. *Arbeits physiologie*, 10,
553-560. cf : Mitchell, H.H. (1962).

Mason, E.D. and Benedict, F.G. (1931).

Basal metabolism of South Indian women.

Ind. J. Med. Res., 19, 75-98.

Mason, H.L. and Williams, R.D. (1942).

The urinary excretion of thiamine as an index of the
nutrition level: Assessment of the value of test dose.

J. Clin. Invest., 21, 247.

Mayoral, L.G., Bolanos, O., Lotero, H. and Duque, E. (1975).

Enteropathy in adult protein malnutrition: A review of
Cali experience. *Am. J. Clin. Nutr.*, 28, 894-900.

Mc Cay, D. (1908).

Scientific memoirs of the Government of India, no. 34.

Mc Donald, J.T. and Margen, S. (1976).

Wine versus ethanol in human nutrition. 1. Nitrogen and
calorie balance. *Am. J. Clin. Nutr.*, 29, 1093-1103.

McLaren, D.S., Loshkajian, H. and Kanawati, A.A. (1970).

Urinary creatinine and hydroxoproline in relation to
childhood malnutrition. *Brit. J. Nutr.*, 24, 641-651.

Mc Millan, M. and Reid, C. (1965).

Body composition, resting oxygen composition and urinary
creatinine in Edinburg students. *Lancet*, 1, 728.

Meador, C.K., Kreisberg, R.A., Friday, J.P., Bowdoin, B.,

Coan, P., Armstrong, J., and Hazelrig, J.B. (1968).

Muscle mass determination by isotopic dilution of creatine-C¹⁴. *Metabol, Clin. Exptl.*, 17, 1104-1108.

Melnick, D. (1942).

Vitamin B₁ (thiamine) requirement of man.

J. Nutr., 24, 139-151.

Mendez, J. and Buskirk, E.R. (1971).

Creatinine height index. *Am. J. Clin. Nutr.*, 24, 385-386.

Mickelsen, O. and Anderson, A.A. (1959).

A method for preparing intact animals for carcass analysis. *J. Lab. Clin. Med.*, 53, 282-290.

Mickelsen, O., Caster, W.O. and Keys, A. (1947).

A statistical evaluation of the thiamine and pyramin excretions of normal young men on controlled intakes of thiamine. *J. Biol. Chem.*, 168, 415-431.

Milenkovic, M., Kucukalic, E. and Muftic, A. (1974).

Changes in content and ratios of some nitrogen compounds in the urine of rats fed with different quantities of the same protein. *Veterinaria, Yugoslavia*, 23, 17-25.
cf : *Nutr. Abstr. Revs.*, 45, ~~4~~ 4208 (1975) p.518.

Miller, A.T. Jr. and Blyth, C.S. (1952).

Estimation of leanbodymass and body fat from basal oxygen consumption and creatinine excretion. *J. Appl. Physiol.*, 5, 73-78.

Miller, L.T. and Linkswiler, H. (1967).

Effects of protein intake on the development of abnormal tryptophan metabolism by men during vitamin B₆ depletion. J. Nutr., 93, 53-59.

Miller, D.S., Mumford, P. and Stock, M.J. (1967).

Gluttony 2. Thermogenesis in over-eating men. Am. J. Clin. Nutr., 20, 1223-1229.

Mills, J.N. (1964).

Circadian rhythms during and after three months in solitude underground. J. Physiol., 174, 217-231.

Milner, J.A., Wakeling, A.E. and Visek, W.J. (1974).

Effect of arginine deficiency on growth and intermediary metabolism in rats. J. Nutr., 104, 1681-1689.

Mitchell, H.H. (1962).

Comparative Nutrition of man and Domestic Animals. Academic Press, New York, Vol. I.

Mitchell, H.H. (1964).

Comparative Nutrition of Man and Domestic Animals. Academic Press, New York, Vol. II.

Mitchell, H.H. and Kruger, J.H. (1928).

The effect of muscular work upon the endogenous catabolism of the tissues. J. Biol. Chem., 76, 55-74.

Mitchell, H.H. and Hamilton, T.S. (1949).

The dermal excretion under controlled environment conditions of nitrogen and minerals in human subjects with particular reference to calcium and iron. J. Biol. Chem., 178, 345-361.

Mitchell, H.H. and Edman, M. (1962).

Nutritional significance of dermal losses of nutrients in man, particularly of nitrogen and mineral.

Am. J. Clin. Nutr., 10, 163-172.

Mueller, A.J. and Cox, W.M. (1947).

Comparative nutritive value of casein and lactalbumin for man. J. Nutr., 34, 285-294.

Muldowney, F.P., Crooks, J. and Bluhm, M.M. (1957).

The relationship of total exchangeable potassium and chloride to leanbody mass, redcell mass and creatinine excretion in man. J. Clin. Invest., 36, 1375-1381.

Munro, H.N. (1969).

Evolution of protein metabolism in mammals. In 'Mammalian Protein Metabolism'. (H.N. Munro, Ed.), Academic Press, New York, Vol. III, Chapter 25.

Munro, H.N. (1964a).

General aspects of the regulation of protein metabolism by diet and by hormones. In 'Mammalian Protein Metabolism'. (H.N. Munro, and J.B. Allison, Eds.), Academic Press, New York, Vol. I, Chapter 10, p. 406.

Munro, H.N. (1964b).

Historical Introduction: The origin and growth of our present concepts of protein metabolism. In 'Mammalian Protein Metabolism'. (H.N. Munro and J.B. Allison, Eds.). Academic Press, New York, Vol. I, Chapter 1.

Muramatsu, K. and Ashida, K. (1955).

J. Agr. Chem. Soc., Japan, 29, 725.

cf : Kiriyama, S. (1970).

Murawski, B.J., Jones, K.J. and Reddy, W.J. (1970).

Steroid output in healthy young men of college age.

Metabol. Clin. Exptl., 19, 341-348.

Murlin, J.R., Edwards, L.E., Hawley, E.E. and Clark, L.C. (1946).

Biological value of proteins in relation to the essential aminoacids which they contain. 1. The endogenous nitrogen of man. J. Nutr., 31, 533-553.

Myers, V.C. and Fine, M.S. (1913).

The creatine content of muscle under normal conditions.

Its relation to urinary creatinine. J. Biol. Chem., 14, 9-26.

Nagamine, S., Yamakawa, K., Isobe, S., Chinose, Y. and

Nakagami, T. (1973).

Relationship of urinary hydroxyproline excretion to growth rate and nutritional status in Japanese school children.

Japanese J. Nutr., 31, 106-112.

cf : Nutr. Abstr. Revs. 45, ~~1516~~ (1975) p.221.

Nagase, S. (1976).

Hydroxyproline and creatinine in the human amniotic fluid: an indicator of fetal growth. Osaka. Shiritsu.

Daigaku Igaku. Zasshi, 25, 117-125.

cf : Chem. Abstr., 86, ~~186732~~ (1977), p. 350.

Nageswara Rao, C., Nadamuni Naidu, A. and Narasingarao, B.S.

(1975). Influence of varying energy intake on nitrogen balance in men on two levels of protein intake.

Am. J. Clin. Nutr., 28, 1116-1121.

Nakagawa, I., Takahashi, T. and Suzuki, T. (1961).

Amino acid requirements of children; Minimal needs of lysine and methionine based on nitrogen balance method.

J. Nutr., 74, 401-407.

Nakagawa, I., Takahashi, T., Suzuki, T. and Kobayashi, K.

(1962). Amino acid requirements of children: Minimal needs of threonine, valine and phenylalanine based on nitrogen balance method. J. Nutr., 77, 61-68.

Nakagawa, I., Takahashi, T., Suzuki, T. and Kobayashi, K.

(1963). Amino acid requirements of children: Minimal needs of tryptophan, arginine, and histidine based on the nitrogen balance method. J. Nutr., 80, 305-310.

Nakagawa, I., Takahashi, T., Suzuki, T. and Kobayashi, K.

(1964). Amino acid requirements of children: Nitrogen balance at the minimal level of essential amino acids.

J. Nutr., 83, 115-118.

Nakagawa, I., Sasaki, A., Kajimoto, M., Fukuyama, T.,

Suzuki, T. and Yameda, E. (1974).

Effect of protein nutrition on growth, longevity and incidence of lesions in the rat. J. Nutr., 104, 1576-1583.

Nakagawa, I., Takahashi, T., Sasaki, A., Kajimoto, M. and

Suzuki, T. (1973). Efficiency of conversion of tryptophan to niacin in humans. J. Nutr., 103, 1195-1199.

Narayan, K.G.A. (1935).

Nitrogen partition in the urine of South Indians.

Curr. Sci., 45, 96.

NAS-~~ERC~~ (1962).

Nutrient requirement of laboratory animals No. 10.

Publ. 990., National Academy of Sciences, National Research Council, Washington, D.C.

Nichols, J., Miller, A.T. and Hiatt, E.P. (1951).

Influence of muscular exercise on uric acid excretion in man. J. Appl. Physiol., 3, 501-507.

Nicol, B.M. and the Late Phillips, P.G. (1976).

Endogenous nitrogen excretion and utilization of dietary protein. Brit. J. Nutr., 35, 181-193.

Nishizawa, N., Shimbo, M., Hareyama, S. and Funabiki, R. (1977).

Fractional catabolic rates of myosin and actin estimated by urinary excretion of N'-methyl histidine; the effect of dietary protein level on catabolic rates under conditions of restricted food intake. Brit. J. Nutr., 37, 345-353.

Niyogi, S.P., Patwardhan, V.N. and Sirasat, M.V. (1941).

Studies on basal metabolism in Bombay Part III. An examination of the factors influencing the basal metabolism. Ind. J. Med. Res., 29, 289-302.

Norris, A.H., Lundy, T. and Shock, N.W. (1963).

Trends in selected indices of body composition in men between the ages 30 and 80 years. Am. N.Y. Acad. Sci., 110, 623-639.

Novak, L.P. (1963).

Age and sex differences in body density and creatinine excretion of high school children. Ann. N.Y. Acad. Sci., 110, 543.

NRC (1968).

Recommended dietary allowances, A report of the Food and Nutrition Board, National Research Council, Publ. No.1694; National Academy of Sciences, Washington, D.C.

NRL (1968).

Annual Report of Nutrition Research Laboratory, Hyderabad, India.

Nutritional Survey Reports (1957-1972).

Interdepartmental Committee on Nutrition for National Defence, Washington, D.C.

Ohnaka, M. (1976).

Changes in energy expenditure of human subjects during starvation in relation to endocrine function.

Nutr. Repts. International, 14, 681-690.

Oldham, H.C., Davis, M.V. and Roberts, L.J. (1946).

Thiamine excretions and blood levels of young women on diets containing varying levels of B vitamins, with some observations on niacin and pantothenic acid.

J. Nutr., 32, 163-180.

O'Neal, R.M., Johnson, O.C. and Schaefer, A.E. (1970).

Guidelines for classification and interpretation of group blood and urine data collected as part of the National Nutrition Survey. Pediatr.Res., 4, 103.

Oomen, H.A.P.C. (1967).

Nitrogen compounds and electrolytes in the urine of New Guinean sweet potato eaters-a study of normal values.

Trop. Geogr. Med., 19, 31-47.

Oomen, H.A.P.C. and Malcolm, S. (1958).

Nutrition of papuan child, Nourmes, New Caledonia, South Pacific Commission Technical Paper No. 118.

Owen, O.E., Felig, P., Morgan, A.P., Warhren, J. and Cahill, G.P. (1969). Liver and kidney metabolism during prolonged starvation. J. Clin. Invest., 48, 574-583.

Pace, N., Grunbaum, B.W., Kodama, A.M., Rahlman, D.F. and Newsom, B.D. (1975).

Effect of 14 days of bed rest on urine metabolite excretion and plasma enzyme levels. Life Sci. Space Res., 13, 41-47.

cf : Chem. Abstr. 86, ~~77~~ 28123 S, (1977) p. 227

Pain, S.K. and Banerjee, S. (1957).

Studies on nitrogenous constituents of urine in normal subjects and in patients suffering from cirrhosis of liver, subacute nephritis and hypertension. Ind. J. Med. Res., 45, 35-39.

Palmer, W.W., Means, J.H. and Gamble, J.L. (1914).

Basal metabolism and creatinine elimination.

J. Biol. Chem., 19, 239-244.

Park, S.Y. (1973).

The effect on the protein metabolism in albino rats by feeding on the rice mixed with wheat or barley diet.

Korean J. Nutr., 6, 179-185.

cf : Nutr. Abstr. Revs., 45, ~~4~~ 4205 (1975) p. 517.

Pasternack, A. and Kuhlback, B. (1971).

Diurnal variation of serum and urine creatine and creatinine. Scand. J. Clin. Lab. Invest., 27, 1-7.

cf : Biol. Abstr., 52, ~~7~~ 70393 (1971).

Paterson, N. (1967).

Relative constancy of 24 hr. urine volume and 24 hr. creatinine output. Clin. Chim. Acta, 18, 57.

Patwardhan, V.N. (1961).

Biochemistry of human protein metabolism.

Fed. Proc., 20, 73-79 (suppl. 7).

Peters, W.H., Grosser, V. and Knapp, A. (1972).

Creatinine excretion of women during therapeutic starvation. Acta. Medica. Academiae Scientiarum Hungaricae, 29, 357-363.

cf : Nutr. Abstr. Revs., 45, ~~4~~ 438 (1975) p. 61.

Pearson, W.N. (1962a).

Biochemical appraisal of nutritional status in man.

Am. J. Clin. Nutr., 11, 462.

Pearson, W.N. (1962b).

Biochemical appraisal of the vitamin nutritional status in man. J.A.M.A., 180 49-55.

Pearson, W.N. (1966).

Assessment of Nutritional Status: Biochemical Methods.

In 'Nutrition'. A comprehensive Treatise' (G.H.Beaton and E.W. McHenry, Eds.), Academic Press, New York, Vol.III, Chapter 7.

Pearson, W.N. (1967).

Blood and urinary vitamin levels as potential indices of body stores. Am. J. Clin. Nutr., 20, 514-525.

Pearson, W.N. (1967).

Thiamine, In 'The Vitamins', (P. Gyorgy and W.N.Pearson, Eds.), Academic Press, New York, Vol. III, p. 53.

Phansalkar, S.V. and Patwardhan, V.N. (1954).

A Partition of urinary nitrogen in Indian adults, relation between Urea-N and Total-N. Ind. J. Med. Res., 42, 363-371.

Picou-Reategu, E. (1962).

Creatinine excretion and body composition.

Am. J. Clin. Nutr., 10, 128-133.

Picou, D., Alleyne, G.A.O. and Seakins, A. (1965).

Hydroxyproline and creatinine excretion in infantile protein malnutrition. Clin. Sci., 29, 517-523.

Picou, D., Halliday, D. and Garrow, J.S. (1966).

Total body protein, collagen and non-collagen protein in infantile protein malnutrition. Clin. Sci., 30, 345-351.

Picou, D., Alleyne, G.A.O., Waterlow, J.C. and Seakins, A. (1965).

Hydroxyproline and creatinine excretion in protein-depleted infants. Biochem. J., 95, 18p.

Picou, D., Reeds, P.J., Jackson, A. and Poulter, N. (1976).

The measurement of muscle κ mass in children using N^{15} creatine. *Pediatr. Res.*, 10, 184-188.

Pierro, A. and Johnson, R.E. (1970).

Creatinine excretion. *Lancet*, 1, 784.

Pike, R.L., Suder, H.B. and Rose, M.L. (1954).

The influence of protein and energy intakes upon nitrogen retention in the pregnant rat. *J. Nutr.*, 52, 297-309.

Pittet, P.H., Gyax, P.H. and Jequier, E. (1974).

Thermic effect of glucose and amino acids in man studied by direct and indirect calorimetry. *Brit. J. Nutr.*, 31, 343-349.

Platt, B.S. (1954).

In 'Malnutrition in African mother, Infants, and Young children'. Rept. Second Inter-African Conf. Nutr. (Gambia) 1952, H.M. Stationery Office, London.

Platt, B.S. (1958).

Malnutrition and the pathogenesis of disease.

Trans. Roy. Soc. Trop. Med. Hyg., 52, 189.

Platt, R., Stewart, A.E. and Energy, E.W. (1958).

The aetiology, incidence and heredity of pre-eclamptic toxemia of pregnancy. *Lancet*, 1, 552-556.

Platt, B.S., Heard, C.R.C. and Stewart, R.J.C. (1964).

Experimental protein calorie deficiency. In 'Mammalian Protein Metabolism' (H.N. Munro and J.B. Allison, Eds.), Academic Press, New York, Vol. II, Chapter 21.

Plough, I.C. and Consolazio, C.F. (1959).

The use of casual urine specimens in the evaluation of the excretion rates of thiamine, riboflavin and N'-methyl-nicotinamide. J. Nutr., 69, 365-370.

Pollack, H. (1970).

Creatinine excretion as an index for estimating urinary excretion of micronutrients or their metabolic end products. Am. J. Clin. Nutr., 23, 865-867.

Pollack, H., Ellenberg, M. and Dolger, H. (1941).

Study of the excretion of thiamin and its degradation products in humans. J. Nutr., 21, 10-15.

Powell, R.C., Plough, I.C. and Baker, E.M. (1961).

The use of nitrogen to creatinine ratios in random urine specimens to estimate dietary protein. J. Nutr., 73, 47-52.

Prasad, L.S. and Rahman, A. (1970).

Urinary excretion of hydroxyproline and creatinine in malnutrition. Ind. Pediatr., 7, 54-56.

Prinsloo, J.G., Du Plessis, J.P., Kruger, H., de Lange, D.J. and de Villiers, L.S. (1968).

Protein nutrition status in childhood pellagra. Evaluation of nicotinic acid status and creatinine excretion. Am. J. Clin. Nutr., 21, 98-106.

Rajalakshmi, R. (1971).

Reproductive performance of poor Indian women on a low plane of nutrition. Trop. Geogr. Med., 23, 117-125.

Rajalakshmi, R. and Ramakrishnan, C.V. (1977).

Formulation and evaluation of meals based on locally available foods for young children. *Wld. Rev. Nutr. Diet.*, 27, 34-104.

Ram, M.M. and Reddy, V. (1970).

Variability in urinary creatinine. *Lancet*, 2, 674.

Ramachandran, K. (1968).

Formulation and evaluation of low cost balanced meals based on locally available foods for children in the postweaning period. Doctoral Thesis, M.S. University, Baroda.

Ramakrishnarao, P., Puri, R.K. and Balakrishnan, S. (1973).

Plasma aminoacids and urinary nitrogen partition in kwashiorkor. *Ind. Pediatr.*, 10, 173-176.

Ramiah, P.V., Sundaram, M. and Narayanayya, Y.V. (1939).

The effect of muscular work on protein metabolism in the ruminant. *Currt. Sci.*, 8, 161-162.

Rammurthi, K. (1955).

Studies on the nitrogen partition in the urine of Rajasthanis. *Ind. J. Med. Res.*, 43, 61-69.

Rao, C.N. and Rao, B.S.N. (1974).

Nitrogen balance in pregnancy and lactation in women whose protein intake is marginal. *Ind. J. Med. Res.*, 62, 1619-1626.

Ray S.N. and Ganguly, R. (1938).

Urinary composition of normal Bengali subjects. *Ind. J. Med. Res.*, 26, 459-463.

Reddy, V. (1964).

Partition of urinary nitrogen in pregnancy and lactation.

Ind. J. Med. Res., 52, 1192-1197.

Reddy, V., Belavady, B. and Srikantia, S.G. (1963).

Partition of urinary nitrogen in marasmus.

Ind. J. Med. Res., 51, 952-957.

Refsum, H.E. and Stromme, S.B. (1974).

Urea and creatinine production and excretion in urine during and after prolonged heavy exercise.

Scand. J. Clin. Lab. Invest., 33, 247-254.

cf : Nutr. Abstr. Revs., 45, ~~7~~ 7252 (1975) p.836.

Reindrop, S. and Whitehead, R.G. (1971).

Change in serum creatine kinase and other biological measurements associated with musculature in children recovering from kwashiorkor. Brit. J. Nutr., 25, 273-283.

Ritchey, S.J., Derise, N.L., Abernathy, R.P. and Korslund, M.K.

(1973). Variability of creatinine excretion in pre-adolescent girls consuming a wide range of dietary nitrogen.

Am. J. Clin. Nutr., 26, 690-695.

Rose, W.C., Ellis, R.H. and Helming, O.C. (1928).

The transformation of creatine into creatinine by the male and female organism. J. Biol. Chem., 77, 171-184.

Rose, W.C., Johnson, J.E. and Haines, W.J. (1960).

The aminoacid requirements of man: The role ^{of} valine and methionine. J. Biol. Chem., 182, 541-556.

Rose, W.C., Haines, W.J., Warner, D.T. and Johnson, J.E.(1951).

The aminoacid requirements of man: The role of threonine and histidine. J. Biol. Chem., 188, 49-58.

Rubin, R.T. (1971).

Urine creatinine excretion: variability and volume dependency during sleep deprivation. Psychosom. Med., 33, 539.

Rubner, M. (1902).

Die Gesetze des Energieverbrauchs bei der Ernährung, Leipzig and Vienna, Deuticke,
cf : Mitchell, H.H. (1964).

Runcie, J. and Hilditch, T.E.(1974).

Energy provision, tissue utilization and weight loss in prolonged starvation. Brit. Med. J., 2, 352-356.

Rutishauser, I.H.E. and Whitehead, R.G. (1969);

Field evaluation of two biochemical tests which may reflect nutritional status in three areas in Uganda. Brit. J. Nutr., 23, 1-13.

Ryan, R.J., Williams, J.D., Ansell, B.M. and Bernstein, L.M. (1957). The relationship of body composition to oxygen consumption and creatinine excretion in healthy and wasted men. Metabolism, 6, 365-377.

Sail, S.S. (1970).

Nutritional studies on school children.

Doctoral Thesis, M.S.University, Baroda.

Sallan, V. and Puri, B. (1962).

The assessment of nutritional status of lactating women and chemical composition of human colostrum. Studies in Nutrition, Lady Irwin College, New Delhi, 1967.

Sauberlich, H.E. (1967).

Biochemical alterations in thiamine deficiency—Their interpretation. Am. J. Clin. Nutr., 20, 528-542.

Sauberlich, H.E. and Canham, J.E. (1973).

Vitamin B₆. In 'Modern Nutrition in Health and Disease' (R.S. Goodhart and E.M. Shils, Eds.), Lea and Febiger, Philadelphia.

Sauberlich, H.E., Herman, Y.F. and Stevens, C.O. (1970).

Thiamin requirement of the adult human. Am. J. Clin. Nutr., 23, 671-672.

Sauberlich, H.E., Canham, J.E., Baker, E.M., Raica, N. Jr., and Herman, Y.F. (1970).

Human B₆ nutriture. J. Sci. Ind. Res., 29, 528.

Schedl, H.P. (1974).

The concept of pool expansion in creatinine excretion. Am. J. Clin. Nutr., 27, 769.

Schendel, H.E. and Hansen, J.D.L. (1965).

Daily urinary nitrogen partition and balance in infants with kwashiorkor. Am. J. Clin. Nutr., 17, 36-50.

Scott, P.J. and Hurley, P.J. (1968).

Demonstration of individual variation in constancy of 24 hr. urinary creatinine excretion. Clin. Chim. Acta, 21, 411.

Scrimshaw, N.S., Taylor, Y. and Young, V.R. (1973).

Lysine supplementation of wheat gluten at adequate and restricted energy intakes in young men. Am. J. Clin. Nutr., 26, 965-972.

Scrimshaw, N.S., Petera, W.D.A. and Young, V.R. (1976).

Protein requirements of man: Obligatory urinary and fecal nitrogen losses in elderly women. J. Nutr., 106, 665-670.

Scrimshaw, N.S., Hussien, M.A., Murray, E., Rand, W.M. and Young, V.R. (1972).

Protein requirements of man: Variations in obligatory urinary and fecal nitrogen losses in young men. J. Nutr., 102, 1595-1604.

Seguin, A. and Lavoisier, A.L. (1789).

"Histoire de L' Academie des Sceinces, Annee 1789"
(published, 1793) p. 566.

cf : Krebs, H.A. (1964).

In 'Mammalian Protein Metabolism' (H.N. Munro and J.B. Allison, Eds.), Academic Press, New York, Vol. I, Chapter 5, p. 165.

Sengupta, N. and Bagchi, K. (1961).

Dietary habits of Bengali Pregnant and Lactating women of low socio-economic groups in Calcutta, Alumni Bull. September, 1-6.

Sertel, H. and Scopes, J. (1973).

Rates of creatinine clearance in babies less than one week of age. Arch. Dis. Child, 48, 717-720.

Shaffer, P. (1908).

The excretion of Kreatinin and Kreatin in health and disease. Am. J. Physiol., 23, 1-22.

Shankar, K. (1962).

The dietary intake and nutritional status of pregnant and nursing women in Hyderabad. Ind. J. Med. Res., 50, 113-119.

Shimazono, N. and Katasura, E. (1965).

Eds. of Review of Japanese Literature on Beriberi and Thiamine, Vitamin B Research Committee of Japan, Clinical Nutrition, Faculty of Medicine, Kyoto University, Kyoto, Japan.

Siersbaek-Nielsen, K., Hansen, J.M., Kampmann, J. and

Kristensen, M. (1971).

Rapid evaluation of creatinine clearance. Lancet 1, 1133.

Simko, V., Merrifield, H.H. and Stouffer, T.R. (1973).

Effect of exercise and body composition on metabolism in young women. Am. J. Clin. Nutr., 26, XXIX

Simmons, W.K. (1972).

Urinary urea nitrogen/creatinine ratio as indicator of recent protein intake in field studies. Am. J. Clin. Nutr., 25, 539-542.

Simmons, W.K. (1973).

Use of inorganic sulfate sulfur/creatinine ratio in field studies. Am. J. Clin. Nutr., 26, 62-76.

Simmons, W.K. and Bohdal, M. (1970).

Assessment of some biochemical parameters related to protein-calorie nutrition in children. Bull. W.H.O., 42, 897-906.

Sitren, H.S. and Fisher, H. (1977).

Nitrogen retention in rats fed on diets enriched with arginine and glycine. Brit. J. Nutr., 37, 195-208.

Slonim, A.R. (1961).

Effect of high accelerations on some biological systems.

Ind. J. Med. Res., 16, 221.

Smiley, J.D. and Ziff, M. (1964).

Urinary hydroxyproline excretion and growth.

Physiol. Rev., 44, 30-44.

Smith, M. (1926).

The minimum endogenous nitrogen metabolism.

J. Biol. Chem., 68, 15-31.

Smuts, D.B. (1935).

The relation between the basal metabolism and the endogenous nitrogen metabolism, with particular reference to the estimation of the maintenance requirement of protein.

J. Nutr., 9, 403-433.

Sorg-Matter, H. (1928).

Arch. Intern. Physiol., 20, 126.

cf : Hegsted, D.M. (1964).

Southgate, D.A.T. and Durmin, J.V.G.A. (1970).

Caloric conversion factors. An experimental reassessment of the factors used in the calculation of the energy value of human diets. Brit. J. Nutr., 24, 517-535.

Spector, H., Peterson, M.S. and Friedmann, T.E. (1954).

Eds. ~~of~~ 'Methods for evaluation of nutritional adequacy and status'. National Academy of Science, National Research, Council Publication, Washington, D.C.

Spencer, H., Lewin, I., Samachson, J. and Laszlo, J. (1966),

Changes in metabolism in obese persons during starvation.
Am. J. Med., 40, 27-37.

Spray, C.M. (1950).

A study of some aspects of reproduction by means of chemical analysis. Brit. J. Nutr., 4, 354-360.

Srikantia, S.G., Paragaonkar, V.U. and Reddy, V. (1965).

Studies on creatinine metabolism in kwashiorkor and marasmus. Am. J. Clin. Nutr., 16, 436-439.

Srivastava, S.S., Mani, K.V., Soni, C.M. and Bhati, J. (1967).

Effect of muscular exercise on urinary excretion of creatine and creatinine. Ind. J. Med. Res., 55, 953-960.

Standard, K.L., Wills, V.G. and Waterlow, J.C. (1959).

Indirect indicators of muscle mass in malnourished infants.
Am. J. Clin. Nutr., 7, 271-279.

Starling, E.K. and Evans, L.C. (1956).

Principles of Human Physiology. J.A. Churchill Ltd., London, p. 959.

Stearns, G., Adamson, L., McKiney, J.B., Linner, T. and

Jeans, P.C. (1958).

Excretion of thiamine and riboflavine by children.
Am. J. Dis. Child, 95, 185-201.

Stearns, G., Newman, K.J., McKinley, J.G. and Jean, P.C. (1958)

The protein requirements of children from one to ten years of age. Ann. N.Y. Acad. Sci., 69, 857-868.

Steinbock, H.L. and Tarver, H. (1954).

Plasma proteins. V. The effect of protein content of the diet on turnover. J. Biol. Chem., 209, 128-132.

Stirling, J.L. and Stock, M.J. (1968).

Metabolic ~~org~~gins of thermogenesis induced by diet.

Nature, 220, 801.

Stuart, H.C. and Stevenson, S.S. (1954).

Physical growth and development. In 'Text Book of Pediatrics' (W.E. Nelson, Ed.), Saunders, Philadelphia, p.10-66.

Stuber, A. and Paksi, A. (1976).

The uric acid creatinine index ^fin normal and mentally retarded population. Orv. Hetil., 117, 2167-2169.

cf : Chem. Abstr., 86, ~~≠~~ 53600 C, p. 289 (1977).

Subbulakshmi, G. (1970).

Studies on pregnancy and lactation.

Doctoral thesis, M.S. University, of Baroda, Baroda.

Sulovic, V. and Poljakovic, L. (1976).

Estriol-creatinine ratio in determining the condition of the fetal-placental system. Jugost. Ginek. Opstel., 16, 25-32. cf : Chem. Abstr., 86, ~~≠~~ 69669 t, (1977) p. 386

Swift, R.W. and Fisher, K.H. (1964).

Energy metabolism. In 'Nutrition, A comprehensive Treatise'. (G.H. Beaton and E.W. McHenry, Eds.), Academic Press, New York, Vol. I, p. 182-257.

Szadkowski, D., Joergensen, A., Essing, H.G. and Schaller, K.H. (1970).

The rate of creatinine excretion as a reference value for the analysis of urine samples. I. The effect of daily urine volume and circadian rhythm on the excretion of creatinine. Z. Klin. Chem. Klin. Biochem., 8, 529-530.
cf : Biol. Abstr., 52, # 35311, (1971).

Talafant, E., Hoskova, A. and Pojervoa, A. (1977).

Negative feed back control of neonatal serum creatine and increased urinary creatinine excretion after enzyme inducing drugs. Pediatr. Res., 11, 880-882.

Talbot, N.B. (1936).

Basal energy metabolism and creatinine in the urine.

I. Observation on children. Am. J. Dis. Child, 52, 16.

Talbot, N.B. (1938).

Measurement of obesity by the creatinine coefficient.

Am. J. Dis. Child., 55, 42-58.

Talbot, N.B., Worcester, J. and Stewart, A. (1939).

New creatinine standard for basal metabolism and its clinical application. Am.J. Dis. Child., 58, 506-511.

Tanner, J.M., Heally, M.J.R., Whitehouse, R.H. and Edgson, A.C.

(1959). The relation of body build to the excretion of 17-ketosteroid and 17-ketogenic steroid in healthy young men. J. Endocrin., 19, 87.

Taylor, R.D. and Swartwort, J.R. (1967).

Biochemical survey of protein sufficiency during pregnancy in urban women. Obst. Gynecol., 29, 244-246.

Ten- State Nutrition Survey Reports, I-V (1972) Centre for Disease Control, Atlanta, Georgia.

Terroine, E.F. and Sorg-Matter, H. (1927).

Loi quantitative de la depense axotee minima des homeo-thermes: validite interspecifique. Arch. Intern. Physiol., 29, 121-132. cf : Mitchell, H.H. (1962).

Thanangkul, O. and Whitaker, J.A. (1966).

Childhood thiamine deficiency in northern Thailand. Am. J. Clin. Nutr., 18, 275.

Theophilus, F. and Barnes, E.H. (1974).

Diet and urolithiasis in rats with special reference to Ca:P ratio. Baroda J. Nutr., 1, 95-99.

Tolbert, B. and Watts, J.H. (1963).

Phenylalanine requirements of women consuming a minimal tyrosine diet and the sparing effect of tyrosine on the phenylalanine requirement. J. Nutr., 80, 111-116.

Tripathy, K., Khar, S. and Lotero, H. (1970).

Utilization of exogenous urea nitrogen in malnourished adults. Metabol, Clin. Exptl., 19, 253-262.

Tyler, F.H. (1972).

Muscular dystrophies. In 'The Metabolic Basis of Inherited Disease'. (J.B. Stanbury, J.B. Wyngaarden and D.S. Fredrickson, Eds.), Mc Graw-Hill Book Company, A Blakiston Publication, 3rd edn., p. 1210.

Unglaub, E.G. and Goldsmith, G.A. (1954).

Evaluation of vitamin adequacy: urinary excretion tests.

In 'Methods for Evaluation of Nutritional Adequacy and Status'. A symposium, Advisory Board on Quartermaster Research and Development Committee on Foods, National Academy of Sciences, National Research Council, Washington, D.C., p. 73.

Upadhyay, C.M. (1974).

Effects of dietary variations on bone composition in rats.
 Doctoral thesis, M.S.University, Baroda.

Van Hoogenhuyze, C.J.C. and Verpoelgh, H. (1906).

Beobachtungen Uber die Kreatinin auscheidung beim
 Menschen. Z. Physiol. Chem., 46, 415-471.

cf: Keys, et al (1950).

Vanpilsun, J.F. and Seljeskog, K.L. (1958).

Longterm ~~and~~ endogenous creatinine clearance in man.
 Proc. Soc. Exptl. Biol. Med., 97, 270-272.

Vasant Gadkar, Venkatachalam, P.S. and Tulpule, P.G. (1963).

Partition of urinary nitrogen in children with kwashiorkor
 treated with animal and vegetable protein.
 Am. J. Clin. Nutr., 12, 150.

Venkatachalam, P.S. (1962).

Maternal nutritional status and its effect on the newborn.
 Bull. W.H.O., 26, 193-201.

Vestergaard, P., Leverett, R. and Grangeburg, N.Y. (1958).

Constancy of urinary creatinine excretion.
 J. Lab. Clin. Med., 51, 211-218.

Viteri, F.E. (1972).

Creatinine-height index in malnourished children.

Nutr. Revs., 30, 24-26.

Viteri, F.E. and Alvarado, J. (1970).

The creatinine height index: its use in the estimation of degree of protein depletion and repletion in protein calorie malnourished children. Pediatrics, 46, 696-706.

Viteri, F.E., Alvarado, J. and Alleyne, G.A.O. (1971).

Reply to Drs. Mendez and Buskirk.

Am. J. Clin. Nutr., 24, 386-387.

Viteri, F.E., Mata, L.J. and Behar, M. (1973).

Methods de evaluation del estado nutricional. Protein-calorico en pre-escolares de condiciones-socio-economicas differetes. Arch. Latinoam. Nutr., 23, 13.

cf : Ghadami, H. (1975).

Vysotskii, V.G., Agureev, A.N. and Brantova, S.S. (1975).

The character of excretion of end products of nitrogen metabolism in urine in human protein deficiency.

Voprosy Pitaniya, No. 5, 27-35.

cf : Nutr. Abstr. Revs., 46, ~~99~~ 9985, (1976) p.1099.

Vysotskii, V.G., Vlasova, T.F., Kochetkova, A.N., Ushakov, A.S. and Shishkina, S.K. (1974).

Composition of urine and faeces of healthy men.

Voprosy Pitaniya, No. 6, 35-38.

cf : Nutr. Abstr. Revs., 45, ~~86~~ 8690 (1975).

Wadsworth, G.R. (1959).

Proc. Nutr. Soc., 18, iv.

cf : Kiriyama, S. (1970).

Watanabe, R. and Sassa, R. (1914).

Die Haranalyse wahrend des zweiwochigen hungerns eines Mannes. Z. Biol., 64, 373-408.

cf : Keys et al (1970).

Waterlow, J.C. (1969).

The assessment of protein nutrition and metabolism in the whole animal, with special reference to man. In 'Mammalian Protein Metabolism' (H.N. Munro, Ed.), Academic Press, New York, Vol. III, Chapter 28.

Waterlow, J.C., Neale, R.J., Rowe, L. and Palin, I. (1972).

Effects of diet and infection on creatine turnover in the rat. Am. J. Clin. Nutr., 25, 371-375.

Watson, R.L. and Langford, H.G. (1970).

Usefulness of overnight urines in population groups.

Pilot studies of sodium, potassium and calcium excretion.

Am. J. Clin. Nutr., 23, 290-304.

West, E.S., Todd, W.R., Mason, H.S. and Van Bruggen, J.T. (1966)

Text Book of Biochemistry, The Macmillan Company,

Collier-Macmillan/London, 4th edn.

Whitehead, R. (1965).

Hydroxyproline creatinine ratio as an index of nutritional status and rate growth. Lancet, 2, 567-570.

Whitehead, R.G. (1966).

Urinary excretion of hydroxyproline in kwashiorkor.

Lancet, 1, 203-204.

Whitehead, R.G. (1967).

Biochemical tests in differential diagnosis of protein and calorie# deficiencies. Arch. Dis. Child., 42, 479.

Whitehead, R.G. (1969).

The assessment of nutritional status in protein-malnourished children. Proc. Nutr. Soc., 28, 1-16.

Widdowson, E.M. and Mc Cance, R.A. (1957).

Effect of low-protein diet on the chemical composition of the bodies and tissues of young rats. Brit. J. Nutr., 11, 198-206.

Widdowson, E.M. and Mc Cance, R.A. (1970).

Use of the random specimens of urine to compare dietary intake of African and British children. Arch. Dis. Child., 45, 547-552.

Williams, R.R. (1961).

Toward the conquest of Beri-beri, Harvard University Press Cambridge, Massachusetts.

Willocks, J. (1977).

'Symposium on nutrition and the foetus'. The assessment of foetal growth. Proc. Nutr. Soc., 36, 1-7.

Wilson, C.S. et al (1964).

A review of methods used in nutrition surveys conducted by the Inter Departmental Committee on Nutrition for National Defence (ICNND). Am. J. Clin. Nutr., 15, 29.

Wolthuis, F.A. (1961).

Balance studies on protein metabolism in normal and uremic man, Scheltema en Holkema, Amsterdam.

cf : Gomen, H.A.P.C. (1967).

Wuest, H.M., Furness, F.N., White, E.W. and Beckel, W.S. (1962)

Eds. ~~of~~ Unsolved problems of thiamine.

Ann. N.Y. Acad. Sci., 98, 384.

Yamada, T. (1972).

Could an increased output of nitrogen in sweat result in an increase of protein requirements in muscular exercise.

Summaria, IX. Intern. Congr. Nutr., Mexico.

Yokota, F. (1964).

Japan J. Nutr., 22, 193.

cf : Kiriyama, S. (1970).

Young, C.M., Blondin, J., Tensuan, R. and Fryer, J.H. (1963).

Body composition studies of older women, thirty to seventy years of age. Ann. N.Y. Acad. Sci., 110, 589-607.

Young, V.R. and Scrimshaw, N.S. (1968).

Endogenous nitrogen metabolism and plasma free aminoacids in young adults given a protein free diet.

Brit. J. Nutr., 22, 9-20.

Young, V.R., Haverberg, L.N., Blimazes, C. and Munro, H.N. (1973).

Potential use of 3-methyl histidine excretion as an index of progressive reduction in muscle protein catabolism during starvation. Metabol. Clin. Exptl. 22, 1429-1436.

- Young, V.R., Taylor, Y.S.M., Rand, W.M. and Scrimshaw, N.S.
(1973). Protein requirements of man: Efficiency of egg protein utilization at maintenance and sub-maintenance levels in young men. J. Nutr., 103, 1164-1173.
- Yess, N., Price, J.M., Brown, R.R., Swan, P.B. and Linkswiler, H.
(1964). Vitamin B₆ depletion in man: urinary excretion of tryptophan metabolites. J. Nutr., 84, 229-236.
- Yuile, C.L., Lucas, F.V., Olson, J.P. and Shapiro, A.B. (1959).
Plasma~~X~~ protein turnover and tissue exchange.
J. Exptl. Med., 109, 173.
- Zorab, P.A. (1969).
Normal creatinine and hydroxyproline excretion in young persons. Lancet, 2, 1164.
- Zorab, P.A., Clark, S, and Harrison, A. (1969).
Creatinine excretion. Lancet, 2, 1254.