

BIBLIOGRAPHY

- Abdul-Alim, M.A. (1971). A chemical study of the leaves of Clerodendron inerme. Planta-Med. 19(4): 318-321.
- Agarwal, S.C., Sarangadharan, M.G. and T.R. Seshadri. (1965). Colouring matter of Teak (Tectona leaves): Isolation and constitution of Tecto-leaf quinone. Tetrahedron Lett. 30: 2623-2626.
- Ahmed, Z.F., EL-Moghazy Shoaib, A.M., Wassel, G.M. and S.M.EL Sayyad. (1972). Phytochemical study of Lantana camara. Planta-Med. 21(3): 282-288.
- Airy Shaw, H.K. (1952). Notes on the taxonomic position of Nyctanthes L. and Dimetra Kerr. Kew Bull. 1952: 271-272.
- Alston, R.E. and Irwin, H.S. (1961). On comparative extent of variation of free amino acids and certain secondary substances in Cassia sp. Amer. J. Bot. 48: 35-39.
- Alston, R.E. and Turner, B.L. (1963). "Biochemical Systematics". New Jersey.
- Alston, R.E. (1967). Biochemical systematics. Evol. Biol. 1: 197-305.
- Amarsingham, R.D., Bisset, N.G., Millard, A.H. and M. C. Woods. Phytochemical survey of Malaya part III. Alkaloids and saponins. Eco. Bot. 18(3): 270.
- Anjaneyulu, A.S.R., Rao, K.J., Rao, V.K., Row, L.K., Subramanian, C., Pelter, A. and R.S. Ward. (1975). The structure of Lignans from Gmelina arborea. Tetrahedron, 31(10): 1277-1285.

- Anjaneyulu, A.S.R., Rao, A.M., Rao, V.K., Row, L.R. and A. Pelter. (1977). Novel hydroxy lignans from the heartwood of Gmelina arborea. Tetrahedron 33(1): 133-144.
- Apparao, M. and Rao, E.V. (1978). Chemical components of Adenocalymma aliaceum. Indian J. Pharm. Sci. 40(6):224.
- Avadhoot, Y., Dixit, V.K. and K.C. Varma. (1978). Preliminary chemical examination of seed of Lantana camara var. aculeata. Indian Drugs Pharm. Ind. 18(3): 23-24.
- Banerji, A., Chadha, M.S. and V.G. Malshit. (1969). 5-OH, 3,6,7,3',4'-penta methoxy flavone from Vitex negundo. Phytochemistry, 8: 511.
- Banerjee, S.K., Chakravarti, R.N., Sachdev, K.S. and S.A. Vasavada. (1969). Constituents of the root bark of Clerodendron serratum. Phytochemistry 8: 515.
- Barua, A.K., Sanyal, P. and P. Chakrabarti. (1967). Steroid glucoside from Clerodendron infortunatum. J. Indian Chem. Soc. 44(6): 549-551.
- Barua, A.K., Chakrabarti, and P.K. Sanyal. (1969). Nodifloretin-A. New flavone from Lippia nodiflora. J. Indian Chem. Soc. 46(3): 271-272.
- Basu, N.K. and Singh, G.B. (1944). Indian J. Pharm. 6: 71.
- Bate-Smith, E.C. and Swain, T. (1966). The asperulosides and the aucubins - In Swain, T. (ed.): "Comp. Phytochem." 159-174. London.
- Bate-Smith, E.C. (1972). Chemistry and phylogeny of the angiosperms. Nature. 236 : 353-354.

- * Bendz, G. and Santesson, J. (eds.) (1973). Chemistry in Botanical classification. Nobel Symposium 25, New York and London.
- Benson, L. (1957). "Plant Classification". Boston.
- Bentham, G. and Hooker, J.D. (1865). Bignoniaceae in "Genera Plantarum". Vol. 2. London.
- Bentham, G. and Hooker, J.D. (1876). Verbenaceae in "Genera Plantarum". Vol. 2: 1131-1137. London.
- Bessey, C.E. (1915). The phylogenetic taxonomy of flowering plants. Ann. Missouri Bot. Gard. 2: 109-164.
- Bhattacharjee, A.K. and Das, A.K. (1969). Phytochemical survey of few Mysore plants. Eco. Botany 23(3): 274-276.
- Block, R.J., Durrum, E.L. and G. Zweig. (1958). A manual of paper chromatography and paper electrophoresis. 2nd ed. Academic Press. New York.
- Bohm, B. and Wilkins, D. (1978). Chemosystematic studies in the Saxifragaceae sensu lato. X. Flavonoids in Elmira racemosa. Brittonia. 30: 327-333.
- Boivin, B. (1956). Les Familles de Tracheophytes. Bull. Soc. Bot. France 103 : 490-505.
- * Bournot, K. (1914). Arch. Pharm. 25 : 351.
- Briquet, J. (1897). Verbenaceae in Engler and Prantl 'Die naturlichen pflanzenfamilien'. IV 3b: 132-182. Leipzig.
- Britton, N.L. (1920). Jour. N.Y. Bot. Gard. 21 : 72-73.
- Campbell, D.H. (1930). The relationships of Paulownia. Bulletin of the Torrey Botanical Club. 57: 47-50.

- Chapman, E., Perkin, A.G. and R. Robinson. (1927). J. Am. Chem. Soc. 3015.
- Chaudhary, A., Bhattacharya, A., Mitra, S.R. and N. Aditya Chaudhary. (1978). Phytochemical investigation on the leaves of Callicarpa macrophylla. Jr. Indian Chem. Soc. 55(6): 628-629.
- Chowdhary, K.A. (1964). Growth rings in tropical trees and Taxonomy. The J. Indian Bot. Soc. Vol. XLIII(2):334-342.
- Clarke, C.B. (1884). In "Flora of British India" (J.D.Hooker). Vol. IV: 376-386.
- Clarke, C.B. (1885). In "Flora of British India" (J.D.Hooker). Vol. IV.
- * Cognin, M.A.H., Gomes, C.M.R., Gottlieb, O.R., Marx, M.C., Rocha, A.I.D.A., De Silva, G.F., Das, M.F. and J.A. Temperini. (1977). Biochemical Systematics : Methods and Principles, Pl. Syst. Evol. Suppl. 1: 53-56.
- * Cosinovi, C.G., Marini-Bettolo, G.B., Dalima, O.G., Dalamaia, M.E. and I.L. Dalbbuquerque (1963). One quinone isolated from the wood of Tabebuia avellonedae (Bignoniaceae). Rend. Ist. Super. Sanita. 26(1/2): 5-10.
- Crawford, D.J. and Dorn, R.D. (1974). "Numerical Chemotaxonomy" and other aspects of chemosystematics. Taxon 23(2/3): 331-338.
- Crawford, D.J. and Mabry, T.J. (1978). Flavonoid chemistry of Chenopodium fremontii : infraspecific variations and systematic implications at the interspecific level. Biochem. Syst. and Ecol. 6: 189-192.
- Crete, P. (1955). L'application der certaines donnees embryo-

- logiques a la systematique des Orobanchacees et de quelques familles voisines. Phytomorphology (Delhi) 5: 422-435.
- Cronquist, A. (1968). "The Evolution and Classification of flowering plants". Houghton Mifflin, Boston.
- Cronquist, A. (1975). Some thoughts on angiosperm phylogeny and taxonomy. Ann. Missouri Bot. Gard. 62: 517-520.
- Cronquist, A. (1977). On the taxonomic significance of secondary metabolites in angiosperms. Plant Syst. Evol. Suppl. 1: 179-189.
- Czaza, A.Th. (1978). Structure of starch grains and the classification of vascular plant families. Taxon, 27 (5/6): 463-470.
- Damtoft, S., Jensen, S. and B.J.Nielson. (1979). Iridoids in Verbena. Taxon. 28(5/6): 525-528.
- Daniel, M. and Sabnis, S.D. (1979). Chemotaxonomy of Loganiaceae. Curr. Sci. 48: 383-385.
- Daniel, M. and Sabnis, S.D. (1979). Chemotaxonomy of Oleaceae. Indian J. of Expt. Biol. 17(9): 995-997.
- Daniel, M. and Sabnis, S.D. (1981). A numerical evaluation of the taxonomic status of Nyctanthes. L. Indian J. Bot. (Prof. Deshpande Commemoration Vol. (in press).
- Darrah, W.C. (1939). Text book of Palaeobotany. Appleton Century, New York.
- DA, Silveira, Jose Chagas, Gottlieb, O.R. and Geovane, G.De Oliveira. (1975). Zeyherol, a dilignol from Zeyhera digitalis. Phytochemistry 14(8): 1829-1830.

- Davis, G.L. (1966). "Systematic embryology of the Angiosperms", John Wiley and Sons, IC. New York.
- Davis, P.H. and Heywood, V.H. (1967). "Principles of Angiosperm Taxonomy". Oliver and Boyd. Edinburgh.
- Desai, H.K., Gawad, D.H., Govindachari, T.R., Joshi, B.C., Kamat, V.N., Modi, J.D., Parthasarthy, P.C., Radhakrishnan, J., Shanbhag, M.N., Sidhaye, A.R. and N. Viswanathan. Chemical investigation of Indian plants. Part VII. Indian J. Chem. 2: 840.
- Dey, A.K., Mukherjee, A., Das, P.C. and A. Chatterjee. (1978). Occurrence of aloe-emodin in the leaves of Oroxylum indicum. Indian J. Chem. Sect. B. Org. Chem. incl. Med. Chem. 16 B(11): 1042.
- Dhruwa, B.R., Ramarao, A.V., Srinivasan, R. and K. Venkataraman. (1972). Structure of a quinone from Teak tissue culture. Indian J. Chem. 10(7): 683-685.
- Dickinson, E.M. and Jones, G. (1969). Pyridine alkaloids from Tecoma stans. Tetrahedron, 25(7): 1523-1529.
- * Dietrichs, H.H. (1964). Desoxylapachol in Tabebuia species. Naturwissenschaften, 51(17): 408-409.
- * Dohnal Barbara (1977). Investigation on some metabolites of Tecoma stans. Juss. callus tissues. III. Chromatographical search for iridoids, phenolic acids, terpenoids and sugars. Acta Soc. Bot. Pol. 46(2): 187-200.
- * Duret, S.H., Jacquenin, and R.R. Paris. (1976). The chemical composition of Stachytarpheta jamaicensis (L.) Vahl (= S. indica Vahl) Verbenaceae. Planta. Med. Phytother. 10(2): 96-104.

- Endlicher, S.L. (1836-40). Genera Plantarum secundum orolines naturales disposita. Vindobonae.
- Endlicher, S.L. (1839). "Genera plantarum". 678.
- Engler, A. and Diels, L. (1936). Engler's Syllabus der Pflanzenfamilien. 11th ed. edited by L. Diels, Berlin.
- Erdtman, G. (1952). Pollen morphology and plant taxonomy. Angiosperms. Almquist and Wiksell, Stockholm.
- Ford, C.W. and M. Robin Bendall. (1980). Identification of the iridoid glucoside theveside in Lantana camara and determination of its structure and stereochemistry by means of NMR. (Cited from Biol. Abst. 70(10). Ref. No. 67403).
- Forsynth, W.G.C. and Simmonds, N.W. (1954). Proc. R. Soc. 142 : 549.
- Gardner, R.O. (1977). Systematic distribution and ecological function of the secondary metabolites of the Rosidae, Asteridae. Biochem. Syst. Ecol. 5: 26-35.
- * Garnier, J. (1979). Chemical study of 2 Verbenaceae of Guina: Stachytarpheta guyanensis Vahl and S. mutabilis Vahl. Plant Med. Phytother. 11(4): 303-305.
- Gates, F.C. (1940). Unnoted list of the plants of Kansas : Ferns and flowering plants. (cited from R.D.Gibbs, Vol. IV).
- Geissman, T.A. (ed.) (1962). "Chemistry of the flavonoid compounds". Pergamon Press. Oxford.
- Gentry, A.H. (1974). Flowering phenology and diversity in tropical Bignoniaceae. Biotropica, 6: 64-68.

- Gentry, A.H. (1979). Bignoniaceae. Part I. Crescentieae and Tourrettieae. Flora Neotropica Monograph 23.
- Ghatak, J. (1956). A contribution to the life history of Oroxylum indicum. Proc. Indian Acad. Sci. 43 B: 72-87.
- Gibbs, R.D. (1958). Chemical Evolution in Plants. J. Linn. Soc. London (Bot.) 56 (365): 49-57.
- Gibbs, R.D. (1974). "Chemotaxonomy of flowering plants". McGill Queens, Univ. Press, Montreal. London.
- Goldblatt, P. (1976). New or noteworthy chromosome records in the Angiosperms. Ann. Missouri Bot. Gard. 63: 889-895.
- Goldblatt, P. and Gentry, A.H. (1979). Cytology of Bignoniaceae. Bot. Notiser 132: 475-482.
- Gornall, R.J. and Bohm, B.A. (1978). Angiosperm Flavonoid evolution : A Reappraisal. Systematic Botany 3(4): 353-368.
- Gornall, R.J., Bohm, B.A. and R. Dahlgren (1979). The distribution of flavonoids in angiosperms. Bot. Notiser 132 : 1-30.
- Govindachari, T.R., Patankar, S.J. and N. Viswanathan. (1971). Isolation and structure of two new Dihydroisocoumarins from Kigelia pinnata. P hytochemistry 10(7): 1603-1606.
- Govindachari, T.R., Parthasarthy, P.C. and H.K. Desai (1972). Arboreol, a new lignan from Gmelina arborea. Indian J. Chem. 10(12): 1120-1122.
- Govindu, H.C. (1950). Studies in the embryology of some members of the Bignoniaceae. Proc. Indian Acad. Sci. 32: 164-178.

- Cross, D., Berg, W.W. and H.R. Schuette. (1973). Δ^8 -dehydrostytanthin and δ^7 -skytanthin in Tecoma stans. Phytochemistry 12(1): 201-202.
- Gujral, V.K., Gupta, S.R. and K.S.Verma. (1979). New chromone glucosides from Tecomella undulata. Phytochemistry, 18(1): 181-182.
- Gundersen, A. (1950). "Families of Dicotyledons". Wattham, Mass.
- Gupta, S.R., Sureshchandra, and V. Mahadevan. (1967). Chemical examination of Clerodendrum phlomidis. Indian J. Pharm. 29(3): 102-103.
- Gupta, S.R., Malik, K.K. and T.R. Seshadri. (1969). Lapachol from the heartwood of Tecoma undulata and a note on its reaction. Indian J. Chem. 7(5): 457-459.
- Gupta, S.S. and Nanda, K. (1978). Studies in Bignoniaceae.I. Ontogeny of dimorphic anther tapetum in Pyrostegia. Amer. J. Bot. 65: 395-399.
- Hallier, H. (1912). L'origine et le systeme phyletique des Angiospermes exposes a laide de leur arbre geneologique. Arch. Neerl. Sci. Exctes et Nat Ser. III. B. 1: 146-234. (Cited from R.D. Gibbs (1974). Vol. IV).
- * Hammouda, Y., Michel, P. and Jean Lemen. (1963). Isolation of Tecostanine. Alkaloids of Tecoma stans (Bignoniaceae). Ann. Pharm. Franc. 21(9/10): 699-702.
- * Hammouda, Y. and Khalafaliah, N. (1971). Extraction studies on the leaves of Tecoma capensis. Pharmazie, 26(10): 640-642.
- Hansel, R., Leuchert, C., Rimpler, H. and K.D. Schaaf. (1965). Phytochemistry 4: 19.

- Harborne, J.B. (1967). "Comparative Biochemistry of flavonoids". Academic Press, London.
- Harborne, J.B. (1967). Comparative biochemistry of the flavonoids VI. Flavonoid pattern in Bignoniaceae and Gesneriaceae. Phytochemistry, 6: 1643-1651.
- Harborne, J.B., Boulter, D. and B.L. Turner (eds.). (1971). "Chemotaxonomy of the Leguminosae". London and New York.
- Harborne, J.B. and Williams, C.A. (1971). 6-Hydroxy luteolin and Scutellarein as phyletic markers in higher plants. Phytochemistry, 10: 367-378.
- Harborne, J.B. (1972). Evolution and function of flavonoids in plants. Recent Adv. Phytochem. 4: 107-141.
- Harborne, J.B. (1973). Phytochemical Methods : A Guide to Modern Techniques of Plant Analysis. Chapman and Hall. London.
- Harborne, J.B. (1973). Flavonoids. In: L.P. Miller (ed.). Phytochemistry. Vol. III, pp. 344-380. Van Nostrand Rheinhold, New York.
- Harborne, J.B. (1975). The Biochemical systematics of flavonoids. In: J.B. Harborne, T.J. Mabry and H. Mabry (eds.). "The Flavonoids", pp. 1056-1095. Chapman and Hall, London.
- Harborne, J.B. (1977). Flavonoids and the evolution of the angiosperms. Biochem. Syst. Ecol. 5: 7-22.
- * Hegnauer, R. (1966). Aucubinartige Glucoside. Uber die Verbreitung und Bedeutung als systematisches Merkmal. Pharm. Acta Helv. 41: 557-587.

- Hegnauer, R. (1969). Chemical evidence for the classification of some plant taxa. - In Swain, T. and Harborne, J.B. (eds.). "Perspectives in Phytochemistry". 1: 121-138.
- * Hegnauer, R. (1971). Pflanzenstoffe und Pflanzensystematik. Naturwiss. 58: 585-598.
- Heywood, V.H. (1973). The role of chemistry in plant systematics. Pure and applied chemistry. 34: 355-375.
- Holttum, R.E. (1967). Comparative morphology, taxonomy and evolution. Phytomorphology, 17(1-4): 36-41.
- Hungund, B.L. and Pathak, C.H. (1971). U.S.D.A. Forest Service Research Paper NE-201.
- Hutchinson, J. (1926). "The families of flowering plants". Volume 1 Dicotyledons. Clarendon Press, Oxford.
- Hutchinson, J. (1959). "The families of flowering plants" 2nd ed. Oxford University Press, New York. Bignoniaceae Vol. 1: 389, Verbenaceae 1: 395.
- Hutchinson, J. (1969) "Evolution and Phylogeny of flowering plants". Academic Press, London.
- Hutchinson, J. (1973). The families of flowering plants, arranged according to a new system based on their probable phylogeny. 3rd ed.
- Ibrahim, R.K. and Towers, G.H.N. (1960). Arch. Biochem. and Biophys. 87 : 125.
- Ibrahim, R.K., Towers, G.H.N. and R.D. Gibbs. (1962). Syringic and sinapic acids as indicators of differences between major groups of vascular plants. J. Linn. Soc. Lond. (Bot.). 58: 223-230.
- Inamdar, J.A. (1967). Studies on the trichomes of some Oleaceae, structure and ontogeny. The Proc. of Indian Acad. Sci. 66: 164-177.

- * Inoue, K., Shiobara, Y., Chang, Chen, C., Sakuyama, S.
and H. Inouye. (1979). Quinones and related compounds
in higher plants : 7th Supplementary. Studies on the
constituents of the wood of Catalpa ovata. Yaku Gaku
Zasshi. 99(5): 500-504.
- * Ismailov, N.M. (1967). Ecological floristic analysis of
alkaloid bearing plants found in Talysh. RAST. RESUR.
3(1): 53-57.
- * Iyer, A. and Joshi, B.C. (1974). Chemical investigation of
the fruit of Kigelia pinnata. Part I. Herba Pol. 20(4):
319-320.
- Jafri, S.M.H. and Ghafoor, A. (1974). Verbenaceae in Flora
of Pakistan..(ed.) Nasir, E. and Ali, S.I. 77: 1-40.
- Jain, D.K. (1978). Studies in Bignoniaceae.III. Leaf
Architecture. J. Indian Bot. Soc. 57: 369-386.
- * Jaurez, B.E. and Seeugmann, P. (1978). Chemosystematic study
of the Flavonoid distribution in the Argentine species of
genus Tecoma (Bignoniaceae). Lilloa. 35(1): 15-22.
- Jensen, S.R., Nielsen, B.J. and R. Dahlgren. (1975). Iridoid
compounds, their occurrence and systematic importance
in the Angiosperms. Bot. Notiser. 128: 148-180.
- Joshi, A.B. and Hardas, M.W. (1956). Ploidy in two
bignoniaceous garden climbers. Indian J. Genet. Plant
Breeding 16: 57-59.
- Joshi, B.C. and Bakuni, D.S. (1959). Chemical examination of
Lippia nodiflora. J. Sci. Industr. Res. 18B: 525-527.
- Joshi, B.C. (1966). Abstract of the 54th session. Indian
Science Congress Association. 161.

- * Joshi, K.C. and Singh, L.B. (1970). Extractives from heart woods. II. Isolation of Gmelinol and n-octacosanol from Gmelina arborea. Z. NATURFORSCH. TEIL-B. 25B(7):693-694.

- * Joshi, K.C., Prakash, L., Bansal, R.K. and P. Singh. (1973). Mass spectrometric studies of dehydro- α -lapachone and dehydro-iso- α -lapachone. from Tabebuia rosea. Z. Naturforsch Teil. C. Biochem. Biophys. Biol. Virol. 28(11/12): 646-649.

- Joshi, K.C., Prakash, L. and P. Singh. (1973). Quinones and other constituents from Phyllarthron comorense. Phytochemistry, 12 : 469-470.

- Joshi, K.C., Prakash, L. and P. Singh. (1973). Quinones and other constituents from Tabebuia rosea. Phytochemistry, 12 : 942.

- Joshi, K.C., Prakash, L. and L.B.Singh. (1975). 6-O Veratryl Catalposide : A new iridoid glycoside from Tecomella undulata. Phytochemistry, 14(5/6): 1441-1442.

- Joshi, K.C., Bansal, R.K., Singh, P. and Girraj-Singh. (1975). Components of the stem bark of Phyllarthron comorense and Jacaranda mimosaefolia and roots of Desmodium pulchellum. Indian J. Chem. 13(8): 869-870.

- Joshi, K.C., Singh, P. and G. Singh. (1976). Crystalline components of roots of Phyllarthron comorense and stem bark of Tabebuia rosea. Indian J. Chem. Sect. B. Org. Chem. Incl. Med. Chem. 14(8): 637-638.

- Joshi, K.C., Singh, P. and R.T. Pardasani. (1977). Quinones and other constituents from the roots of Tecomella undulata. Planta-Med. 31(1): 14-16.

- Joshi, K.C., Singh, P. and R.T. Pardasani. (1977). Chemical components of the roots of Tectona grandis and Gmelina arborea. Planta-Med. 32(1): 71-75.
- Joshi, K.C., Prakash, L. and R. Shah. (1977). Chemical examination of the roots of Tabebuia rosea and heartwood of Oroxylum indicum. Planta-Med., 31(3): 257-258.
- Joshi, K.C., Singh, B. and R.T. Pardasani. (1978). Chemical constituents of stem heartwood of Markhamia stipulata. Planta-Med., 34(2): 219-221.
- Joshi, K.C., Singh, P., Pardasani, R.T. and B. Singh. (1979). Quinones and other constituents from Haplophragma adenophyllum. Planta-Med., 37(1): 60-63.
- Joshi, K.C., Singh, P. and Alka Mehra. (1979). Chemical investigation of the roots of different Clerodendrum species. Planta-Med., 37(1): 64-66.
- Joshi, K.C., Prakash, L., Singh, L.B. and SR. P. Rastogi. Paper presented at the 8th International Symposium of the chemistry of natural products. Abstract 163.
- Joshi, V., Merchant, J.R., Nadkarny, V.V., Namboori, K. and D.D. Vaghani. (1974). Chemical components of some Indian Medicinal Plants. Indian J. Chem. 12(2): 226.
- * Juarez, B.E. and Seeugmann, P. (1978). Chemosystematic study of the flavonoid distribution in the Argentine sp. of genus Tecoma (Bignoniaceae). Lilloa. 35(1): 15-22.
- * Junell, S. (1934). Zur Gynaceummorphologie und systematik der Verbenaceen and Labiaten. Symb. Bot. Upsal. 4: 1-219.
- Kameswaramma, A. and Sheshadri, T.R. (1947). Proc. Indian Acad. Sci. 25 A : 43.

- Kapil, R.S. (1960). J. Indian Chem. Soc. 37: 697.
- Kapil, R.N. and Vani, R.S. (1966). Nyctanthes arbor-tristis Linn.: Embryology and relationships. Phytomorphology, 16(4): 553-563.
- Kate, T.L. (1978). The male and female gametophyte of Tecoma radicans. Proc. Indian Sci. Congress Association. Abst. : 78.
- Kedharnath, S. (1950). Chromosome numbers of plants. A note on the chromosome number of some plants. Indian J. Genet. Plant Breed. 10: 96.
- Kodanda Rao, E., Rao, E.V. and D.V. Rao. (1977). Phenolic constituents of the bark of Vitex negundo. Indian J. Pharm. 39(2): 41.
- Kshetrapal, S. and Tiagi, Y.D. (1970). Structure, vascular anatomy and evolution of the gynoecium in family Oleaceae and their bearing on the systematic position of genus Nyctanthes L. Acta Bot. Acad. Sci. Hung. Tomas 16 (1-2), 143-151.
- Kubitzki, K. (1969). Chemosystematische Betrachtungen zur Grossgliederung der Dicotylen. Taxon. 18: 360-368.
- Kutney, J.P. and Hanssen, H.W. (1971). 5,6,7,tri-methoxy and 5,6,7,8 tetra-methoxy flavone from Zeyhera tuberculosa. Phytochemistry, 10(12): 3298-3302.
- Lawrence, G.H.M. (1951). "Taxonomy of Vascular Plants". Oxford.
- Lindley, J. (1847). The Vegetable Kingdom. 2nd ed. London.
- Mabry, T.J., Markham, K.R. and M.B. Thomas. (1970). The systematic identification of flavonoids. Springer-Verlag, Berlin.

- Maheshwari, P. and Kapil, R.N. (1966). Some Indian contributions to the embryology of Angiosperms. Phytomorphology, 16: 239-291.
- Maheshwari, J.P. and Banerjee, S.K. (1970). Isolation of β -sitosterol from Tecoma stans. Indian J. Pharm. 32(6): 159.
- * Mangelot, S. and Mangelot, G. (1962). Enquete sur les nombres chromosomiques dans une collection d'especes tropicales. Rev. Cytol. Biol. Veg. 25: 411-447.
- Manners, G.D., Jurd, L., Wong, R. and K. Palmer. (1975). Constituents of Tabebuia guayacan : The structure of guayacanine. Tetrahedron, 31(24): 3019-3024.
- Manners, G.D. and Jurd, L. (1976). A new naphthaquinone from Tabebuia guayacan. Phytochemistry, 15(1): 225-226.
- Manske, R.H.F. (1944). The alkaloids. Ann. Rev. Biochem. 13: 533-548.
- * Mauritson, J. (1935). Etwas uber die Embryologie der Bignoniaceen. Bot. Notiser : 60-77.
- * Meeuse, A.D.J. (1970). The descent of the flowering plants in the light of new evidence from phytochemistry and from other sources. I and II - Acta Bot. Neerl. 19: 61-72, 133-140.
- Mehra, P.N. and Bawa, K.S. (1969). Chromosomal evolution in tropical hardwoods. Evolution 23: 466-481.
- Melchior, H.A. (ed.) (1964). Tubiflorae In: Engler's Syllabus der pflanzenfamilien (12th, ed.), II Band Angiospermen Gebruder Borntrager, Berlin.

- Metcalfe, C.R. and Chalk, L. (1950). "Anatomy of the Dicotyledons", Vol. II. Oxford.
- Milz, S. and Rimpler, H. (1978). Pulchelloside II, a new iridoid of Verbena pulchella. Tetrahedron Lett. No.38. 3549-3552.
- * Milz, S. and Rimpler, H. (1979). Iridoids in Verbena and some other Verbenaceae. Z. Natur. Forsch. Sect. C. Biosci. 34(5/6): 319-329.
- Misra, G.S. and Subramanian, P.M. (1980). 3 new flavone glycosides from Vitex negundo. Planta-Med. 38(2): 155-160.
- Mitra, K. (1968). Pollen morphology in Bignoniaceae in relation to taxonomy. Bull. Bot. Surv. India 10(3/4): 319-326.
- * Moldenke, H.N. and Moldenke, A.L. (1946). Brief historical survey of the Verbenaceae and the related families. Plant Life 2: 13-98.
- Moldenke, H.N. (1959). "A resume of the Verbenaceae, Avicenniaceae, Stilbaceae, Symphoremaceae and Eriocaulaceae of the world as to valid taxa, geographic distribution and synonymy". pp. 1-495, Publ. by the Author, New Jersey.
- Mukherjee, J. and Chanda, S. (1973). Biosynthesis of Avicennia L. in relation to taxonomy. Geophytology, 3: 85-88.
- Nair, P.K.K. and Rehman, K. (1962). Pollen grains of Indian plants - V. Verbenaceae. Bulletin of the Nat. Bot. Gard. No. 76.

- Nair, A.G.R., Ramesh, P., Nagarajan, S. and S.S.Subramanian. (1973). New flavone glycosides from Lippia nodiflora. Indian J. Chem. 11(12): 1316-1317.
- Nair, A.G.R. and Subramanian, S.S. (1975). Quercetagenin and other flavones from Gmelina arborea and G. asiatica. Phytochemistry, 14(4): 1135-1136.
- Nair, A.G.R., Vedantham, T.N.C., B. Kannabiran, (1979). Polyphenolic components of Clerodendrum serratum. Curr. Sci. 48(10): 440-441.
- Nanda, P.C. (1962). Chromosome numbers of some trees and shrubs. J. Indian Bot. Soc. 41: 271-277.
- Nanda, K. and Gupta, S.S. (1978). Endothecium in Pyrostegia venusta. Curr. Sci. 40 : 470-471.
- Nanda, K. and Gupta, S.S. (1978). Studies in the Bignoniaceae. II. Ontogeny of dimorphic anther tapetum in Tecoma. Amer. J. Bot. 65: 400-405.
- Narsinga, Rao, U. (1936). Chromosome number in Millingtonia hortensis Linn. f. (Bignoniaceae). Curr. Sci. 4: 654.
- Ogura, Y. (1964). Comparative morphology and classification of plants. Phytomorphology, 14: 240-247.
- * Okigawa, M., Hatanaka, H., Kavaró, N., Matsonaya, J. and Z. Tamura (1971). Chem. Pharm. Bull. 19(1): 148-152.
- Padmanabhan, D. (1960). The embryology of Avicennia officinalis. I. Proc. Indian Acad. Sci. (B). L11: 131-145.
- Pal, N. (1951). Studies in the embryology of some Verbenaceae. J. Indian Bot. Soc. 30: 59-74.

- Pandey, V.B. and Dasgupta, B. (1971). Chemical investigation of the bark of Tecomella undulata. Isolation of new ester glucoside Tecomine. J. Indian Chem. Soc. 48(10): 937-941.
- Perkin, A.G. (1900). J. Chem. Soc. 77: 422.
- * Plouvier, V. (1971). Research on heterosides; catalpol from Paulownia and Catalpa arbutoside from Sarbaria and Knautioside a spananoside from Knautia arvensis Coult. C.R. Hebd. Seances. Acad. Sci. Ser. D. Sci. Natur. (PARIS). 272(10): 1443-1446.
- Price, J.R. (1963). The distribution of alkaloids in the Rubiaceae. In: "Chemical Plant Taxonomy" T. Swain (ed.), pp. 429-452. Academic Press, London.
- Pulle, A. (1952). Compendium van de Terminologie, Nomenclatuur en Systematiek der Zaadplanten, 3rd ed. Utrecht. (Cited from R.D. Gibbs (1974) Vol. IV.).
- Raffauf, R.F. (1970). "A Handbook of Alkaloids and Alkaloidal containing plants". Wiley-Interscience. New York.
- Raghavan, T.S. and Venkatasubban, K.R. (1940). Studies in the Bignoniaceae. I. Chromosome number and epidermal hydathodes in Spathodea campanulata Beauv. J. Indian Bot. Soc. 19: 293-298.
- Rao, C.B. and Venkateswarlu, V. (1962). J. Sci. Indi. Res. 21 B: 313.
- Rao, Ch.B., Rao, T.N. and E.K.S. Vijaykumar. (1978). Chemical examination of the fruit of Duranta plumeri. Indian J. Chem. 16 B(9): 844-845.
- Rao, D.V., Rao, E.V. and N. Vishanadham. (1967). Occurrence of Luteolin in the leaves. of Gmelina arborea. Curr. Sci., 36(3): 71-72.

- Rao, D.V. and Rao, E.V. (1970). Chemical components of the leaves of Gmelina arborea. Indian J. Pharm. 32(5): 140-141.
- * Rao, P.S. (1965). Naturwissenschaften. 34: 262.
- Rao, V.S. (1952). The floral anatomy of some Verbenaceae with special reference to the gynoecium. J. Indian Bot. Soc. 31: 292-315.
- * Reichert, E.T. (1913-1919). Publicn. No. 173 and 270 respectively. Cornege Inst. Washington.
- * Rimpler, H. (1972). Phytoecdysones and iridoids from Vitex megapotamica. Arch. Pharm. (Weinheim.) 305(10): 746-751.
- * Rimpler, H. and Helmut, T. (1974). Iridoids and ecdysones from Verbenaceae. Z. Naturforsch. 29: 111.
- * Rimpler, H. and Schafer, B. (1979). Hastatoside a new iridoid from Verbena hastata and V. officinalis. Z. Naturforsch. Sect. C. Biosci. 34(5/6): 311-318.
- Rizvi, S.A.I. and Sultana, T. (1973). Chemical studies of a fixed oil, terpenoids and other compounds from the pods of Heterophragma adenophyllum. Planta-Med., 23(2): 125-131.
- Rendle, A.B. (1938). "The classification of flowering plants", 2nd ed. (1975) Vol. II (Dicotyledons). Bignoniaceae: 538-541; Verbenaceae: 500-505.
- * Runemark, H. (1968). Critical comments on the use of statistical methods in chemotaxonomy. Bot. Notiser 121: 29-43.

- * Sander mann, W. and Simatupang, M.H. (1965). New quinone from Tectona grandis. NATURWISSENSCHAFTEN. 52(10): 262-263.
- Sastry, M.S. and Mahadevan, V. (1963). Chemical investigation of Lantana camara. Curr. Sci. 32(2): 71.
- Saxena, M.R. (1973). Pollen morphological studies in the Verbenaceae. Ph.D. thesis, Osmania University, Hyderabad, India. (Cited from Saxena, M.R. 1981).
- Saxena, M.R. (1975). Pollen morphology of the Nyctanthoideae (Verbenaceae). J. Indian Bot. Soc. 54: 71-74.
- Saxena, M.R. (1981). Contribution to the palynotaxonomy of Avicenniaceae. Endl. J. Indian Bot. Soc. 60: 28-32.
- Schauer, J.C. (1847). Verbenaceae, in De Candolles prodramus. Vol. XI. Paris. (Cited from Saxena, 1981).
- Schuman, K. (1895). Bignoniaceae in Engler and Prantl 'Die natuerlichen pflanzenfamilien'. IV. 3b: 189-252.
- * Schnarf, K. (1931). "Vergleichende Embryologie der Angiospermen". Berlin.
- * Seemann, B. (1860). Synopsis Crescentiacearum : an enumeration of all the Crescentiaceous plants at present known. Trans. Linn. Soc. 23: 1-22.
- Sen, A.B. and Singh, S.P. (1964). Chemical examination of Clerodendron infortunatum (Verbenaceae). Indian J. Chem. 2(4): 172.
- Sen, M. and Sarkar, V. (1978). Chemical investigation of leaves of Callicarpa arborea (Verbenaceae). J. Indian Chem. Soc. 55(7): 744-745.

- Sharma, R.C., Zaman, A. and A.R. Kidwai. (1968). Chemical examination of Millingtonia hortensis. Phytochemistry 7(10): 1891-1982.
- * Siebold and Zuccharini (1835). Flora Japonica 1: 27.
- Simmonds, N.W. (1954). Chromosome behavior in some tropical plants. Heredity 8: 129-146.
- Singh, P., Prakash, L. and K.C. Joshi (1972). Lapachol and other constituents from Bignoniaceae. Phytochemistry 11(4): 1498.
- Sinha, N.K., Seth, K.K., Pandey, V.B., Dasgupta, B. and A.H.Shah (1981). Flavonoids from the flowers of Clerodendrum infortunatum. Planta-Med. 42(3): 296-298.
- * Skottsberg, C. (1940). Vaxternas L.V. Stockholm.
- Smith, D.C.C. (1955). p-Hydroxybenzoic group in the lignin of Populus tremula. J. Chem. Soc. 2347.
- * Soo, R. (1953). Die modernen Grundsätze der phylogenie im neuen system der Blütenpflanzen. Acta Biol. Acad. Scient. Hung 4: 257-306.
- Sporne, K.R. (1956). The phylogenetic classification of the angiosperms. Biol. Rev. 31: 1-29.
- Stant, M.Y. (1952). Anatomical evidence for including Nyctanthes and Dimetra in the Verbenaceae. Kew Bull. 1952: 273-276.
- Subramanian, S.S., Nagarajan, S. and N. Sulochana. (1971). Flavonoids of Millingtonia hortensis. Curr. Sci. 40: 194.
- Subramanian, S.S. and Nair, A.G.R. (1972). Flavonoids of

- the leaves of Oroxylum indicum and Pajanelia longifolia. Phytochemistry, 11(1): 439-440.
- Subramanian, S.S., Nagarajan, S. and N. Sulochana. (1972). Flavonoids of Bignoniaceous plants. Phytochemistry, 11(4): 1499.
- Subramanian, S.S. and Nair, A.G.R. (1972). Scutellarein-4'-arabinoside from the leaves of Clerodendron nereifolium. J. Indian Chem. Soc. 49(10): 1061-1062.
- Subramanian, S.S. and Nair, A.G.R. (1972). Scutellarein and pectolinarigenin from the leaves of Clerodendron phlomidis and Duranta repens. Phytochemistry, 11: 3095-3096.
- Subramanian, S.S., Nagarajan, S. and N. Sulochana. (1972). Chrysin-7-rutinoside from the leaves of Dolichandrone falcata. Phytochemistry, 11(1): 438-439.
- Subramanian, S.S., Nagarajan, S. and N. Sulochana. (1973). Hydroquinone from the leaves of Jacaranda mimosaeifolia. Phytochemistry, 12(1): 220-221.
- Subramanian, S.S. and Ramesh, P. (1973). Rutin from the leaves of Bignonia magnifica. Indian J. Pharm. 35(6): 207.
- Subramanian, S.S. and Nair, A.G.R. (1973). Scutellarein and Hispidulin 7-o glucuronide from the leaves of Clerodendrum indicum and Clerodendrum infortunatum. Phytochemistry, 12: 1195.
- Subramanian, S.S., Nair, A.G.R. and T.N.C. Vedantham. (1973). Chemical examination of the leaves of Clerodendrum inerme. Indian J. Pharm. 35(6): 191-192.
- Subramanian, S.S., Nair, A.G.R. and T.N.C. Vedantham. (1974).

- Chemical examination of the leaves of Stachytarpheta indica. Indian J. Pharm. 36(1): 15-16.
- Subramanian, S.S. and Vedantham, T.N.C. (1974). Chemical components of Avicennia officinalis. Indian J. Pharm. 36(4): 105-106.
- Subramanian, M., Koziparambil, K.P., Ayyappath, S. and J.D. Connolly. (1976). A new flavone 6-glucoside from Citharexylum suberratum. Phytochemistry, 15: 838.
- Subramanian, P.M. and Misra, G.S. (1978). Leucoanthocyanidins of Vitex negundo. Indian J. Chem. 16B(7): 615-616.
- Sugiura, T. (1931). A list of chromosome numbers in angiospermous plants. Bot. Mag. (Tokyo) 45 : 353-355.
- Sugiura, T. (1936). Studies on the chromosome numbers in higher plants, with special reference to cytokinesis. Cytologia 7: 544-595.
- Swain, T. (ed.) (1963). "Chemical Plant Taxonomy". Academic Press. London.
- Swain, T. (ed.) (1966). "Comparative phytochemistry". Academic Press, London and New York.
- Swain, T. (1975). Evolution of flavonoid compounds. In: J.B. Harborne, T.J. Mabry and H. Mabry (eds.). "The Flavonoid", pp. 1096-1129. Chapman and Hall, London.
- Takhtajan, A. (1969). "Flowering plants, origin and dispersal". Edinburgh.
- Takhtajan, A. (1980). Outlines of the classification of flowering plants (Magnolio phyta). Botanical Review. 46(3): 225-358.

- Taneja, S.C., Bhatnagar, R.P. and H.P. Tiwari. (1975).
Chemical constituents of flowers of Tecomella undulata.
Indian J. Chem. 13(4): 427-428.
- Tantisewie, B. and Ottosticher. (1975). Isolation of
ipolamide from Stachytarpheta indica. Phytochemistry
14(5/6): 1462-1463.
- * Thunberg, (1784). Flora Japonica. 252.
- Tillequein, F., Melle-Ehenri and R.R. Paris. (1977). Isolation
of 5-hydroxy-6,7,4'-tri methoxy flavone from leaves of
Phyllarthron madagascariense. Planta Med. 31(1): 76-79.
- Tomaszewski, M. (1960). Bull. Acad. Polon. Sci. 8: 61.
(Cited from Biochemistry of phenolic compounds.
J.B. Harborne (1964) pp. 80. Academic Press, London and
New York).
- Trim, A.R. and Hill, R. (1951). Biochem. J. 50: 310.
- Venkatasubban, K.R. (1944). Cytological studies in Bignoniaceae.
Annamalai University, Annamalainagar. (Cited from
Goldblatt and Gentry, 1979).
- Venkatasubban, K.R. (1945). "Cytological studies in
Bignoniaceae". IV. Proc. Ind. Acad. Sci., 21B, 77-92.
- Vedantham, T.N.C. and S.S. Subramanian. (1976). Nonflavonoid
components of Vitex trifolia. Indian J. Pharm. 38(1):13.
- Wagner, H. (1966). Flavonoid c-glycosides. In: "Comparative
phytochemistry" (ed. T. Swain). Academic Press, London.
- * Wasicky, R., Akisue, M.K. and T. Saito. (1967). The phyto-
chemistry of species of the genus Tabebuia L. Analysis
of some compounds. Rev. Fac. Farm. Bioquim. Sao Paulo
5(2): 383-395.

- Weifferring, J.H. (1966). Phytochemistry 5: 1053.
- Weimark, G. (1972). On "Numerical Chemotaxonomy". Taxon 21: 615-619.
- Wernham, H.F. (1911-12). Floral evolution; with particular reference to the sympetalous dicotyledons. New Phyl. 10: 73-83, 109-120, 145-159, 217-226, 293-305, (1911). 11: 145-166, 217-235, 290-305, 373-397 (1912). (Cited from R.D. Gibbs (1974). Vol. IV.).
- Westfall, J.J. (1949). Cytological and embryological evidences for the reclassification of Paulownia. Amer. J. Bot. 36: 805.
- Wettstein, R. (1935). "Handbuch der systematischen Botanik" (4th ed.) Leipzig and Viena.
- Yousef, F., Wahbakhalil, S.K. and S.P. Pappas. (1973). Separation and characterisation of a new alkaloid from the fruit of Duranta repens. Planta-Med. 23(2): 173-175.
- * Zirvi, K.A. and Amir, F. (1973). Chemical investigation of Jacaranda acutifolia. Pak. J. Sci. Ind. Res. 16(5): 178.

* * * * *

* Not referred to the original.