

List of Tables

2.1	“m scheme” for the configuration $ (7/2)^2J)^*$	32
3.1	Nuclear charge radii for Fermi and homogeneous mass distribution	43
5.1	Add-back ratio for Clover-3 used in the IUAC experiment	62
5.2	Energies of the Doppler shifted γ -rays (E_γ in keV) for ^{58}Ni and ^{112}Sn excitation as a function of relative angle $\phi_{12} = \phi_\gamma - \phi_p $ between γ -ray and particle detection. The detection of nickel and tin particles in the PPAC are calculated separately.	63
5.3	Gamma-ray detection angles ($\vartheta_\gamma, \varphi_\gamma$) for each Ge-crystal extracted from the Doppler shift data of the Ni excitation (effective beam energy $E_{lab}=167\text{MeV}$).	71
5.4	Peak areas and intensity ratios for Sn and Ni excitation (exc) in $^{112}\text{Sn}+^{58}\text{Ni}$ and $^{116}\text{Sn}+^{58}\text{Ni}$ systems using the DDL analysis.	73
5.5	Peak areas and intensity ratios for Sn and Ni excitation (exc) in $^{112}\text{Sn}+^{58}\text{Ni}$ and $^{116}\text{Sn}+^{58}\text{Ni}$ systems using the SDL analysis.	74
5.6	φ -dependence of particle- γ intensities (Clover-2)	74
5.7	Cross sections for the $^{58}\text{Ni}+^{116}\text{Sn}$ system at 175MeV for two different angular ranges.	77
5.8	Cross sections for the $^{58}\text{Ni}+^{112}\text{Sn}$ system at 175MeV for two different angular ranges.	77
5.9	Double ratio of $^{112}\text{Sn}/^{116}\text{Sn}$ as determined from tab. 5.7 and tab. 5.8	77
5.10	Measured reduced transition probabilities for ^{112}Sn and ^{114}Sn	78
6.1	$I^\pi=2^+$ energies and E2 strengths in $^{102-130}\text{Sn}$. The experimental data of the neutron-deficient isotopes are averaged values of Refs. [6, 14, 15, 16].	84
6.2	Relativistic mean field parameterizations NL1, NL-SH and NL3. K_{nm} corresponds to the incompressibility of the matter for each set of parameters. The mass of the ρ meson is fixed to the experiment value, $m_\rho=763.0$ MeV.	87
6.3	$I^\pi=2^+$ energies and E2 strengths in $^{102-130}\text{Sn}$. The experimental data of the neutron-deficient isotopes are averaged values of Refs. [6, 14, 15, 16]	89
7.1	The measured energy resolutions with ORTEC 572 and CAEN N568BC	100

