

LIST OF FIGURES		
1.1	Diagrammatic representation of various antibiotic targets in bacterial cell.	04
1.2	Diagrammatic representation of popular mechanisms of antibiotic resistance in bacteria	11
1.3	Schematic diagram of representative drug exporting systems in Gram-positive bacteria, highlighting different families of pumps involved in resistance.	14
1.4	Generalized pathway of Fatty acid biosynthesis and Polyketide biosynthesis, suggesting flexibility at various levels in formation of polyketide molecules.	23
1.5	Schematic presentation of biosynthetic pathways of some polyketides	31
1.6	Chemical structures of members of the aureolic acid family	33
1.7	Novel antitumor compounds generated by combinatorial biosynthesis.	17
3.1	Zones of inhibition of <i>M. luteus</i> in agar well assay, at increasing (20ng, 100ng, 200ng, 300ng, 500ng) concentrations of the polyketide.	96
3.2	Nucleic Acid binding study of the polyketide (Test) in relation to known DNA binding molecule, Chromomycin (Control)	108
3.3	Agarose gel electrophoresis of DNA incubated with or without polyketide	109
4.1	Agar well assay for Chromomycin using a co-inoculum of wildtype and Chromo <sup>r</sup> showing two distinct zones of inhibitions marked by difference in carotenoid levels of the two.	125

4.2	Difference in inhibition zones of Chromo <sup>r</sup> vs wild type <i>M. luteus</i> against Chromomycin and Daunomycin	130
4.3	Restriction digestion pattern using different REs.	138
4.4	Comparison of SDS-PAGE of total proteins prepared from colonies and broth.	140
4.5	Effect of increasing chromomycin concentration on Chromo <sup>r</sup> against wild type.	140
5.1	Growth of <i>S. griseus</i> (left) and <i>S. flaviscleroticus</i> (right) on different media.	158
5.2	Comparative TLC of metabolite extracts of <i>S. flaviscleroticus</i> (left) and <i>S. griseus</i> (right) from various media, observed under ultraviolet light (365nm)	162
5.3	Relative spectrophotometric signatures of <i>S. flaviscleroticus</i> (SF) and <i>S. griseus</i> (SG) extracts from different media	163
5.4	Restriction Enzyme Digestion Pattern of Genomic DNA from <i>S. flaviscleroticus</i> (left) and <i>S. griseus</i> (Right) using different enzymes.	165
5.5	16S rDNA phylogenetic tree of <i>S. flaviscleroticus</i>	169
5.6	Phylogenetic tree of ketosynthase gene of <i>S. flaviscleroticus</i>	171
5.7	CLUSTLW phylogram of selected 16s rDNA sequences	173
6.1	Diagrammatic representation of the conjugation mediated transposition process furnished by the presence of F plasmid containing Tn1000	184
6.2	Physical map of Tn1000.	185
6.3	Diagrammatic representation of <i>Eco</i> R1 sites in physical map of PKS DNA	186

6.4	The Identical Cluster organization of two chromomycin producers compared to a much different cluster of structurally related Mithramycin	189
6.5	Putative chromomycin biosynthesis pathway showing role of various proteins in formation of molecule.	201
6.6	Minimal PKS region of different type II polyketide clusters	203
6.7	Structure and position of different sugars in structure of Chromomycin A <sub>3</sub>	205
6.8	Putative biosynthesis pathway of different sugars of chromomycin A <sub>3</sub> as proposed for <i>S. griseus</i> cluster.	206
7a.1	Schematic presentation of 17kb EcoR1 fragment, containing resistance genes, with restriction sites of <i>Kpn</i> 1, <i>Xho</i> 1 and BglII represented on nucleotide scale.	224
7a.2	Relative zone of inhibition against Crude ethyl acetate extract from <i>S. flaviscleroticus</i> (K20ul, K30ul) and pure chromomycin (C10, C20)	227
7a.3	Phenotypic characterization by expressing <i>sfr</i> B, <i>sfr</i> A and <i>sfr</i> X in heterologous host	227
7a.4	Confirmation of various clones by checking restriction digestion pattern.	228
7b.1	Clone construction and confirmation of <i>actRp</i> -EGFP transcriptional fusion system.	235
7b.2	Diagrammatic representation of the proposed strategy.	237
7b.3	Construction and confirmation of pTRACT clone	239
7b.4	Conserved island of promoter region in a totally non-conserved intergenic DNA upstream of activator gene in <i>S. flaviscleroticus</i> and <i>S. griseus</i>	241

7b.5	Thiostrepton mediated <i>tipAp</i> induction in <i>S. lividans</i> transformed with pIJ8655 resulting in concentration dependend increase in fluorecence of EGFP.	245
7b.6	Fluorescence detection in pTRACT transformed <i>S. lividans</i> after 2 days and 4 days of incubation. An overall increase in fluorescent mycelia was observed with time but the intensity remained constant.	247
7b.7	Pattern generated using MEME software, for the bidirectional promoter complexes associated with the both regulatory genes of chromomycin synthase clusters of the two producers.	250a