BIBLIOGRAPHY

.

1. Abdulbari et al (2006)

Prevalence of under weight, over weight and obesity in Qatar adolescents. Food & Nutrition Bulletin; 27(1): 39-45.

2. Aboshkair et al (2011)

Assessment of Weight Status of Secondary School Children in Selangor, Malaysia. Australian Journal of Basic and Applied Sciences; 5(9): 1675-1682.

3. Abrantes et al (2002)

Recommendations for the use of body mass index for the classification of over weight and obese children and adolescents. Food and Nutrition Bulletin; 23 (3): 262-273.

- Adolphus et al (2013)
 The effects of breakfast on behavior and academic performance in children and adolescents. Frontiers in Human Neuroscience; 7(425): 1-28.
- 5. Agarwal and Agarwal (2003)

The growth infancy to adolescence. CBS Publishers and Distributors- New Delhi.

- Aggarwal et al (2008)
 Prevalence of obesity and overweight in affluent adolescents from Ludhiana,
 Punjab. Indian Pediatr; 45: 500-502.
- 7. Akinpelu et al (2008)

Overweight and Obesity: Does It Occur In Nigerian Adolescents in an Urban Community? Int J of Biomedical and Helath Sciences; 4(1): 11-17.

8. Alaimo et al (2001)

Food insufficiency and American school-aged children's cognitive, academic and psychosocial developments. Pediatrics; 108(1):44-53.

9. Ali et al (2009)

Smokeless tobacco use among adult patients who visited family practice clinics in Karachi, Pakistan. *J Oral Pathol Med*; 38(5): 416- 421.

10. Allensworth et al (1997)

Institute of Medicine. Schools and health: our nation's investment. Washington, DC: The National Academies Press.

Bibliography

11. American Dietetic Association(2003) Society for Nutrition Education; American School Food Service Association (2003) Position of the American Dietetic Association, Society for Nutrition Education, and American School Food Service Association. Nutrition services: an essential component of comprehensive school health programs. J Nutr Educ Behav; 35:57-67. 12. Anderson et al (2005) The impact of a school-based nutrition education intervention on dietary intake and cognitive and attitudinal variables relating to fruits and vegetables. Public Health Nutrition; 8(6):650-656. 13. Anderson et al (2005) The impact of a school-based nutrition education intervention on dietary intake and cognitive and attitudinal variables relating to fruits and vegetables. Public Health Nutrition; 8(6):650-656. 14. Angelopoulos et al (2009) Changes in BMI and blood pressure after a school based intervention: The CHILDREN study. European Journal of Public Health; 19(3): 319-325. 15. Armstrong et al (2006) Obesity and overweight in South African primary school children - the Health of the Nation Study. S Afr Med J; 96: 439-444. 16. Austin et al (2006) Facilitating Change in School Health: A Qualitative Study of Schools' Experiences Using the School Health Index. Prev Chronic Dis; 3(2): A35 -40.Available at: http://whqlibdoc.who.int/hq/1998/WHO NUT NCD 98.1 (p1-158).pdf 17. Aziz et al (2009) Prevalence of OW& OB among children and adolescents of affluent schools in Karanchi. J Pak Med Assoc JPMA; 59:35. 18. Balvin Frantzen et al (2013) Association between frequency of ready-to-eat cereal consumption, nutrient intakes, and body mass index in fourth- to sixth-grade low-income minority

225

children. J. Acad. Nutr. Diet; 113: 511-519.

19. Baranowski et al (2000)
Gimme 5 Fruit, Juice, and Vegetables for Fun and Health: Outcome Evaluation. Health Education & Behavior; 27(1): 96-111.

- Bartholomew and Jowers (2006)
 Increasing frequency of lower-fat entrees offered at school lunch: an environmental change strategy to increase healthful selections. Journal of the American Dietetic Association; 106(2):248–252.
- 21. Bauer et al (2006)

Swimming upstream: faculty and staff members from urban middle schools in low-income communities describe their experience implementing nutrition and physical activity initiatives. Prev Chronic Dis; 3:A37–45.

- Baya et al (2010)
 International BMI-for-age references underestimate thinness and overestimate overweight and obesity in Bolivian adolescents. Nutr Hosp; 25:428-436.
- 23. Bayne-Smith et al (2004)Improvements in heart health behaviors and reduction in coronary artery disease risk factors in urban teenaged girls through a school-based intervention: the PATH program. American Journal of Public Health; 94 (9): 1538-1543.
- 24. Bayne-Smith M et al (2004)

Improvements in heart health behaviors and reduction in coronary artery disease risk factors in urban teenaged girls through a school-based intervention: the PATH program. American Journal of Public Health; 94(9):1538–1543.

- 25. Beech et al (2003)
 Child- and parent-targeted interventions: the Memphis GEMS pilot study.
 Ethn Dis; 13(1):S40-53.
- Bellizi and Dietz (1999)
 Workshop on childhood obesity: summary of the discussion. American J of Clinical Nutrition; 70: 173S-175S.
- 27. Bharati, Deshmukh and Garg (2008)
 Correlates of overweight & obesity among school going children of Wardha city, Central India Indian J Med Res; 127:539-543.
- 28. Bhardwaj et al (2008)

Childhood obesity in Asian Indians: a burgeoning cause of insulin resistance, diabetes and sub-clinical inflammation. Asia Pac J Clin Nutr; 17(S1):172-175. 29. Bhardwaj et al (2008) Childhood obesity in Asian Indians: a burgeoning cause of insulin resistance, diabetes and sub-clinical inflammation. Asia Pac J Clin Nutr; 17(S1):172-175. 30. Bhave ei al (2004) IAP National Task Force for Childhood Prevention of Adult Diseases.Available at: http://medind.nic.in/ibv/t04/i6/ibvt04i6p559.pdf 31. Bhave et al (2004) IAP National task force for childhood prevention of adult disease: Childhood obesity. Indian Paediatr; 41: 559-75. 32. Bourne et al (2002) Where does the black population of South Africa stand on the nutrition transition? Public Health Nutr; 5(1A):157-162. 33. Bovet et al (2011) Prevalence of thinness in children and adolescents in the Seychelles: comparison of two international growth references. Nutrition Journal; 10:65. 34. Bray (1992) Pathophysiology of obesity. Am J Clin Nutr; 55(2): 488S-494S. 35. Briefel et al (2009) School food environments and practices affect dietary behaviors of U.S. public school children. J Am Diet Assoc; 109(2):S91-107. 36. Bruner et al (1996) Randomised study of cognitive effects of iron supplementation in non anaemic and iron deficient adolescent girls. The Lancet; 348 (9033): 992-996. 37. Butte and Garza (2006) Development of an international growth standard for preadolescent and adolescent children. Food Nutr Bull; 27:S169-S326. 38. Butte et al (2007) Evaluation of the feasibility of international growth standards for school-aged children and adolescents. J Nutr; 137:153-57. 39. Caballero et al (2003)

Pathways: a school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren1-3. Am J Clin Nutr; 78:1030-8. 40. Caballero et al (2003) Pathways: a school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren. American Journal of Clinical Nutrition; 78(5):1030-1038. 41. Cater and Carr (2004) Competencies, knowledge, and skills of effective school nutrition managers. R-66-03. University, MS: National Food Service Management Institute; 2004. 42. CD (2007) Can eating fruits and vegetables help people manage their weight? Available at: http://www.cdc.gov/nccdphp/dnpa/nutrition/pdf/rtp_practitioner_10_07.pdf 43. CDC (2005) A school health index (middle / high school), a self assessment and planning guide. Available at: http://www.cdc.gov/healthyyouth/shi/pdf/MiddleHigh.pdf 44. CDC (2008) Competitive foods and beverages available for purchase in secondary schools-selected sites, United States, 2006. MMWR ;57:935-38. 45. CDC (2012) Health Coordinated School Program. Available at: http://www.cdc.gov/healthyyouth/cshp/index.htm 46. CDC (2012) PARENT ENGAGEMENT: Strategies for Involving Parents in School Health. Available at: www.cdc.gov/HealthyYouth/ 47. Center for Disease Control (2005) A school health index (middle / high school), a self assessment and planning guide. Available at: http://www.cdc.gov/healthyyouth/shi/pdf/MiddleHigh.pdf 48. Center for Disease Control (2011) School health guidelines to promote healthy eating and physical activity. Recommendations and reports, vol. 60, No.5. 49. Chandrashekhar et al (2013) Blood pressure and its associated factors among primary school children in sub urban in Selangor, Malaysia: A cross sectional survey. Journal of Family and Community Medicine; 20(2): 90-97.

50. Chatterjee P (2002)

India sees parallel rise in malnutrition and obesity. Lancet; 360: 1948.

51. Chhatwal et al (2004)

Obesity among preadolescent and adolescents of a developing country (India). Asia Pacific J Clin Nutr; 13(3): 231-235.

52. Cho et al (2003)

The effect of breakfast type on total daily energy intake and body mass index: results from the Third National Health and Nutrition Examination Survey (NHANES III). J Am Coll Nutr; 22: 296–302.

53. Choudhary and Dhage (2008)

A study of anemia among females in the urban area of Nagpur. Indian journal of community medicine; 33(4): 243-245.

54. Coly et al (2006)

Preschool stunting, adolescent migration, catch-up growth, and adult height in young senegalese men and women of rural origin. J Nutr 2006; 136:2412-2420.

55. Condon, Crepinsek and Fox (2009)

School meals: types of foods offered to and consumed by children at lunch and breakfast. J Am Diet Assoc; 109(2):S67–78.

56. Davis and Pamela (2005)

The Influence of Parent Education and Family Income on Child Achievement: The Indirect Role of Parental Expectations and the Home Environment. Journal of Family Psychology; 19(2): 294-304

57. Davis et al (2003)

Pathways curriculum and family interventions to promote healthful eating and physical activity in American Indian schoolchildren. Preventive Medicine; 37:S24–S34.

58. Davis et al (2005)

The relationship between qualified personnel and self-reported implementation of recommended physical education practices and programs in U.S. schools. Res Q Exerc Sport; 76:202–11.

59. Davis et al (2005)

The relationship between qualified personnel and self-reported implementation of recommended physical education practices and programs in U.S. schools. Res Q Exerc Sport; 76:202–11.

60. De Benoist et al (2004)

Iodine status worldwide. WHO Global Database on Iodine Deficiency. Geneva: World Health Organization, 2004.

 $61. \qquad \text{de la Hunty et al (2013)}$

Does regular breakfast cereal consumption help children and adolescents stay slimmer? A systematic review and meta-analysis. Obes. Facts; 6: 70–85.

62. de Onis et al (2007)

Development of a WHO growth reference for school-aged children and adolescents. Bulletin of the World Health Organization; 85 (9): 660-667.

63. De Onis M (2004)

The use of anthropometry in the prevention of childhood overweight and obesity. Int J Obes Relat Metab Disord; 28:S81-S85.

64. Deshmukh-Taskar et al (2010)

The relationship of breakfast skipping and type of breakfast consumption with nutrient intake and weight status in children and adolescents: the National Health and Nutrition Examination Survey 1999-2006. J. Am. Diet. Assoc; 110(6): 869–878.

65. Dewey (1999)

Reviewing the relationship between school factors and substance use for elementary, middle, and high school students. J Prim Prev; 19(3):177–225.

66. Dubois et al (2007)

Regular sugar-sweetened beverage consumption between meals increases risk of overweight among preschool-aged children. J Am Diet Assoc; 107(6):924-34.

67. Dunkle and Nash (1991)

Beyond the health room. Washington, DC: Council of Chief State School Officers, Resource Center on Educational Equity. Available at http://www.eric.ed.gov/PDFS/ED340681.pdf.

68. Durrani and Waseem (2011)

Blood pressure distribution and its relation to anthropometric measurements among school children in Aligarh. Indian journal of public health; 55 (2): 121-124.

69. Eggal and Lopriore (2006)

Agriculture/Health collaboration: The Key to fighting malnutrition in all its forms. In Diet related chronic diseases and the double Burden of malnutrition in West Africa. SCN News No. 33.

70. Ella et al (2011)

Prevalence of over weight and obesity, and status of chronic noncommunicable diseases and some related risk factors among Egyptian adolescents. Journal of Diabetes and Endocrinology; 2(4): 41-52.

71. Elmadfa and Weichselbaum (2004)

European Nutrition and Health Report. Vol 58, Basel, Karger Publishing, 2005.

72. Engels et al (2005)

Promoting healthful diets and exercise: efficacy of a 12-week after-school program in urban African Americans. Journal of the American Dietetic Association; 105(3):455–459.

73. Epstein (2001)
 School, family, and community partnerships: preparing educators and improving schools. Boulder, CO: Westview Press; 2001.

74. Epstein et al (2009)

School, family, and community partnerships: your handbook for action. 3rd ed. Thousand Oaks, CA: Corwin Press; 2009.

75. Epstein JL (2011)

School, Family, and Community Partnerships: Preparing Educators and Improving Schools Second Edition. Boulder, CO: Westview Press.

Fairclough and Stratton (2005)
 Improving health-enhancing physical activity in girls' physical education.
 Health Educ Res; 20:448–57.

77. FAO (2007)

Report of the technical meeting of the Asia Pacific network for foo'd and nutrition on school based nutrition. RAP publication, 14, Bangkok, Thailand.

78. Ferna'ndez et al (2004)

Waist circumference percentiles in nationally representative samples of African-American, European-American, and Mexican-American children and adolescents. J Pediatr; 145(4): 439–444.

79. Field (2006)

Epidemiology of the health and economic consequences of pediatric obesity. In *Handbook of pediatric obesity Etiology, Pathophysiology and Prevention* Edited by: Goran MI, Sothern MS. Boca Raton: Taylor & Francis Group, LLC.

80. Forshee and Storey (2003)

Total beverage consumption and beverage choices among children and adolescents. Int J Food Sci Nutr; 54(4):297-307.

81. Foster et al (2008)

A Policy-Based School Intervention to Prevent Overweight and Obesity. *Pediatrics;* 121: e794- e802.

82. Foster et al (2008)

A Policy-Based School Intervention to Prevent Overweight and Obesity. *Pediatrics;* 121: e794- e802.

83. French et al (2004)

An environmental intervention to promote lower-fat food choices in secondary schools: outcomes of the TACOS Study. American Journal of Public Health; 94(9):1507–1512.

84. Fulkerson et al (2002)

Foodservice staff perceptions of their influence on student food choices. J Am Diet Assoc; 102:97–9.

85. Gales-Camus (2006)

Nutrition: an input and foundation for Health Diet. Tackling the double burden of Malnutrition. A global Agenda. SCN News. No. 32.

86. Gargari et al (2004)

Prevalence of overweight and obesity among high-school girls in Tabriz, Iran, in 2001: Food Nutr Bull; 25(3):288-91.

87. Gibson S (2003)

Micronutrient intakes, micronutrient status and lipid profiles among young people consuming different amounts of breakfast cereals: further analysis of data from the National Diet and Nutrition Survey of Young People aged 4 to 18 years. Public Health Nutr; 6: 815–820.

88. Gills and Gills (2005)

Nutrient inadequacy in obese and non-obese youth. Can J Diet Pract Res; 66: 237-242.

89. Gittelsohn et al (2003)

Pathways: lessons learned and future directions for school-based interventions among American Indians. Preventive Medicine; 37(1):S107–S112.

90. Gleason and Dodd (2009)
 School breakfast program but not school lunch program participation is associated with lower body mass index. J Am Diet Assoc; 109(2):S118–28.

91. Going et al (2003)

The effects of the Pathways Obesity Prevention Program on physical activity in American Indian children. Preventive Medicine; 37(1):S62–S69.

92. Golan and Crow (2004)

Targeting parents exclusively in the treatment of childhood obesity: long-term results. Obes Res; 12 (2): 357-361.

93. Gomes et al (2009)

Influence of different body mass index cut-off values in assessing the nutritional status of adolescents in a household survey. Cad Saude Publica; 25:1850-1857.

- 94. Gonzalez, Jones and Frongillo (2009)
 Restricting snacks in U.S. elementary schools is associated with higher frequency of fruit and vegetable consumption. J Nutr; 139:142–4.
- 95. Gopalan, Sastri Rama and Balasubramanian (2004)
 Nutritive value of Indian foods. National Institute of Nutrition, Indian Council of Medical Research, Hyderabad.

96. Gordon et al (2009)
The third school nutrition dietary assessment study: summary and implications. J Am Diet Assoc; 109(1):S129–135.

97. Gorely et al (2009)

Effect of a school-based intervention to promote healthy lifestyles in 7–11 year old children. *International Journal of Behavioral Nutrition and Physical Activity*; 6:5-16.

98. Gorely et al (2009)

Effect of a school-based intervention to promote healthy lifestyles in 7–11 year old children. International Journal of Behavioral Nutrition and Physical Activity; 6:5.

99. Goyal et al (2010)

Prevalence of Overweight and Obesity in Indian Adolescent School Going Children : Its Relationship with Socioeconomic Status and Associated Lifestyle Factors. JAPI; 58: 151-158.

100. Goyal et al (2010)

Prevalence of Overweight and Obesity in Indian Adolescent School Going Children : Its Relationship with Socioeconomic Status and Associated Lifestyle Factors. JAPI; 58: 151-158.

101. GSEB (2008)

Gujarat State School Textbook Board - Education Department (GSEB). Available at:http://gujarat-education.gov.in/textbook/

Halterman et al (2001)
 Iron deficiency and cognitive achievement among school-aged children and adolescents in the United States. Pediatrics; 107: 1381-1386.

- 103. Hastings et al (2006)The Extent, Nature and Effects of Food Promotion to Children: A Review of the Evidence. Available at:
- Hedley et al (2002)
 Overweight and obesity among US children, adolescents, and adults, 1999-2002. JAMA; 291:2847-50.

105. Henderson and Mara (2002)

A new wave of evidence: the impact of school, family, and community connections on student achievement. Austin, TX: Southwest Educational Development Laboratory; 2002.

106. Hoelscher et al (2003)

How the CATCH eat smart program helps implement the USDA regulations in school cafeterias. Health Education & Behavior; 30(4):434–446.

- 107. Hoelscher et al (2004)
 School-based health education programs can be maintained over time: results from CATCH Institutionalization study. Preventive Medicine; 38(5):594–606.
 108. Hong et al (2007)
- Gender and SES differences in BMI of secondary high school students in Ho Chi Minh City. Asia Pac J Clin Nutr; 16 (1):74-83.

109. Hooshyar et al (2001)

Using body mass index for determination of obesity, overweight and underweight in Iranian groups. In: Iranian 6th Nutritional Congress Executive Committee, 1st ed. Ahwaz, Islamic Republic of Iran, Mehr Publications.

110. Hoyland et al (2009)

A systematic review of the effect of breakfast on the cognitive performance of children and adolescents. Nutr Res Rev; 22(2):220–243. Available at: http://www.who.intdietphysicalactivitypublicationsHastings_paper_marketing. pdf

111. Hunt (2001)

Investing in children: Child protection and economic growth. In: Ortiz I (ed). *Social protection in Asia and the Pacific.* Manila, Philippines:Asian Development Bank; 2001:579–98.

112. IDF (2007)

The International Diabetes Federation consensus definition of the metabolic syndrome in children and adolescents. IDF Communications.Available at: http://www.idf.org/webdata/docs/Mets_definition_children.pdf

113. Iyer et al (2006)

Obeisty in school children. Magnitude and risk factor analysis. ICMR funded Project report, Department of Foods and Nutrition. WHO collaborating centre for diet related non-communicable diseases and anemia control, MS University of Baroda, India. 2006.

114. Iyer et al (2010)

An exploratory study on the nutritional status and Determinants of malnutrition of urban and rural adolescent children (12-16) years of vadodara city. International Journal of Applied Biology and Pharmaceutical Technology; 2(1): 102-107.

115. Iyer et al (2010)

An exploratory study on the nutritional status and Determinants of malnutrition of urban and rural adolescent children (12-16) years of vadodara city. International Journal of Applied Biology and Pharmaceutical Technology; 2(1): 102-107.

116. Iyer et al (2011)

An exploratory study on the nutritional status and determinants of malnutrition of urban and rural adolescent children (12-16) years of vadodara city. International Journal of Applied Biology and Pharmaceutical Technology; 2(1): 102-107.

117. Iyer et al (2011)

An exploratory study on the nutritional status and Determinants of malnutrition of urban and rural adolescent children (12-16) years of vadodara city. International Journal of Applied Biology and Pharmaceutical Technology; 2(1): 102-107.

118. Iyer et al (2011)

An exploratory study on the nutritional status and Determinants of malnutrition of urban and rural adolescent children (12-16) years of vadodara city. International Journal of Applied Biology and Pharmaceutical Technology; 2(1): 102-107.

119. Iyer et al (2011)

Magnitude and Determinants of Overweight and Obesity in 6-12 year old school children of Vadodara City. *Curr Pediatr Res; 15 (2): 105-109.*

120. Jaisheeba et al (2012)

Influence of Westernized culture and changed dietary habits on the BMI status of the school children of Tirunelveli. Int.J.PharmTech Res; 4(3): 1065-1077.

121. James (2004)

Assessment of obesity for Asians. Available at: www.obesity.org

122. Jazayeri (2005)

Overweight and Obesity among School-Aged Children of Metropolitan Tehran, Iran Pakistan Journal of Nutrition; 4 (5): 342-344.

123. Jeemon et al (2009)

- 124. Double burden of underweight and overweight among children (10–19 years of age) of employees working in Indian industrial units. The National Medical Journal of India; 22(4): 172-176.
- 125. Jeemon et al (2009)
 Double burden of underweight and overweight among children (10–19 years of age) of employees working in Indian industrial units. The national medical journal of India; 22(4): 172-176.

126. Jeemon et al (2009)

Double burden of underweight and overweight among children (10–19 years of age) of employees working in Indian industrial units. The national medical journal of India; 22(4): 172-176.

127. Jimenez et al (2003)

Comparison of the dietary intakes of two different groups of children (grades 4 to 6) before and after the Kahnawake Schools Diabetes Prevention Project. Journal of the American Dietetic Association; 103(9):1191–1194.

128. JNC VII Report.

The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation and Treatments of High Blood Pressure. JAMA 19: 2560-2572.

- Juliusson et al (2010)
 Overweight and obesity in Norwegian children: prevalence and sociodemographic risk factors. Acta Pædiatrica; 0803–5253.
- 130. Kain et al (2004)
 School-based obesity prevention in children; methodology and evaluation of a controlled study. International Journal of Obesity; 28(4):483–493.
- 131. Kanani and Poojara (2006)
 Supplementation with IFA enhances growth in adolescent Indian girls. J.
 Nutrition; 130: 452S-455S.
- 132. Kaneria et al (2006)
 Prevalence of Overweight and Obesity in Relation to Socio-economic Conditions in Two Different Groups of School-age Children of Udaipur City (Rajasthan). JIACM; 7(2): 133-135.

133. Kanien'keha (2005)

Children 6 to 11 years old: 8 year results from the Kahnawake Schools Diabetes Prevention Project. Pediatrics; 115(2):333-339.

134. Kapil et al (2002)
Prevalance of obesity amongst affluent adolescent school children in Delhi. *Indian Paediatr; 39*: 449-52.

135. Kapil et al (2002)Prevalence of obesity amongst affluent adolescent school children in Delhi.Indian Pediatrics; 39: 449-452.

- 136. Kapil et al (2002)
 Prevalence of obesity amongst affluent adolescent school children in Delhi. Indian Pediatr; 39:449–52.
- 137. Kaur et al (2008)

Prevalence of overweight and obesity amongst school children in Delhi, India Asia Pac J Clin Nutr; 17 (4):592-596.

138. Kaur et al (2013)

Association of Obesity with Hypertension Amongst School-Age Children Belonging to Lower Income Group and Middle Income Group in National Capital Territory of Delhi. Indian Journal of Community Medicine; 38(3): 175-179

139. Kaur et al (2013)

Association of Obesity with Hypertension Amongst School-Age Children Belonging to Lower Income Group and Middle Income Group in National Capital Territory of Delhi. Indian Journal of Community Medicine; 38(3): 175-179.

140. Kaur et al (2013)

Association of Obesity with Hypertension Amongst School-Age Children Belonging to Lower Income Group and Middle Income Group in National Capital Territory of Delhi. Indian Journal of Community Medicine; 38(3): 175-179.

141. Kelder et al (2003)

Long-term implementation of the CATCH physical education program. Health Educ Behav; 30:463–75.

142. Kelder et al (2003)

Long-term implementation of the CATCH physical education program. Health Educ Behav; 30:463–75.

 143. Kelishadi et al (2007)
 Association of anthropometric indices with cardiovascular disease risk factors among children and adolescents: CASPIAN Study. Int J Cardiol; 117:340-48.

144. Keski-Rahkonen et al (2003)
Breakfast skipping and health-compromising behaviors in adolescents and adults. Eur J Clin Nutr; 57: 842–853.

- 145. Khadilkar and Khadilkar (2004)
 Prevalence of obesity in affluent school boys in Pune. Indian Pediatr; 41: 857-858.
- 146. Khadilkar et al (2010)Overweight and obesity prevalence and body mass index trends in Indian children. Int J Pediatr Obes; 6(2): e216-e224.

147. Khadilkar VV, Khadilkar AV (2004) Prevalence of obesity in affluent schoolboys in Pune. Indian Pediatr; 41: 857-858.

148. Khan (2010)

A study of the risk Factors and the prevalence of hypertension in the adolescent school boys Of ahmedabad city. Journal of Clinical and Diagnostic Research; 4: 3348-3354.

149. Khasnutdinova et al (2010)

Prevalence of stunting, underweight, overweight and obesity in adolescents in Velsk district, north-west Russia: a cross-sectional study using both international and Russian growth references. Public Health; 124:392-397.

- 150. Khor (2008)
 Cardiovascular epidemiology in the Asia–Pacific region. Asia Pacific Journal of Clinical Nutrition; 10(2): 76–80.
- 151. Khowaja et al (2010)
 Quit smoking for life--social marketing strategy for youth: a case for Pakistan.
 J Cancer Educ; 25(4): 637-642.
- 152. Khuwaja et al (2007)

Noncommunicable diseases and injuries: action needed in South Asia too. PLoS Med; 4(1): e38.

153. Kleinman et al (2002)

Diet, breakfast, and academic performance in children. Ann Nutr Metab; 46(1): S24–S30.

154. Kollias et al (2012)

Tracking of blood pressure from childhood to adolescence in a Greek cohort. Eur J Public Health; 22 (3):389-393.

155. Koukounari et al (2008)

Relationships between anaemia and parasitic infections in Kenyan Schoolchildren: a Bayesian hierarchical modelling approach. International Journal for Parasitology; 38: 1663–1671.

156. Kovacs et al (2010)

Role of waist circumference in predicting the risk of high blood pressure in children. *Int J Pediatr Obes;* 5(2): 143 – 150.

157. Krebs et al (2007)

Assessment of Child and Adolescent Overweight and Obesity. Available at: http://www.pediatrics.org/cgi/content/full/120/Supplement_4/S193

158. Kreuter et al (2003)

Achieving Cultural Appropriateness in Health Promotion Programs: Targeted and Tailored Approaches. Health Educ Behav; 30(2): 133-146.

159. Kuczmarski et al (2000)

Criteria for definition of overweight in transition: background and recommendations for the United States. Am J Clin Nutr; 72:1074–1081.

- 160. Kumar et al (2007)Prevalence of Obesity and Its Influencing Factor among Affluent School Children of Davangere City. Ind J of Community Med; 32(1): 15-17.
- 161. Kumar et al (2007)

Prevalence of obesity and its influencing factor among affluent school children of Davangere city. Indian J Community Med; 32:15-7.

162.	Kumari and Krishna (2011)
	Prevalence and risk factors for adolescents (13-17 years): overweight and
	obesity. CURRENT SCIENCE; 100(3): 373-377.
163.	Laxmaiah et al (2007)
	Factors affecting prevalence of over weight among 12 to 17 years old urban
	adolescents in Hyderabad, India. OBESITY; 15 (6): 1384-1390.
164.	Li et al (2005)
	Disease risks of childhood obesity in China. Biomed Environ Sci; 18: 401-
	410.
165.	Lob stein et al (2004)
	Obesity in children and young people: a crisis in public health. Obes Rev;
	5(1): 4 – 85.
166.	Lob stein et al (2004)
	Obesity in children and young people: A crisis in public health. Obesity
	Reviews; 5(1): 4-85.
167.	Ludwig et al (2001)
	Relation between consumption of sugar-sweetened drinks and childhood
	obesity: a prospective, observational analysis. The Lancet; 357(9255): 505-
	508.
168.	Lytle et al (2004)
	School-based approaches to affect adolescents' diets: results from the TEENS
	study. Health Education & Behavior; 31(2):270-287.
169.	Lytle LA et al (2006)
	Influencing healthful food choices in school and home environments: Results
	from the TEENS study. Preventive Medicine; 43(1):8–13.
170.	Madhusudan et al (2012)
	A study of blood pressure profile in school Going adolescents of district jaipur
	(rajasthan). National Journal of Community Medicine; 3(2): 242-246.
171.	Manios et al (2002)
	Health and nutrition education in primary schools of Crete: changes in chronic
	disease risk factors following a 6-year intervention programme. British Journal
	of Nutrition; 88(3):315-324.

.

172. Matijasevich et al (2009)

.

Socioeconomic position and overweight among adolescents: data from birth cohort studies in Brazil and the UK. *BMC Public Health*; 9: 105.

- 173. McCarthy and Ashwell (2006) A study of central fatness using waist-to-height ratios in UK children and adolescents over two decades supports the simple message-'keep your waist circumference to less than half your height. Int J Obesity; 30: 988-92.
- McGarvey et al (2004)
 Feasibility and benefits of a parent-focused preschool child obesity intervention. Am J Public Health; 94:1490–5.
- McKenzie et al (1995)
 Children's activity levels and lesson context during third-grade physical education. Res Q Exerc Sport; 66:184–93.
- McKenzie et al (1997)
 Long-term effects of a physical education curriculum and staff development program: SPARK. Res Q Exerc Sport; 68:280–91.
- McKenzie et al (1997)
 Long-term effects of a physical education curriculum and staff development program: SPARK. Res Q Exerc Sport; 68:280–91.
- McKenzie et al (2000)
 Student activity levels, lesson context, and teacher behavior during middle school physical education. Res Q Exerc Sport; 71:249–59.
- McKenzie et al (2000)
 Student activity levels, lesson context, and teacher behavior during middle school physical education. Res Q Exerc Sport; 71:249–59.
- McKenzie et al (2001)
 Effects of the CATCH Physical Education Intervention: Teacher Type and Lesson Location. American Journal of Preventive Medicine; 21(2):101–109.
- McKenzie et al (2004)
 Evaluation of a two-year middle-school physical education intervention: M-SPAN. Med Sci Sports Exerc; 36:1382–8.
- 182. McLean et al (2003)

Family involvement in weight control, weight maintenance and weight-loss interventions: a systematic review of randomised trials. Int J Obes Relat Metab Disord; 27:987–1005.

183. Mcmurray et al (2002)

A school-based intervention can reduce body fat and blood pressure in young adolescents. J. of Adol Health; 31(2): 125-132.

184. Mehan et al (2012)

Study of school environment and prevalence Of obesity & its predictors among adolescent (10-13 years) belonging to a private school in An urban indian city. National Journal of Community Medicine; 3(3): 400-407.

185. Misra et al (2011)

The High Burden of Obesity and Abdominal Obesity in Urban Indian Schoolchildren: A Multicentric Study of 38,296 Children. Ann Nutr Metab; 58(3):203–211.

- 186. Moayeri et al (2006)Over weight and obesity and their associated factors in adolescents in Tehran,Iran.2004-2005. Eur J Pediatr; 165(7):489-93.
- 187. Mogra and Kaur (2012) Association of body fat and blood pressure among school children (9-12 years). Indian Journal of Fundamental and Applied Life Sciences; 2 (1): 139-144.
- 188. Mogra et al (2012)

Association of body fat and blood pressure among school children (9-12 Years). Indian Journal of Fundamental and Applied Life Sciences; 2 (1): 139-144.

- 189. Montazerifar et al (2009)
 Prevalence of under weight, over weight and obesity among high school girls in Sistan va Baluchistan. Eastern Mediterranean Health Journal; 15(5): 1293-1300.
- 190. Monteiro et al (2003)
 Birth size, early childhood growth and adolescent obesity in a Brazilian birth cohort. *Int J Obes*; 27:1274-1282.
- 191. Morris and Zidenberg-Cherr (2002)

Garden-enhanced nutrition curriculum improves fourth-grade school children knowledge of nutrition and preferences for some vegetables. J of Am Diet Ass; 102 (1): 91-93.

- 192. Mostafavi et al (2005)Prevalence of obesity and over- weight in adolescents and adult population in Shiraz: Iran J Endcorinol Metab; 7(1):57-66.
- 193. Moyer (2013)

Screening for Primary Hypertension in Children and Adolescents: U.S. Preventive Services Task Force Recommendation Statement. Annals of Internal Medicine 2013. Available at: www.annals.org

- 194. Murphy et al (1998)The relationship of school breakfast to psychosocial and academic functioning. Arch Pediatr Adolesc Med; 152:899–107.
- 195. Murphy et al (1998)

The relationship of school breakfast to psychosocial and academic functioning. Arch Pediatr Adolesc Med; 152:899–107.

- 196. Murphy, Drake and Weineke (2005) Academics and breakfast connection pilot: final report on New York's Classroom Breakfast Project. Albany, NY: Nutrition Consortium of New York; 2005.
- 197. Murphy, Pagano and Bishop (2001)
 Impact of a universally-free, in-classroom school breakfast program on achievement: results from the Abell Foundation's Baltimore Breakfast Challenge Program. Boston, MA: Massachusetts General Hospital; 2001.
- 198. Musaiger et al (2011)The Paradox of Nutrition-Related Diseases in the Arab Countries: The Need for Action Int. J. Environ. Res. Public Health; 8(9): 3637-3671.
- Mushtaq et al (2011)
 Waist circumference, waist-hip ratio and waistheight ratio percentiles and central obesity among Pakistani children aged five to twelve years. BMC Pediatrics; 11:105.
- 200. Mushtaq et al (2011)

Waist circumference, waist-hip ratio and waistheight ratio percentiles and central obesity among Pakistani children aged five to twelve years. BMC Pediatrics; 11:105.

201. Must et al (1991)

Reference data for obesity: 85th and 95th percentiles of body mass index and triceps skin fold thickness. Am J Clin Nutr; 53:839-846.

202. Nader et al (1996)

The effect of adult participation in a school-based family intervention to improve Children's diet and physical activity: the Child and Adolescent Trial for Cardiovascular Health. Prev Med; 25:455–64.

- 203. Nader et al (2006)
 Identifying Risk for Obesity in Early Childhood. *Pediatrics*; 118(3): e594-601.
- 204. Narayanappa et al (2012)

Prevalence of Prehypertension and Hypertension among Urban and Rural School Going Children. Indian Pediatr: 49(9): 755-756.

- 205. National Association for Sport and Physical Education. National standards for beginning physical education teachers. 2nd ed. Reston, VA: National Association for Sport and Physical Education; 2009.
- 206. Naylor et al (2006)

Action Schools! BC: A socioecological approach to modifying chronic disease risk factors in elementary school children. Preventing Chronic Disease: Public Health Research, Practice and Policy; 3(2):156-160.

207. NCF (2005)

National Curriculum Framework (NCF, 2005) Available at: http://www.ncert.nic.in/rightside/links/pdf/framework/english/nf2005.pdf

- 208. Nead et al (2004)
 Overweight children and adolescents: A risk group for iron deficiency.
 Pediatrics; 114: 104-108.
- 209. NFHS III (2005-2006) Available at: http://www.rchiips.org/nfhs/pdf/India.pdf.
- 210. NFHS III 2005-2006.Available: http://www.rchiips.org/nfhs/pdf/India.pdf.
- 211. NIH (2005)

National Institute of Health. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. Available at: http://www.nhlbi.nih.gov/health/prof/heart/hbp/hbp_ped.pdf

212. NIH (2005)

The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. Available at: http://www.nhlbi.nih.gov/health/prof/heart/hbp/hbp_ped.pdf

213. NIN (2011)

National Institute of Nutrition. Dietary guidelines for Indians- A Manual.Available at: http://ninindia.org/DietaryGuidelinesforNINwebsite.pdf

- 214. Noronha et al (2012)
 High blood pressure in overweight children and adolescents. Journal of Human Growth and Development; 22(2): 196-201.
- 215. Noronha et al (2012)High blood pressure in overweight children and adolescents. Journal of Human Growth and Development; 22(2): 196-201.
- 216. O'Toole et al (2007)

Nutrition services and foods and beverages available at school: results from the school health policies and programs study 2006. J Sch Health; 77:500–21.

217. O'Toole et al (2007)

Nutrition services and foods and beverages available at school: results from the school health policies and programs study 2006. J Sch Health; 77:500–21.

218. Ogden et al (2010)

Prevalence of Obesity Among Children and Adolescents: United States, Trends 1963–1965 Through 2007–2008. Centre for disease control, Division of Health and Nutrition Examination Surveys 2010.

- 219. Önera et al (2004)
 Prevalence of under weight, over weight and obesity in Turkish adolescents.
 Swiss Med Wkly; 134: 529 533.
- 220. Önera et al (2004)
 Prevalence of under weight, over weight and obesity in Turkish adolescents.
 SWISS MED WKLY; 1 3 4 : 5 2 9 5 3 3.
- 221. Opara et (2010)

Prevalence of stunting, underweight and obesity in school aged children in Uyo, Nigeria. Pakistan Journal of Nutrition; 9 (5): 459-466.

- 222. Opara et al (2010)
 Prevalence of Stunting, Underweight and Obesity in School Aged Children in
 Uyo, Nigeria. Pakistan Journal of Nutrition; 9(5): 459-466.
- 223. Opara et al (2010)
 Prevalence of stunting, underweight and obesity in school aged children in Uyo, Nigeria. Pakistan Journal of Nutrition; 9 (5): 459-466.
- Ory, Jordan and Bazzarre2 (2002)
 The Behavior Change Consortium: setting the stage for a new century of health behavior-change research. HEALTH EDUCATION RESEARCH; 17(5):500-511.
- 225. Osganian et al (1996)Changes in the nutrient content of school lunches: results from the CATCHEat Smart Food service Intervention. Prev Med; 25:400–12.
- 226. Özgüven et al (2010)
 Evaluation of nutritional status in Turkish adolescents as related to gender and socio economic status. J Clin Res Ped Endo; 2(3): 111-116.
- 227. Özgüven et al (2010)

Evaluation of nutritional status in Turkish adolescents as related to gender and socio economic status. J Clin Res Ped Endo; 2(3):111-116.

228. Pande et al (2012)

Industrial city of western region (Case control study & Conclusion: screening of children for bp and fbg at early age). *Int J Pharm Biomed Sci;* 3(4): 229-232.

229. Patton et al (2012)
Health of the World's Adolescents: A Synthesis of International Comparable
Data. The Lancet; 379(9286): 1665-1675.

230. Perry et al (2004)

A randomized school trial of environmental strategies to encourage fruit and vegetable consumption among children. Health Education & Behavior; 31(1):65-76.

231. Pinhas-Hamiel et al (2003)

Greater prevalence of iron deficiency in overweight and obese children and adolescents. Int J Obesity; 27: 416-418.

232. Poskitt (2009)Countries in transition: underweight to obesity non- stop? Annals of Tropical

233. Powell et al (1998)
Nutrition and education: a randomized trial of the effects of breakfast in rural primary school children. Am J Clin Nutr; 68:873–9.

- 234. Prabhjot et al (2005)
 Variation in Blood Pressure Among School Children of Amritsar (Punjab).
 Anthropologist; 7(3): 201-204.
- 235. Prell et al (2005)

Paediatrics; 29: 1-11.

A school-based intervention to promote dietary change, Journal of Adolescent Health; 36(6):515–529.

- 236. Rachael et al (2007)
 APPLE Project: 2-y findings of a community-based obesity prevention program in primary school-age children1-3. Am J Clin Nutr; 86:735-42.
- 237. Rainville and Carr (2001)
 Competencies, knowledge, and skill statements for district school nutrition directors/supervisors. R-50-01. University, MS, National Food Service Management Institute; 2001.
- 238. Raja'a and Bin Mohanna (2005) Over weight and obesity among school children in Sana's City, Yemen. Ann Nutr Metab; 49(5):342-5
- 239. Rajalakshmi et al (2010)

A novel school-based intervention to improve nutrition knowledge in children: cluster randomised controlled trial. BMC Public Health; 10:123-131.

- 240. Rajeshwari et al (2005)
 Secular trends in children's sweetened-beverage consumption (1973 to 1994):
 the Bogalusa Heart Study. J Am Diet Assoc; 105(2):208-214.
- 241. Ralph et al (2006)
 Age at Drinking Onset and Alcohol Dependence: Age at Onset, Duration, and Severity. *Archives of Pediatric Adolescent Medicine*; 160(7)6): 739-46.
- 242. Ralph et al (2006)

Age at Drinking Onset and Alcohol Dependence: Age at Onset, Duration, and Severity. *Archives of Pediatric Adolescent Medicine;* 160(7)6): 739-46.

- 243. Ramachandran et al The Indian Diabetes Prevention Program shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia*; 49(2): 289-297.
- Ramachandran et al (2002)
 Prevalence of overweight in urban Indian adolescent school children. *Diabetes Res Clin Pract*; 57(3):185–90.
- 245. Ramchandran et al (2002)
 Prevalence of overweight in urban Indian adolescent school children. *Diabetes Res Clin Pract*; 57(3): 185-90.
- Rampersaud et al (2005)
 Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. J Am Diet Assoc; 105:743–60.
- 247. Ransdell et al (2003)

Daughters and mothers exercising together: effects of home- and communitybased programs. Med Sci Sports Exerc; 35:286–96.

- 248. Rees et al (1999)
 Improving the Nutritional Health of Adolescents –Position Statement Society for Adolescent Medicine. *Journal of Adolescent Health*; 24:461-462.
- 249. Rema and Vasanthamani (2011)

Prevalence of nutritional and lifestyle disorders among school going children in urban and rural areas of Coimbatore in Tamil Nadu, India. Indian Journal of Science and Technology; 4(2): 131-140.

 250. Rinderknecht and Smith (2004)
 Social cognitive theory in an after-school nutrition intervention for urban Native American youth. Journal of Nutrition Education & Behavior; 36(6):298–304. 251. Saha (2008)

Prevalence of hypertension and variation of blood pressure with age among adolescents in Chetla, India. Tanzania Journal of Health Research; 10(2): 108-111.

252. Saksvig et al (2005)

A pilot school-based healthy eating and physical activity intervention improves diet, food knowledge, and self-efficacy for native Canadian children. Journal of Nutrition; 135(10):2392–2398.

253. Sallis et al (1997)

The effects of a 2-year physical education program (SPARK) on physical activity and fitness in elementary school students. Sports, Play, and Active Recreation for Kids. Am J Public Health; 87:1328–34.

254. Sandercock et al (2010)

Associations between habitual school-day breakfast consumption, body mass index, physical activity and cardiorespiratory fitness in English schoolchildren. Eur. J. Clin. Nutr; 64(10): 1086–1092.

255. Sarma et al (2006)

Effect of micronutrient supplementation on health and nutritional status of school children: Growth and morbidity. Nutrition; 22: S8-14.

- 256. Schaff et al (2000) Prevalence of iron deficiency in Auckland high school students. N Z Med J; 113: 347-350.
- 257. Sen and Kanani (2006)

Deleterious Functional Impact of Anemia on Young Adolescent School Girls. Indian Pediatrics; 43: 219-226.

258. Seshadri et al (1999)

Studies on Nutritional Anaemia in pre-adolescent and adolescent children (6-15 y) of upper income groups from three urban centers in India. Project report jointly from Department of Foods and Nutrition, MS University of Baroda, Lady Irwin College, New Delhi and Women's Christian College, Chennai, 1999.

259. Shah et al (2008)

Assessment of Obesity in School Children. Calicut Medical Journal; 6(3):e2-e8.

- 260. Shah et al (2008) Assessment of Obesity in School Children. Calicut Medical Journal; 6(3):e2-6. 261. Shah et al (2008) Assessment of Obesity in School Children. Calicut Medical Journal; 6(3):e2e8. 262. Shah et al (2008) Assessment of Obesity in School Children. Calicut Medical Journal; 6(3):e2. 263. Sharbatti (2003) Anemia among adolescents. Saudi Med J; 24 (2): 189-194. 264. Sharma et al (2007) Growth pattern and prevalence of obesity in affluent school children of Delhi. Public Health Nutr; 10: 485-491. 265. Shephard (1996) Habitual physical activity and academic performance. Nutr Rev; 54(4):S32–S36. 266. Shrimpton (2006) Life Cycle and Gender Perspective on the Double Burden of Malnutrition and the prevention of Diet Related Chronic diseases in Diet-related Chronic diseases and the double burden of malnutrition in West Africa SCN News No. 33. 267. Sidhu et al (2005) Prevalence of overweight and obesity among the affluent adolescent school children of Amritsar. Punjab Coll Antropol; 29(1): 53-55. 268. Sidhu et al (2005) Prevalence of Overweight and Obesity among the Affluent Adolescent School Children of Amritsar, Punjab. Coll. Antropol; 29(1): 53-55. 269. Silva et al (2008) Diagnosing the nutritional status of schoolchildren: a comparison between Brazilian and international criteria. J Pediatr (Rio J); 84:550-555.
- 270. Simon et al (2004)

Intervention centred on adolescents' physical activity and sedentary behaviour (ICAPS): concept and 6-month results. International Journal of Obesity and Related Metabolic Disorders; 28(3):S96–S103.

271. Simon et al (2006)

- 272. ICAPS: a multilevel program to improve physical activity in adolescents.Diabetes & Metabolism; 32(1):41-49.
- 273. Sivakumar et al (2006)
 Effect of micronutrient supplementation on health and nutritional status of school children: study design. Nutrition; 22(1): S1-7.
- 274. Soudarssanane et al (2008)
 Tracking of Blood Pressure Among Adolescents and Young Adults in an Urban Slum of Puducherry. Indian journal of community medicine; 33(2): 107-112.
- 275. Srihari et al (2007)
 Nutritional status of affluent Indian school children: What and how much dowe know? *Indian Pediatr*; 44:204–13.
- 276. Srihari et al (2007)Nutritional Status of Affluent Indian School Children: What and How MuchDo We Know? Indian Pediatrics; 44:199-203.
- 277. Srihari et al (2007)

Nutritional Status of Affluent Indian School Children: What and How Much Do We Know? Indian Pediatrics; 44:199-203.

278. Steckler et al (2003)

Pathways process evaluation results: a school-based prevention trial to promote healthful diet and physical activity in American Indian third, fourth and fifth grade students. Preventive Medicine; 37(6):S80–S90.

279. Stettler et al (2002)

Prevalence and risk factors for overweight and obesity in children from Seychelles, a country in rapid transition: the importance of early growth. International Journal of Obesity; 26: 214–219.

- 280. Stevens et al (2003)
 - The impact of the Pathways intervention on psychosocial variables related to diet and physical activity in American Indian schoolchildren. Preventive Medicine; 37(6):S70–S79.
- 281. Steyn et al (2006)

Chronic diseases of lifestyle in South Africa: 1995-2005. Technical Report. Cape Town: South African Medical Research Council.Available at: http://www.mrc.ac.za/chronic/cdl1995-2005.pdf 282. Steyn et al (2009)

A review of school nutrition interventions globally as an evidence base for the development of the HealthKick programme in the Western Cape, South Africa. S Afr J Clin Nutr; 22(3):145-152.

283. Stigler et al (2011) Measuring Obesity among School-aged Youth in India: A Comparison of Three Growth References. Indian Pediatr; 48: 105-110.

284. Subramanyam et al (2003)

Prevalence of overweight and obesity in affluent adolescent girls in Chennai in 1981 and 1998, Indian Pediatr; 40: 332-336.

285. Sudhagandhi et al (2011)

Prevalence of anemia in the school children of Kattankuathur, Tamil Nadu, India. Int J of Nutrition, Pharmacology, Neurological Diseases; 1 (2): 184-188.

nicia. In 5 of Full tion, 1 harmacology, feetological Disea

286. Sundar et al (2013)

Prevalence and determinants of hypertension among urban school children in the age group of 13- 17 years in, Chennai, Tamilnadu. *Journal of Dental and Medical Sciences; 8(3): 14-20.*

287. Sundar et al (2013)

Prevalence and determinants of hypertension among urban school children in the age group of 13- 17 years in, Chennai, Tamilnadu. IOSR Journal of Dental and Medical Sciences; 8(3): 14-20.

288. Szajewska and Ruszczynski (2010)

Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. Crit. Rev. Food Sci. Nutr; 50: 113–119.

289. Taheri and Kazemi (2009)
Prevalence of over weight and obesity in 7 to 18 years old children in Birjand/Iran. Iranian Journal of Pediatrics; 19 (2): 135-140.

290. Tang et al (2007)Gender and SES differences in BMI of secondary high school students in HoChi Minh City. Asia Pac J Clin Nutr; 16 (1):74-83.

291. Tang et al (2007)Gender and SES differences in BMI of secondary high school students in HoChi Minh City. Asia Pac J Clin Nutr; 16 (1):74-83.

292. Taras (2005)

Nutrition and student performance at school. J Sch Health; 75:199-213.

- 293. Taveras et al (2005)
 Association of consumption of fried food away from home with body mass index and diet quality in older children and adolescents. Pediatrics; 116(4):18-24.
- 294. Thane, Bates and Prentice (2003)Risk factors for low iron intake and poor iron status in a national sample ofBritish young people aged 4-18 years. Public health Nutr; 6: 485-496.
- 295. Thiam et al (2006)

Diet related Chronic diseases in the west African Region in DIET-related chronic disease and the Double Burden of Malnutrition in West Africa. UNSCN News 33.

296. Thompson et al (2004)Food purchased away from home as a predictor of change in BMI z-score among girls, Int J Obes Relat Metab Disord; 28(2):282-289.

297. Tiwari (2000)

A study on anaemia control among adolescent girls - development of a school based intervention programme in Kathmandu, Nepal. PhD Thesis, M. S.University of Baroda, Vadodara, India 2000

298. Tremblay and Willms (2003)

Is the Canadian childhood obesity epidemic related to physical inactivity? International Journal of Obesity; 27: 1100–1105.

299. Trevino et al (2004)

Impact of the Bienestar school-based diabetes mellitus prevention program on fasting capillary glucose levels: a randomized controlled trial. Archives of Pediatrics & Adolescent Medicine; 158(9):911–917.

300. Trevino et al (2005)

Effect of the Bienestar Health Program on physical fitness in low-income Mexican American children. Hisp J Behav Sci; 27:120–32.

301. Uddin et al (2008)

Hemoglobin level among adolescent girls and it's relation to intestinal parasites. Bangladesh J. Zool; 33(2): 183-187.

302. UNICEF (2010)

United Nations, Department of Economic and Social Affairs, Population Division, World Population Prospects: The 2008 revision. Available at: www.esa.un.org/unpd/wpp2008/index.htm>

303. UNICEF (2010)

United Nations, Department of Economic and Social Affairs, Population Division, World Population Prospects: The 2008 revision. Available at: www.esa.un.org/unpd/wpp2008/index.htm>

304. UNICEF (2011)

State of world children 2011. Available at: http://www.unicef.org/sowc2011/

305. US (2010)

US Department of Education, Institute of Education Sciences. Educational indicators, indicator 24: time in formal instruction. Washington, DC: US Department of Education; 2010. Available at <u>http://nces.ed.gov/pubs/</u>eiip/eiipid24.asp.

306. US (2010)

US Department of Education, Institute of Education Sciences. Educational indicators, indicator 24: time in formal instruction. Washington, DC: US Department of Education; 2010. Available at <u>http://nces.ed.gov/</u>pubs/eiip/eiipid24.asp.

- 307. US Government Accountability Office. School meal programs: competitive foods are widely available and generate substantial revenues for schools. GAO-050563. Washington, DC: US Government Accountability Office; 2005.
- Werma and Singh (2012)
 Prevalence of hypertension in gujarati School going children and adolescents in Anand district. National Journal of Community Medicine; 3 (3): 452-457.
- 309. Verma and Singh (2012)
 Prevalence of hypertension in gujarati school going children and adolescents in anand district. National Journal of Community Medicine; 3(3): 452-457.
- 310. Verma et al (1998)
 Prevalence of anemia among urban school children of Punjab. Indian Pediatr;
 35: 1181-1186.
- 311. Wang and Lob stein (2006)

Worldwide trends in childhood overweight and obesity International Journal of Pediatric Obesity; 1: 11- 25.

- 312. Wang and Lob stein (2006)Worldwide trends in childhood overweight and obesity International Journal of Pediatric Obesity; 1(1): 11-25.
- Wang et al (2002) Trends of obesity and underweight in older children and adolescents in the United States, Brazil, China, and Russia. Am J Clin Nutr; 75 (6): 971-977.
- 314. Wardle et al (2006)
 - Development of adiposity in adolescence: five year longitudinal study of an ethnically and socioeconomically diverse sample of young people in Britain. BMJ 332:1130-1135
- 315. Warren et al (2003)

Evaluation of a pilot school programme aimed at the prevention of obesity in children. HEALTH PROMOTION INTERNATIONAL Vol. 18. No. 4 © Oxford University Press 2003.

- 316. Weiderpass (2010)Lifestyle and cancer risk. J Prev Med Public Health; 43(6): 459-471.
- 317. Whitaker et al (1997)
 Predicting obesity in young adulthood from childhood and parental obesity. N
 Engl J Med; 337(13):869-873.
- 318. WHO (1986)

Ottawa Charter for Health Promotion. WHO, Geneva. Available at: http://www.who.int/healthpromotion/conferences/previous/ottawa/en/

- WHO (1995)
 Physical status: the use and interpretation of anthropometry. Report of a WHO
 Expert Committee. World Health Organ Tech Rep Ser; 854:161-262.
- 320. WHO (1997) Obesity:

Preventing and managing the global epidemic. Report of a WHO consultation Geneva 3-5 January 1997.

321. WHO (2002)

The world health report 2002: Reducing risks, promoting healthy life. Geneva: World Health Organization.Available at: http://www.who.int/whr/2002/en/

322. WHO (2002)

The world health report 2002: Reducing risks, promoting healthy life. Geneva: World Health Organization. Available at: http://www.who.int/whr/2002/en/

323. WHO (2003)

Diet, Nutrition and Prevention of chronic diseases, WHO technical report series, Report of a joint WHO/FAO expert consultation, Geneva, 2003.

324. WHO (2004)

Global strategy on diet, physical activity and health. Geneva, World Health Organization, 2004 (WHA57.17). Available at: <u>http://apps.who.int/gb/ebwha/</u> pdf_files/WHA57/A57_R17-en.pdf

325. WHO (2004)

World Population Prospects: The 2004 Revision and World Urbanization Prospects.Available at: http://esa.un.org/unpp

- 326. WHO (2005) Nutrition in adolescence Issues and Challenges for the Health Sector. Available at: <u>http://whqlibdoc.who.int/publications /2005/</u> <u>9241593660 eng.pdf</u>
- 327. WHO (2005)

Nutrition in adolescence Issues and Challenges for the Health Sector.Available at: http://whqlibdoc.who.int/publications/2005/9241593660_eng.pdf

328. WHO (2006)

Nutrition Friendly School Initiative. Available at: <u>http://www.who.int/</u> nutrition/ topics/nut_school_aged/en/

329. WHO (2006)

NutritionFriendlySchoolInitiative.Availableat:http://www.who.int/nutrition/topics/nut_school_aged/en/

330. WHO (2006)

Adolescent Nutrition: A Review of the situation in selected south-east Asian Countries. World Health Organization, Regional office for South East-Asia, New Delhi.

- 331. WHO (2006)
- 332. Multicentre Growth Reference Study Group: WHO Child Growth Standards: Length/ height-for-age, weight-for- age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. Geneva, World Health Organization, 2006. Available at: <u>http://www.who.int/childgrowth/ standards/technical_report/en/</u> index.html

333. WHO (2006)

Nutrition Friendly School Health Initiative. Available at: http://www.who.int/nutrition/topics/nut_school_aged/en/

334. WHO (2007)

Assessing the iron status of populations. Second Edition. Publisher: WHO & CDC

335. WHO (2007)

Child and adolescent health and development. The adolescent with a chronic condition. WHO discussion paper on adolescence. Available at: http://whqlibdoc.who.int/publications/ 2007/9789241595704_eng.pdf.

336. WHO (2008) 2008-2013:

Action Plan for the Global Strategy for the Prevention and Control of Noncommunicable Diseases. Available at: <u>http://whqlibdoc.who.int/</u>publications/2009/9789241597418_eng.pdf

337. WHO (2008)

World wide prevalence of anemia 1993-2005. WHO global database on anemia. Available at: <u>http://www.who.int/vmnis/publications/ anaemia_prevalence/en/</u>

338. WHO (2009)

Global School Based Student Health Survey (GSHS) Core Questionnaire Modules. Available at: <u>http://www.who.int/chp/gshs/ GSHS_Core_</u> <u>Modules_2009_English.pdf</u>

339. WHO (2009)

Milestones in Health Promotion. Statements from Global Conferences. Available at: <u>http://www.who.int/healthpromotion/ Milestones Health</u> <u>Promotion 05022010.pdf</u>.

```
    WHO (2010).
    Global status report on non communicable diseases 2010. Available at:

    <u>http://www.cabdirect.org/abstracts/ 20113168808.html; jsessionid=</u>

    <u>BE97CD458133ED00700510A17046C3EC</u>
```

341. WHO (2011)

Global status report on non-communicable diseases 2010. Geneva. Available at: http://www.who.int/nmh/publications/ncd_report2010/en/

342. WHO (2011)

Global status report on non-communicable diseases 2010. Geneva. Available at: http://www.who.int/nmh/publications/ncd_report2010/en/

 343. WHO (2012)
 Child and adolescent health and development. Available at: http://www.who.int/child_adolescents_health/topics/ prevention_care/adolescents/en.

344. WHO (2012)

Child and adolescent health and development.Available at: http://www.who.int/child_adolescents_health/topics/

prevention_care/adolescents/en.

345. WHO (2012)

Tobacco Free Initiative, Geneva. Available at: http://www.who.int/tobacco/en/

346. Widenhorn-Muller et al (2008)

Influence on having breakfast on cognitive performance and mood in 13- to 20-year old high school students: Results of a crossover trial. Pediatrics; 122:279–284.

- Williams et al (2004)
 Cardiovascular risk reduction in preschool children: The "Healthy Start" project. Journal of the American College of Nutrition; 23(2):117–123.
- 348. Williamson et al (2007)

Wise Mind Project: A School-based Environmental Approach for Preventing Weight Gain in Children. OBESITY; 15(4): 906-917.

- 349. Wilsgaard et al (2005)
 Determining lifestyle correlates of body mass index using multilevel analyses: The Tromsø Study 1979–2001. Am J Epidemiol; 162(12): 1179–1188.
- Woodruff and Duffield (2000)
 Adolescents: assessment of nutritional status in emergency-affected adolescents. ACC/SCN: Geneva, 2000.