

Bibliography

- Adatia, R.D. and S.G. Gavde. 1962. Embryology of the Celastraceae. In *Plant Embryology - A symposium*, CSIR, New Delhi-11.
- Adinarayana, D. 1987. A new aromatic nitrosulphone from *Cardiospermum corindum*. *Lloydia*, 50 (4) : 620-622.
- Adkinson, J. 1913. Some features of the anatomy of the Vitaceae. *Ann. Bot.* 27: 133-139.
- Agnihotri, V.K., Srivastava, S. and S.K. Srivastava. 1987. A new limonoid, Amoorin, from the stem bark of *Amoora rohituka* Wall. *Curr. Sci.*, 56 (15) : 770-771.
- Ahmed, A.A. and N.A.M. Saleh. 1987. Peganetin, a new branched acylated tetraglycoside of acacetin from *Peganum harmala*. *Lloydia*, 50 (2) : 256-258.
- Ahmed, V.U., Bano, N., Bano, S., Fatima, A. and L. Kenne. 1986. Guaiainin, a new saponin from *Guaiacum officinale*. *Lloydia*, 49 (5) : 784-786.
- Airy Shaw, H.K. 1973. *A Dictionary of Flowering Plants and Ferns*. Willis. 7th edn., Cambridge Univ. Press, Cambridge.
- Amalraj, V.A. 1986. Floral biology of *Balanites roxburghii* Pl. J. Econ. Tax. Bot. 8 (1): 203-208.
- Amarasingham, R.D., Bisset, N.G., Millard, A.H. and M.C. Wood. 1964. Phytochemical survey of Malaya, Part III. Alkaloids and saponins. *Econ. Bot.* 18 (3) : 270.

- Anjaneyulu, A.S.R. and M.N. Rao. 1980. Elaeodendrol and elaeodendradiol, new triterpenes from *Elaeodendron glaucum*. *Phytochemistry* 19 : 1163-1170.
- Anjaneyulu, V., Prasad, K.H., Ravi, K. and J.D. Connolly. 1985. Triterpenoids from *Mangifera indica*. *Phytochemistry* 24 : 2359-67.
- Ansari, F.R., Ansari, W.H. and W. Rahman. 1978. A biflavone from *Garuga pinnata* Roxb. (Burseraceae). *Indian J. Chem.* 16 B : 846.
- Arndt, R.R. and W.H. Baarschers. 1964. The alkaloids of *Phyllia rogersii* Pillans. *J. Chem. Soc.* 2244-2248.
- Arora, N. 1953. The embryology of *Zizyphus rotundifolia* Lamk. *Phytomorphology*. 3 : 88-98.
- Aurada, E., Jurenitsoh, J. and W. Kubelka. 1984. Struktur der triterpen - sapogenine von *Aesculus glabra*. *Planta Med.* 50 (5) : 391-394.
- Baer, H., Hooton, M., Fales, H., Wu, A. and F. Schaub. 1980. Catecholic and other constituents of the leaves of *Toxicodendron radicans* and variation of urushiol concentrations within the plant. *Phytochemistry* 19 : 799-802.
- Bagehi, A., Sahai, M. and A.B. Ray. 1985. Phenolic constituents of *Rhus semialatus* leaves. *Planta Med.* (5) : 467-468.
- Bailey, A.E. and R.O. Aspluno. 1986. Isolation of methylgallate as the antitumor principles of *Acer saccharum*. *Lloydia* 49 (6) : 1149.

- Baillon, H.E. 1887. *Hist. Plantes*. 4 : 415.
- Bandaranayake, W.M. 1980. Terpenoids of *Canarium zeylanicum*. *Phytochemistry*. 19 : 255.
- Barbetti, P., Grandoline, G., Fardella, G. and I. Chiapinni, 1987. Indole alkaloids from *Quassia amara*. *Planta Med.* 53 (3) : 289-291.
- Basak, R.K. 1967. Studies on pollen morphology of Simarouba-
ceae. *Bull. Bot. Surv. India* 9 : 63-67.
- Basak, R.K. 1980. *Fascicles of Flora of India*, 4. BSI, Howrah.
- Bate-Smith, E.C. 1962. The phenolic constituents of plants
and their taxonomic significance. 1. Dicotyle-
dons. *J. Linn. Soc. (Bot)*. 58 : 95-110.
- Bate-Smith, E.C. 1972. Ellagitannin content of leaves of
Geranium species. *Phytochemistry* 11 : 1755-1757.
- Bate-Smith, E.C. 1978. Sytematic aspects of astringent tannins
of *Acer* species. *Ibid.* 17 : 1945-1948.
- Bate-Smith, E.C. 1973. Chemotaxonomy of *Geranium*. *J. Linn.
Soc. Bot.* 67 : 347-359.
- Bate-Smith, E.C. 1981. Astringent tannins of the leaves of
Geranium species. *Phytochemistry*. 20 : 211-216.
- Bate-Smith, E.C. 1983. A chemotaxonomic study of some Gerania-
ceae. *Ibid.* 22 : 2501-2505.
- Bate-Smith, E.C. and T. Swain 1966. The asperulesides and
the aucubins. In *Comparative Phytochemistry*.

(T.Swain ed.) Academic Press. London, 159
- 174

- Baxter, R.L. Crombie L., Simmonds, D.J. and D.A. Whiting.
1976. Extractives of *Catha edulis* (khat).
Occurrence of Celastraceous alkaloids having
mono-and bismacrolide bridges. J. Chem.Soc.
Chem.Comm. 12 : 463-464
- Baxter, R.L., Crombie, L., Simmonds, D.J., Whiting D.A., Braen-
den, O.J. and K. Szendrei. 1979. Alkaloids
of *Catha edulis* (khat) Part I. Isolation and
characterisation of eleven new alkaloids with
sesquiterpene cores (cathedulins): Identifica-
tion of quionone-methide root pigments. J.Chem.
Soc. Perkin Trans. I. 0 (12). 2965-2971.
- Behnke, H.D. 1981. Sieve-element character. Nord. J. Bot.
1 : 381-440.
- Behnke, H.D. 1982. Sieve-element plastids of Connaraceae
and Oxalidaceae, p-type plastids in dicotyle-
dons and their significance. Bot. Jahrb. Syst.
Pflanzenges. Pflanzengeogr. 103 (1) : 1-8.
- Bell, E.A. 1980. Non-protein amino acids in plants. In *Encyclo-
pedia of Plant Physiology*. New Series. (A.
Pirson and M.H. Zimmerman ed.) Vol. 8. Springer-
verlag, New York.
- Bennett, A.W. 1875. In *Flora of British India* (J.D. Hooker
ed.) Vol. 1, L. Reeve and Co. Ltd. London.
- Benthal, A.P. 1933 *The Trees of Calcutta*. Thacker Spink
and Co. Ltd. Calcutta.

- Bentham, G. and J.D. Hooker, 1862. *Genera Plantarum* Vol. 1 London.
- Berg, Van Den, R.G., Pollen morphology of genera *Pometia*, *Cubelia*, *Otonephelium* and *Litchi* (Sapindaceae -Nephelieae. *Blumea* 24 (2) : 369 - 394
- Bergman, G., Ganduik, E., Nyman, P. and L. Strid. 1977. The aminoacid sequence of stellacyanin from the lacquer tree. *Biochem. Biophys. Res. Commun.* 77 (3) : 1052-1959.
- Berikashivili, V. Sh., Paseshnichenko, V.A. and A.R. Gusava. 1982. - Terpenyl - - D- Glucopyranoside in leaves of *Pelargonium roseum*. *Prikl. Biokhim. MikroBiol.* 18 (3) : 431 - 433.
- Berkeley, E. 1953. Morphological studies in the Celastraceae *J. Elisha Mitchell Sci. Soc.* 69 : 185 - 206.
- Bessey, E.E. ¹⁹¹⁵ _A The phylogenetic taxonomy of flowering plants. *Ann. Missouri Bot. Gdn.* 2 : 109 - 164.
- Bestman, H., Classen, B., Kobald, V., Vostrowsky, O., Klingauf, F., and V. Stein. 1988. Steam volatile constituents from leaves of *Rhus typhina*. *Phytochemistry* 27 : 85.
- Betancor, C., Freire, R., Gonzalez, A.G., Salazar, J.A., Pascard, C. and T. Prangl. 1980. 3 Triterpenes and other terpenoids from *Catha cassinoides*. *Phytochemistry* 19 : 1989-1994.
- Bhandari, M. M. and V.S. Sharma. 1977. A new *Tribulus* (Zygophyllaceae) from India. *Bot. Notisier* 129 : 367 - 369.

- Bhatin K., Lah, J. and K.S. Ayyar. 1985. Barks as source of oxalic acid. *Indian Forester* III 7 : 538 - 541
- Bhatt, S.K., Saxena V.K. and S.S. Nigam. 1981. A new saponin from seeds of *Amoora rohituka*. *Phytochemistry* 20 : 1749 - 1750.
- Bhattacharya, M.K., Ghosh, P.K. and K.S. Mukherjee. 1981. Chemical investigation of the leaves of *Aesculus indica*. *J.Indian Chem. Soc.* 58 : 1011-1012.
- Bhide, K.S., Mujumdar, R.B. and A.V.R. Rao, 1977. Phenolics from the bark of *Chloroxylon swietenia* DC. *Indian J. Chem.* 15 B : 440 - 444.
- Bhutani, S.P., Chibber, S.S. and T.R. Seshadri, 1969. Flavonoids of the fruits and leaves of *Tribulus terrestris* : constitution of tribuloside. *Phytochemistry* 8 : 299-303
- Bianchi, E., Caldewell, M.E. and J.R. Cole. 1968. Antitumor agents from *Bursera fagaroides* (Burseraceae) (β - peltatin-methyl ether and 5-dimethoxy - β -peltatin). *Pharm. Sci.* 57 : 696 - 697
- Biesboer, D.D. 1975. Pollen morphology of the Aceraceae. *Grana* 15 : 19 - 27
- Biswas, H.G. 1948, A note on the saponin from *Sapindus laurifolius* Vahl. *J. Indian Chem. Soc.* 25 : 151 - 152
- Bohm, B.A. 1975. Chalcones, aurones and dihydroxychalcones. In *Flavonoids* (J.B. Harborne, T.J. Mabry and H. Mabry ed.) Chapman and Hall, London, 442.

- Bohm, B.A. and G.H.N. Towers. 1962. A study on phenolic compounds in *Impatiens*. *Can. J. Bot.* 40 : 677 - 683.
- Brandis, D. 1906, *Indian Trees* B.S.M. Pal Singh, Dehradun.
- Brieskorn, C.H. and P. Noble 1982. Constituents of essential oil myrrh. 11. Sesquiterpenes and furosesquiterpenes *Planta Med.* 44 : 87.
- Brown N.R. 1862, *A Handbook of the Trees, Shrubs and Heraceous Plants Growing in the Madras Horticultural Society and Garden and Neighbourhood of Madras.* Government press, Madras.
- Brown, S.A. 1965. Chemotaxomic aspects of lignin. *Lloydia* 28 (4) : 332 - 341
- Brown, W.A.C. and G.A. Sim. 1964. The constitution and absolute stereochemistry of Simarolides, the bitter principles of *Simarouba amara*. *Proc. Chem. Soc.* 293-294.
- Brunning, R. and H. Wagner. 1978. Review on the substances contained in the Celastraceae : Chemistry, chemotaxonomy, biosynthesis and pharmacology. *Phytochemistry* 17:1821 - 1858
- Campello, J.P. and A.J. Maraiota. 1975 Terebinthifolic acid and bracurenone : New triterpenoid ketones from *Schinus terebinthifolius*, *Ibid* 14 : 2300 - 2302.

- Carman, R.M. and D.E. Cowley. 1964. *Tetrahedron Lett.* 627.
- Carpenter, R.C., Satheeswaran, S., Sultanbawa, M.V.S. and S. Balsubramanian. 1980. (-) -5-Methyellein and catechol derivatives from four *Semecarpus* species. *Phytochemistry*, 19 : 445 - 448.
- Casinovi, C.G., Bellavita, V., Grandoline, G. and P. Cecchorelli. 1965. Occurence of bitter substances related to quassin in *Ailanthus glandulosa*. *Tetrahedron Lett.* 27 : 2273-2279
- Chalk, L. 1983. Wood anatomy, phylogeny and taxonomy. In *Anatomy of Dicotyledons* (C.R. Metcalfe and L. Chalk ed). 2nd Edn., Clarendon Press, Oxford, Vol. II : 108 - 125
- Chakraborty, B. and N.C. Neogi. 1978. Pharmacological properties of *Tribulus terrestris*. *Indian J. Pharm. Sci.* 40 (2) : 50 - 52
- Chaor, A., Charles, L. and S.N. Shamlan. 1980. A chromatographic study of anthraquinones of *Rhamnus alaternus* : Extraction, isolation and identification of the aglycones. *J. Crude Drug Res.* 18 (1) : 49 - 55.
- Chatterjee, B., Banerjee, S.N., Ganguly S.N. and S.N. Sircar. 1974. Triterpenes and coumarin from *Chukrasia tabularis*. *Phytochemistry*. 13 : 2012 - 2013.
- Chatterjee, A., Chakraborty, T. and S. Chandrasekharan. 1971. Chemical investigation of *Cedrela toona*. *Phyto-*

chemistry 10 : 2533.- 2535

Chen, F. and Y. Lin. 1975. Succedeneafavanone. A new, 6, 6"- binaringenin from *Rhus succedenea*. *Phytochemistry* 14: 1644 - 1647.

Chen, F. Y. and J. Wu. 1974. Rhusflavone-A new flavanoflavone from *Rhus succedenea*. *Ibid.* 18 : 1571 - 1574.

Chitnis, M.P., Bhatia, K.G., Phatak, M.K. and K.V.K. Rao. 1986. *Indian J. Exp. Biol.* 18 (1) : 6 - 8

Clair-Maczulajtys, D. and G. Bory. 1983. Pediceled extrafloral nectaries in *Ailanthus glandulosa*. *Can. J. Bot.* 61 (3) : 683 - 691

Cole, J.R. and R.M. Wledhopf. 1978, In *Chemistry of Lignans* (G.B.S. Rao ed). Andhra Univesity Press, India : 39-65.

Connolly, J.D., Okorie, D.A., Dewit, D.L. and D.A.H. Taylor. 1976. *J. Chem. Soc. Chem. Comm.* 909

Connolly J.D., Labbe, C. and D.S. Rycroft. 1978. Tetranortriterpenoids and related substances. Part 20. New tetranortriterpenoids from the seeds of *Chuckrasia tabularis* (Meliaceae); Simple esters of phragmalin and 12- α - acetoxypthagmalin. *J. Chem. Soc. Perkin Trans. 1* : 285 - 288

Cooke, R.G., Liv. A., Raston, C.L. and A.H. White. 1980. coloring matter of Australian plants : 22. *Ventilago*: crystalline structure and an approach to synthesis. *Aus. J. Chem.* 33 (2) : 303 - 312

- Copeland, H.F. 1966. Morphology and embryology of *Euonymus Japonica*. *Phytomorphology* 16 : 326 - 334.
- Corcilius, V.F. 1955. Papier chromatographische untersuchungen uber die oxycumarine und flavone aus der rosskastanie (*Aesculus hippocastanum* L.). *Planta Med.* 3 (1) : 55 - 61.
- Corner, E.J. H. 1976. *The Seeds of Dicotyledons*, Cambridge University Press, Cambridge, Vol. 2.
- Crawford, D.J. 1978. Flavonoid chemistry and angiosperm evolution. *Bot. Rev.* 44 (4) : 431 - 456.
- Crespi-Perrillina, N., Guicciardi, A., Malyszkó, G. and A. Minghetti. 1986. Biosynthetic relationship between indole alkaloids produced by cell cultures of *Ailanthus altissima*. *Lloydia* 49 (5) : 814.
- Crombie, L. 1980. The Cathedulin alkaloids. *Bull. Narc.* 32 (3) : 37 - 50
- Crombie, L., Crombie, M. and D.A. Whiting. 1978. Structure of cathedulin alkaloids from *Catha edulis* (khat), Kenyan and Ethiopian origin. *J. Chem. Soc. Chem. Commun.* 3 : 107 - 108.
- Cronquist, A. 1968. *The Evolution and Classification of Flowering Plants*, Nelson, London.
- Cronquist, A. 1981. *An Integrated System of Classification of Flowering Plants*, Colombia University Press, New York.

- Cronquist, A. 1987. A botanical critique of cladism. *Bot. Rev.* 53 (1) : 1 - 52.
- Cruz, M., Vitorina Da, A. and T. Santanna. 1984. Pollen studies in Sapindaceae. *Rev. Bras. Bot.* 7 (1) : 5 - 26.
- Dahlgren, R.M.T. 1980. A revised system of classification of the angiosperms. *Bot. J. Linn. Soc.* 80 : 91 - 124.
- Dahlgren, R. and K. Bremer, 1985. Major clades of angiosperms. *Cladistics* 1 : 349 - 368.
- Dahlgren, R.M.T., Jensen, S.R. and B.J. Nelson. 1981. A revised classification of the angiosperms. In *Phytochemistry and Angiosperm Phylogeny*. (D.A. Young and S. Sieglar ed.), Praeger Publishers, New York.
- Daniel, M. and S.D. Sabnis, 1979 . Chemotaxonomy of Loganiaceae *Curr. Sci.* 48 : 383 - 385.
- Darwish, F.A., Evans, F.J. and J.D. Philipson. 1980. Bruceolides and dehydrobruceolides from Fijian *Brucea javanica*. *Planta Med.* 39 : 232 - 233.
- Dawidar, A.A.M. and M.B.E. Fayez. 1969. Steroidal sapogenin-XIII. The constituents of *Balanites aegyptiaca*. *Phytochemistry* 8 (1) : 261 - 265.
- De Eds, F. 1968. In *Comprehensive Biochemistry* (M. Florkin and E.H. Stotz ed.) Elsevier Publishing Co., Amsterdam.

- De Jussieu, A.L. 1825 *Mem. Mus. Hist. Nat.* 12 : 450.
- Delvalle, D.M. and G. Schwenker. 1987. Preisocalamenediol. A constituent of *Schinus molle*. *Planta Med.* 53 (2) : 230.
- Demuth, G., Hing, H., Seligmann, O. and H. Wagner. 1978. Investigation on anthraquinoneglycosides of *Rhamnus* species V. Emodin-8-O- β -gentiobioside, a new O-glycoside from *Rhamnus frangula*. *Planta Med.* 33 (1) : 53 - 56.
- Deng, F., Jian-Hong, C., Zhi-Lin, X., Sui, L. and X. Wang. 1987. studies on the sesquiterpene alkaloids of *Tripterygium wilfordii* H.F. *Acta Bot. Sind.* 29 (5) : 523 - 526.
- Dev, K.P. 1972. The use of *Sapindus trifoliatus* in the treatment of chronic rheumatic arthritis. *Indian J. Pharm.* 34 (6) : 176.
- Dickson, W.C. 1986. Floral morphology and anatomy of Staphyleaceae. *Bot. Gaz.* 147 (3) : 312-326.
- Dickson, W.C. 1987. A palynological study of the staphyleaceae. *Grana* 26 : 11 - 24
- Dreyer, D. L., Pickering M.V. and P. Cohan. 1972. Distribution of limonoids in the Rutaceae. *Phytochemistry* 11 : 705 - 713
- Dwivedi, S.P.D., Pandey V.B., Shah, A.H. and G. Eckhart. 1987. Cyclopeptide alkaloids from *Zizyphus nummularia*. *Lloydia* 50 (2) : 235 - 237

- Eames, A.J. 1961. **Morphology of the Angiosperms**. Tata McGraw-Hill Publishing Co. Ltd. Bombay
- Edgeworth, M.P. and J. D. Hooker. 1875. **Flora of British India** (J.D. Hooker ed.) L. Reeve and Co. Ltd. London
- Ekong, D.E.L. and J.I. Okogun. 1969. Terpenoids of **Dacryoides edulis**. **Phytochemistry** 8 : 669 - 671
- El Hadidi, M.N. 1977. Tribulaceae as a distinct family. **Publ. Cairo Univ. Herb.** 7 & 8 : 103 - 108.
- El Negoumy, S.I., Al-Wakeel, S.A.M., El Hadidi, M.N. and N.A.M. Saleh. 1986. The flavonoids of the **Fagonia arabica** - complex (Zygophyllaceae). **Phytochemistry** 25 : 2423.
- El Oqlah. 1983. Pollen morphology of the genus **Erodium** in the Middle East. **Pollen Spores** 25 (3/4): 383 - 394.
- El Sissi, H.I. and N.A.M. Saleh. 1964. Phenolic components of **Mangifera indica** (Part-I). **Plant Med.** 12 : 421 - 427
- El Sissi, H.I. and N.A.M. Saleh. 1965. Phenolic components of **Mangifera indica** (Part-II). **Ibid** 13 : 346 - 352
- Engler, A. Anacardiaceae in Engler and Prantl, **Die Natürlichen Pflanzen Familien** III (5) : 138 - 178.
- Engler, A. 1931. In **Die Natürlichen Pflanzen Familien** (A. Engler and K. Prantl ed.) 2nd ed. 19a.1

- Erdtman, G. 1952. **Pollen Morphology and Plant Taxonomy in Angiosperms.**
Stockholm.
- Fahn, A. and R.F. Evert. 1974. Ultrastructure of the secretory ducts of *Rhus glabra* L. *Amer J. Bot.* 61 (1) : 1 - 14.
- Fagoonee, I. and G. Lauge. 1981. Noxious effects of neem extract on *Corcidolomi binotalis*. *Phytoparasitica* 9 (2): 111 - 118.
- Farooqui, P., and N.V. Subramanian. 1986. Epidermal structure of some important forest plants 4. *Sapindus*. *Indian Forester* 112 (6) : 543 - 553.
- Fong, H.H.S., Trojankova, M., Trojanek, T. and N.R. Farnsworth. 1972. Alkaloids screening. II. *Lloydia* 35 (2) : 117 - 149.
- Fowden L., Anderson, J.W. and M. Sung. 1970. Unusual amino acids of Sapindaceae. Biosynthesis and analogue function. *Phytochemistry* 9 : 916.
- Fowden, L. and A. Smith. 1969. Cyclopropane amino acids from *Aesculus* and *Blighia*. *Phytochemistry* 8: 437 - 443
- Fukuyama, Y., Mizuta, K., Nakagawa, K., Nenjuan, Q. and Waxiue. 1986. A new neo-lignan, a prostaglandin I_2 inducer from leaves of *Zizyphus jujuba*. *Planta Med.* (6): 501 - 502.
- Gamble, J.S. 1957. **Flora of Presidency of Madras**, BSI, Calcutta Vol. 1.

- Garg, H. S. and D.S. Bhakuni. 1984. An isoprenylated flavanone from leaves of *Azadirachta indica*. *Phytochemistry* 23 : 2115 - 2118.
- Gellert, M., Szendrei, K. and J. Reisch. 1980. Dihydromyricetin 3-O-rhamnoside from leaves of *Catha edulis* Ibid 20: 1759 - 1760.
- Ghanin, A., Jayaraman, I., Khan, H.A. and V.A. Amalraj. 1980. *Balanites roxburghii* fruits as source of diosgenin. *Indian J. Pharm. Sci.* 42 (5) : 149 - 150
- Gibbs, R.D. 1974. *Chemotaxonomy of Flowering Plants*, McGill-Queens, Univ. Press, London, Vol. III.
- Gieger, H. and C. Quinn. 1988. Biflavonoids In *The Flavonoids* (J.B. Harborne ed.) Chapman and Hall, London. 99 - 124.
- Giesman, T.A. 1962. *Chemistry of Flavonoid Compounds* Pergamon Press, Oxford.
- Gilbert, S.G. 1940. Evolutionary significance of ring porosity in woody angiosperms. *Bot. Gaz.* 102 : 105 - 120.
- Gilg, E. 1896. Vitaceae In *Die Natürlichen Pflanzen Familien* (A. Engler and K. Prantl ed.) Leipzig.
- Gilg, E. and M. Brandt, 1912. Afrikan Vitaceae. *Engler's Bot. Jahrb.* 46 : 547 - 549
- Gonazalez, A.G., Ferro, E.A. and A.G. Ranelo. 1987. Triterpenes from *Maytenus horrida*. *Phytochemistry* 26 : 2785 - 2788.

- Gorecki, P. and A. Mscisz. 1985. Determination of fraxin and intermediate products in the bark of the common horsechestnut in the presence of their coumarin glycosides. *Herba Pol.* 31 (3/4) : 197 - 200.
- Gornall, J. and B.A. Bohm. 1978. Angiosperm flavonoid evolution : A reappraisal. *Syst. Bot.* 3 : 353 - 368
- Gornall, J., Bohm, B.A. and R. Dahlgren. 1979. The distribution of flavonoids in the angiosperms. *Bot. Notiser.* 132 : 1-30.
- Govindappa, D.A. and G. Boriah. 1956. The development of female gametophyte in *Biophytum sensitivum* (L.) DC. *Curr. Sci.* 25 : 403 - 404
- Gray, D.D. and L. Fowden. 1962. 2-(Methylene cyclo-propyl) glycine from Litchi seeds. *Biochem. J.* 82 : 385 - 389.
- Grey-Wilson, C., 1980. Studies in Balsaminaceae : 6. Some observations on the floral vascular anatomy of *Impatiens*. *Kew. Bull.* 35 (1) : 221 - 228
- Grieve, C.M. and R.W. Soora. (1980) *Syst. Bot.* 5 : 39.
- Gulati N. and Mathur, S. 1977. Embryology and taxonomy of *Filicium decipiens* Thw. *Phytomorphology* 27 (3) : 261 - 266.
- Gunderson, A. 1950. *Families of Dicotyledons*. Chronica Botanica Company, Mass. U.S.A.

- Hall, B.A. 1954. Variability in the floral anatomy of *Acer negundo*. *Amer. J. Bot.* 41 (6) : 529 - 532
- Hallier, H. 1912. L' origine et le systeme phyletique des angiosperms
exposes a laide de leur arbre geneologique.
Arch. Néerl. Sci. Exáct. Nat. B. III I : 146
- 234.
- Harborne, J.B. 1967. *Comparative Biochemistry of the Flavonoids*. Academic Press, London.
- Harborne, J.B. 1977. Flavonoids and the evolution of the angiosperms. *Biochem. Syst. Ecol.* 5 : 7 - 22.
- Harborne, J.B. 1983. The flavonoids of the Rutales. In *Chemistry and Chemical Taxonomy of the Rutales* (P.G. Waterman and M.F. Grondon, ed.) Academic Press, London, 147 - 173
- Harborne, J.B. 1984. *Phytochemical Methods*. II edn. Chapman and Hall, London.
- Harborne, J.B. 1988. *The Flavonoids*. Chapman and Hall, New York.
- Harborne, J.B. and D.M. Smith. 1978. Correlations between anthocyanin chemistry and pollination ecology of the Polemoniaceae. *Biochem. Syst. Ecol.* 6 : 127 - 130.
- Harborne, J.B. and C.A. Williams. 1971. 6-Hydroxy luteolin and scutellarein as phyletic markers in higher plants. *Phytochemistry* 10 : 367 - 378.

- Hardin, J. W. 1956. Studies in the Hippocastanaceae. II. Inflorescence structure and distribution of perfect flowers. *Amer. J. Bot.* 43 : 418 - 424.
- Harms, H. 1931. Meliaceae, In *Die Natürlichen Pflanzen familien* (A. Engler and K. Prantl. ed.) Engelman, Leipzig.
- Harms, H. 1940. *Ibid.* 2, Bd. 19B 1 - 172.
- Harris, A., Anderson. L.A., Phillipson, J.D. and R.T. Brown. 1985. Canthin-6-one alkaloids from *Brucea antidysenterica* root bark. *Planta Med.* 151 - 153
- Hartley, T.G. 1969. A revision of the genus *Flindersia* (Rutaceae) *J. Arnold Arb.* 50 (4) : 481 - 526.
- Haslam, E. 1965. Galloyl esters in the Aceraceae. *Phytochemistry* 4 : 495 - 498.
- Hassall, C.H. and K. Reyle. 1955. Hypoglycin A and B, 2 biologically active polypeptides from *Blighia sapida*. *Biochem. J.* 10 : 334 - 339.
- Hattari, S. and H. Matsuda. 1952. Rhoifolin, a new flavone glycoside isolated from the leaves of *Rhus succedanea*. *Arch. Biochem. Biophys.* 37 : 85 - 89.
- Hayashi, N., Lee, K., Hall, I., Mophail, A.T. and H. Huang. 1982. Structure and stereochemistry of (-)-odorinol, an antileukemic diamide from *Aglaia odorata*. *Phytochemistry* 21 : 2371.

- Hegnauer, R. 1971. Pflanzenstaffe and pflanzen системати k. Naturwiss. 58 : 585 - 598
- Heimsch, C. 1942. Comparative anatomy of the secondary xylem in the 'Gruinales' and Terebinthales of Wettstein with reference to taxonomic grouping. Lilloa 8 : 83 - 198.
- Henning, W. 1966. Phylogenetic Systematics Translated by D.D. Davis and R. Xangeri, University Illinois Press, Urbana.
- Hiern, W.P. 1875. In Flora of British India (J.D. Hooker ed.) L. Reeve and Co., London Vol. I.
- Hill, A.F. 1952. Economic Botany, MacGraw - Hill Book Co. Inc. New York.
- Hinge, V.K., Wahg, A.D., Paknikar, S.K. and S.C. Bhattacharya. 1965. Terpenoids LXXI. Constituents of Indian black dammar resin. Tetrahedron 21 : 3197.
- Hindelang, D.M., Shamma, M., Miana, G.A., Shah, A.H. and B.K. Cassels. 1980. The carbon 13-NMR spectra of cyclopeptide alkaloids. Liebigs Ann. Chem. O (3) : 447 - 450.
- Hooker, J.D. 1879. Flora of British India L. Reeve and Co. Kent. Vol. II.
- Hopkins C.Y., Jevans, A.W. and Chisch. 1968. Fatty acids of Aceraceae seed oils. Canadian J. Biochem. 46 (9) : 999 - 1002.
- Hübel, W. and A. Nahrstedt. 1979. Cardiosperminsulphate, a sulphur containing cyanogenic glycoside

from *Cardiospermum grandiflorum*. *Tetrahedron Lett.* 45 : 4395 - 4396.

Hungund, B.L. and C.H. Pathak. 1971. A survey of plants in Gujarat, India, for alkaloids saponins and tannins. U.S.D.A. Forest Service Research paper NE - 201.

Hui-Zhenz. X., Lu, Z., Konno, C., Soejarta, D.D., Cordell, G.A., Fong, H.H.S. and W. Hodgson. 1988. 3- β -(3,4-dihydroxycinnamoyl) - Erythrodiol and 3- β -(4-hydroxycinnamoyl) erythrodiol from *Larrea tridentata*. *Phytochemistry* 27 (1) : 233 - 236.

Hutchinson, J. 1926. *The Families of Flowering Plants. Dicotyledons* Macmillan, London 328.

Hutchinson, J. 1969. *Evolution and Phylogeny of Flowering Plants*. Academic Press, London.

Hutchinson, J. 1973. *The Families of Flowering Plants*. Clarendon, Oxford. 3rd edn.

Ibrahim, R.K. and G.H.N. Towers, 1960. The identification, by paper chromatography of plant phenolic acids. *Arch. Biochem. Biophys.* 87 : 125 - 128.

Ikran, M. and S.M. Gilani. 1986. Antiinflammatory activity of *Aesculus indica* fruit oil. *Fitoterapia* 57 (6) : 455 - 456.

- Ikran, M., Khan, M.I. and N. Kawano. 1978. Chemical investigation of *Aesculus indica*. *Planta Med.* 34 (3): 337 - 338.
- Imamura, H., Shigerkiryama, H. and H. Ohasi. 1979. Heartwood constituents of *Rengas Gluta* species. *Res. Bull. Fac. Agric. GIFU Univ.* 0 (42) : 117 - 122.
- Inada, A., Kobayashi, M. and T. Nakanishi. 1988. Phytochemical studies on Meliaceae plants. III Structures of two new pregnane steroid, toosendansterols A & B, from leaves of *Melia toosendan*. *Chem. Pharma. Bull. (Tokyo)* 30 (2) : 609 - 612.
- Inamdar, J.A. 1969. Epidermal structure, stomatal ontogeny and relationship of some Zygophyllaceae and Simaroubaceae. *Flora B.* 158 : 360 - 368.
- Inamdar, J.A., Subramanian, R.B. and J.S.S. Mohana. (1986) Studies on resin glands of *Azadirachta indica* A. Juss (Meliaceae). *Ann. Bot.* 58 : 425 - 429.
- Inoue, T., Ishidate, Y., Fujita, M., Kubo, M., and M. Fukushima. 1978. Studies on constituents of Aceraceae plants : 1. Constituents of the leaves and the stem bark of *Acer nikoense* Maxim. *Yakugaku Zasshi.* 98 (1) : 41 - 55.
- Ishibashi, M., Murae T., Hirota, H., Tsuguki, T., Takahashi, T., Itai, A. and Y. Iitaka. 1982. Shingulactone C, a new quassinoid with 1 α , 12 α : 5 α , 13 α - dicyclo - 9 β -H-Picrasane skeleton from *Ailanthus altissima* Swingle. *Tetrahedron Lett.* 23 (1) 1205 - 1206.

- Ishiwata, H., Shizuri, Y. and K. Yamada. 1983. 3-Sequiterpene alkaloids from *Euonymus alatus* var. *striatus*. *Phytochemistry* 22 : 2843 - 2846.
- Ishratullah, K.H. Ansari, W.H. and W. Rahman. 1977. Biflavonoids from *Semecarpus anacardium* L. (Anacardiaceae). *Indian J. Chem. Sect. B Org. Chem. Incl. Med. Chem.* 15 (7) 615 - 619.
- Iyer V.S., and S. Rangaswami. 1972. Indacanthinone, a crystalline chemical component from the wood of *Samadera indica*. *Curr. Sci.* 41 : 140 - 141.
- Iwu, M., Onyechlobido, M. and M. Anazodo, 1986. Biochemical mechanism of the antimalarial activity of *Azadirachta indica* leaf extract. *Pharmacol. Res. Commun.* 18 (1) 81-91.
- Jefferies, P.R. and T.G. Payne, 1967. Diterpenes of *Dodonea* species. *Tetrahedron Lett.* 48 : 4777-4782.
- Jensen, R.D. 1981. Wagner networks and wagner trees : A presentation method for estimating most parsimonious solution. *Taxon* 30 (3) : 576-590.
- Jensen, S.R., Nielson, B.J. and R.D. Dahlgren. 1975. Iridoid compounds, their occurrence and systematic importance in Angiosperms. *Bot. Notiser* 128:148-180
- Joel, D.M. 1981. The duct system of base and stalk of the mango fruit. *Bot. Gaz.* 142 (3) : 329-33.
- Joel, D.M. and A. Fahn. 1980. Ultrastructure of resin ducts of *Mangifera indica* L. (Anacardiaceae) III.

Secretion of the protein-polysaccharide mucilage in the fruit. *Ann. Bot.* 46 : 785-790.

Johri, B.M. and R. Ahuja. 1957. A contribution to the floral morphology and embryology of *Aegle*. *Phytomorphology* 7 : 10-24.

Joshi, A.C. 1938. Parthenocarpy in *Dodonea viscosa*. *J. Indian Bot. Soc.* 17 : 97-99.

Joshi, B.S. and V.R. Hegde. 1979. Extraction of *Balsamodendron pubescens* Hook. Isolation and a new synthesis of siderin. *Proc. Indian Acad. Sci.*, 88A (3) : 185-190.

Joshi, K.C., Prakash, L. and P. Singh. 1973. Quinones and other constituents from *Tabebuia rosea*. *Phytochemistry* 12 : 942-943.

Jussieu, A. 1791. *De Genera Plantarum Secundum Ordines Naturales Disposita : Turici Helvitorum, Sumtibus Ziegleri Q. Filliorum*, 405.

Kakarni, H.K. 1982. *Fitoterapia* 5 : 221.

Kapil, R.N. and K. Ahluwalia, 1963. Embryology of *Peganum harmala* L. *Phytomorphology* 13 : 127-140.

Kapoor, S.K., Ahmed, R.I. and A. Zaman, 1971. Chemical constituents of *Ailanthus excelsa*. *Phytochemistry* 10 : 3333.

Karnik, M.G. and A.K. Misra. 1964. Maple wood (*Acer caesium*) for tooth pick-industry. *Indian Forester* 90 (5) : 310-311.

- Kasai, R., Mishi, M., Mizutani, K., Mirahara, I., Moriya, T. and K. Miyahar. 1988. Trifolioside 11. An acyclic sesquiterpene oligoglycoside from pericarp of *Sapindus trifoliatus*. *Phytochemistry* 27 (7) : 2209-2211.
- Kashyap, G. 1957. Studies in the Vitaceae II. Floral anatomy of *Vitis trifolia* L., *vitis latifolia* Roxb. and *Vitis himalayana* Brandis. *J. Indian Bot. Soc.* 36 : 317-323.
- Kawai, K., Akiyama, T., Ogihara, Y. and S. Shibata. 1974. A new sapogenin in the saponins of *Zizyphus jujuba*, *Hovenia dulcis* and *Bacopa monniera*. *Phytochemistry* 13 : 2829-2832.
- Khalid, S.A. 1983. Chemistry of Burseraceae. In *Chemistry and Taxonomy of Rutales*, (P.G. Waterman and M.E. Grondon ed.) Academic Press, London.
- Khar M., Shah, G.N., Prakar, S. and A. Zaman. 1986. Pallidol, a new reservatrol dimer from *Cissus pallida*. *Phytochemistry* 25 : 1945-1948.
- Khan, S.A. and K.M. Shamsuddin. 1978. Quassinoids from *Ailanthus excelsa*. *Ind. J. Chem.* 16 B : 1045-1046.
- Khan, S.A. and K.M. Shamsuddin. 1981. I-Hydroxy canthin-6-one, an alkaloid from *Ailanthus giraldii*. *Phytochemistry* 20 : 2062-2063.
- King, F.E. and J.W.W. Morgan. 1960. The chemistry of extraction from hardwood part XXX. The constituents of Katonic acid, a triterpene from *Sandoricum indicum*. *J. Chem. Soc.* 4738-4747.

- Koike, K. and T. Ohmoto, 1988. Picrasidie-U, dimer alkaloid from *Picrasma quassioides*. *Phytochemistry* 27 : 3029-3030.
- Korie, D.A. and D.A.H. Taylor, 1971. Limonoids from *Swietenia humilis*. *Ibid* 10 : 469-470.
- Kraus, W. and R. Cramer. 1981. Pentanortriterpenoids from *Azadirachta indica* (Meliaceae) *Chem. Ber.* 114 (7) : 2415-2422.
- Kraus, W. and K. Kypke. 1979. Surenone and surenin, two novel tetranortriterpenoids from *Toona sureni* (Blume) Merrill. *Tetrahedron. Lett.* 2715-2716.
- Kribs, D.A. 1930. Comparative anatomy of the woods of the Meliaceae. *Amer. J. Bot.* 17 : 724-738.
- Krishna, J.C., Singh, B. and T. Parthasani. 1978. Chemical constituents of *Gymnosporia montana* and *Euonymus pendulus*. *Planta Med.* 34 (2) : 211-214.
- Kumar, A. 1976. Studies on Geraniales VII. The embryology of *Erodium stephanianum* Willd. *Acta Bot. Indica* 4 (12) : 105-110.
- Kumar, A. 1977. Studies in Geraniales V. The embryology of some Indian species of *Geranium* L. *Indian J. Bot. Soc.* 56 (4) : 328.
- Knuth, C.S. 1824. *Annals of Science Naturala* Series 1. Part-II : 333.
- Kupchan, S.M., Takasugi, M., Smith, R.M. and P. S. Steyn. 1971. Tumor inhibitors. LXII. The structures

of acerotin and acerocin, novel triterpene ester aglycones from the tumor inhibiting saponins of *Acer negundo*. *J. Org. Chem.* 36 (14) : 1972-1976.

Lagarias, J.C. Goff. D. and H. Rapoport. 1979. Cyclopeptide alkaloids : Rhencyclopeptides from *Ceanothus sanguineus*. *Lloydia* 42 (6) : 663-668.

Lassak, E.V., polonsky, J. and H. Jacquemin 1977. 5 - Hydroxy canthin-6-one from *Simarouba amara*. *Phytochemistry* 16 : 1126.

Lawrence, G.H.M. 1951. *Taxonomy of Vascular Plants*. Oxford and IBH Publishing Co., New Delhi.

Lawson, M.A. 1875. In *Flora of British India* (J.D. hooker ed.) L. Reeve and Co. Ltd. London Vol. I.

Lee, S.M., Olsen., J.I., Schweizer, M.P. and J. Kloeche. 1988. 7-Deacetyl-17- β -hydroxyazadiradione, a new limonid insect growth inhibitor from *Azadirachta indica*. *Phytochemistry* 27 : 2773-2775.

Lersten, N.R. and S.R. rugentsein. 1982. Foliar nectaries in mahogany (*Swietenia* Jacq.). *Annals Bot.* 49 : 397-401.

Lin, Y. and F. Chen, 1974. Agathisflavone from drupes of *Rhus succedanea*. *Phytochemistry* 13 : 657-658.

Linde, H. 1979. Constituents of pericarps of *Sapindus mukorossi* Gaertn. *Arch. Pharm.* 312 (5) : 416-425.

Lindley, J. 1830. *Introduction to the Natural System of Botany*.

- Lindley, J. 1853. **The Vegetable kingdom**, London.
- List, A. and E.C. Steward. 1965. The nucellus, embryosac, endosperm and embryo of *Aesculus* and their interdependence during growth. **Annals Bot.** II 29 : 1-15.
- Lloyd, H.A., Jaouni, T.M. Evans, S.L. and J.F. Morton. 1977. Terpenes of *Schinus terebinthifolius*. **Phytochemistry** 16 : 1301-1302.
- Lösner, Th. 1897. In **Die Natürlichen Pflanzenfamilien** (A. Engler and K. Prantl ed.) Wilhelm Engelman, Leipzig. 3 (15).
- Lu King, M., Chiang, C., Ling. H., Fujita, E. Ochiai, M. and A.T. McPhail. 1982. X-ray crystal structure of rocaglamide, a novel antileukemic 1H-cyclopenta-(b)-benzofuran from *Aglaia elliptifolia*. **J. Chem. Soc. Chem. Commun.** 1150-1151.
- Luo, A., Merksuriyen, D., Erdelmeier, D.A.I., Fong, H.H.S. and G.A. Cordell. 1988. Larreantin, a novel cytotoxic naphthaquinone from *Larrea tridentata*. **J. Org. Chem.** 53 (10) : 2183-2185.
- Mabry, T.J., Markham, K.R. and M.B. Thomas. 1970. **The Systematic Identification of Flavonoids**. Springer - Verlag, Berlin.
- Mahato S.B., Sahu, N.P. Ganguly, A. N., Miyahara, K. and T. Kawasaki. 1981. Steroidal glycosides of *Tribulus terrestris*. **J. Chem. Soc. Perkin Trans, I.** 20 (9) : 2405-2410.
- Maheswari. P. 1964. **Recent Advances in Embryology of Angiosper-**

ms, Catholic Press, Ranchi.

- Maksoud, S.A. and M.N. El Hadidi. 1987. The flavonoids of *Fagonia bruguieri* complex (Zygophyllaceae) Plant Syst. Evol. 155 (1-4) : 311-318.
- Maksoud, S.A. and M.N. El Hadidi. 1988. The flavonoids of *Balanites aegyptiaca* (Balanitaceae) from Egypt. Plant Syst. Evol. 160 (3-4) : 153-158.
- Mangoni, L. Monaco, P. and L. Previtera. 1982. A new monoterpene from bled resin of *Pistacia vera*. Phytochemistry 21 : 811-812.
- Manske, R.H.F. 1944. The Alkaloids. Ann. Rev. Biochem. 13 : 533-548.
- Maradufu, A. 1982. Furosesquiterpenes of *Commiphora erythraea* and *C. myrrh*. Phytochemistry 21 : 677 - 678.
- Marker, R.E. 1943. Diosgenin from *Balanites aegyptiaca*. J. Chem. Soc. 65 : 1248.
- Markham, K.R. 1982. Techniques of Flavonoid Identification, Academic Press, London.
- Masand, P. 1963. Embryology of *Zygophyllum fabago* L. Phytomorphology 13 : 293.
- Masayoshi, K., Nagai, M. and T. Inoue. 1983. The constituents of Aceraceae plants. 4. C-13 NMR of acerogenin A, rhododendrol and related compounds and structure of aceroside IV from *Acer nikoense*. Chem. Pharm. Bull. (Tokyo) 31 (6) : 1917 - 1922.

- Mathur, A. and M.M. Bhandari. 1983. Studies on pollen grain of *Fagonia* and *Seetzenia* sps. *J. Econ. Tax. Bot.* 4 (2)
- Mathur, S. and Gulati, 1980. Embryology and taxonomy of *All-ophylus alnifolius* Radlk. Ex Engl. (Sapindaceae). *Indian J. Bot.* 3 (1) : 103 - 112
- Melville, G.N. and J.N. Addae. 1988. Effects of ackee fruit extracts on bronchomotor tone in rats. *West Indian Med. J.* 37 (2) 97 - 99.
- Merkuza, V.M., Mascaretti, O.A., Crohare, R and E.A. Ruveda. 1971. Triterpenoids from *Discaria longispina* and *Colletia paradoxa*. *Phytochemistry* 10 : 908 - 909.
- Merkuza, V.M., Sierra, M.G., Mascaretti, O.A., Ruveda, E.A., Chang, C. and E. Wenkert. 1974. Peptide alkaloids of *Discaria longispina* and *Scutia buxifolia*. *Phytochemistry* 13 : 1279 - 1282.
- Mester, I. 1983. Structural diversity and distribution of alkaloids in the Rutales. In *Chemistry and Chemical Taxonomy of the Rutales* (P.G. Waterman and M.F. Grundon ed.) Academic Press, London, 31 - 96.
- Metcalf, C.R. and L. Chalk. 1950. *Anatomy of the Dicotyledons*. Clarendon Press, Oxford.
- Metcalf, C.R. and L. Chalk. 1983. *Anatomy of the Dicotyledons* 2nd edn. II, Clarendon Press, Oxford.
- Meyer, S.F.H., Jawetz, E. and Goldfine. 1972. Review of Medi-

cal Pharmacology, 3rd edn. Lange Medical Publications, California.

- Mikolajizak, K.L. and D. Weisleder. 1978. ¹³C NMR spectroscopy of cyanolipids and cyanolipid containing seed oils. *Lipids* 13 (7) : 514 - 516.
- Milagros, M. and S. Machado. 1987. Pollen grains of Simaroubaeae, *Acta Bot. Cabana* 0 (45) : 1 - 7
- Milner, R.D.G. and P.K. Wirdnam. 1977. Hypoglycin stimulates insulin secretion. *Diabetologia* 13 (6) 637 - 238.
- Misra, G.S. and S.M. Chandhok. 1981. Calyculatone a new anthraquinone dye from stem bark of *Ventilago calyculata*. *Indian J. Chem.* 20 B : 721 - 722.
- Mitra C.R., Garg H.S. and G.N. Pandey. 1970. Constituents of *Melia indica* : Nimbidic acid and nimbidinin. *Tetrahedon Lett.* 2761.
- Moore, J.D. 1936. Floral anatomy and phylogeny in the Rutaceae. *New phytologist*, 25 : 318 - 322.
- Moore, M.O. 1986. A study of selected taxa of *Vitis* cultivated in the Southeastern U.S.A. *Rhodora* 89 (857) : 75 - 92.
- Moore, M.O. and E.D. Giannasi, 1987. Foliar flavonoids of selected *Vitis* taxa in South Eastern U.S.A. *Biochem. Syst. Ecol.* 15 (1) : 79 - 84.
- Mukherjee, K.S. and M.K. Bhattacharya. 1983. Phytochemical

- investigations of *Aesculus indica* L. and *Fagonia indica* Andr. J. Indian Chem. Soc. 60 : 507 - 508.
- Mukherejee, S.K. and D. Chandra, 1983. An outline of the revision of Indian Anacardiaceae Bull. Bot. Surv. Ind. 25 (1) : 52 - 24.
- Mulay, B.N., Nair, N.C. and M.S.R. Sastry. 1953. Contribution to the embryology of Vitaceae I. Male and female gametophyte of *Vitis pedata*. Proc. Rajasthan Acad. Sci. 4 : 17 - 28.
- Müller, J. 1979. Form and function of Angiosperm pollen. Ann. Miss. Bot. Gar. 66 (4) : 593 - 632.
- Müller, J. and P.W. Leenhouts. 1976. A general survey of pollen types in Sapindaceae in relation to taxonomy. Linn. Soc. Symp. Ser. No.1 : 407 - 445.
- Murray, R.D.H., Mendez., J. and S.A. Brown, 1981. The Natural Coumarins, Occurence, Chemistry and Biochemistry, John Wiley and sons Ltd.London.
- Murthy, S.S.N., Anjaneyulu, Rao, L.R., Peter, A. and R.S. Ward. 1982. Chemical examination of *Anacardium occidentale* : Isolation and structure determination of a novel biflavonoid C-glycoside. Planta Med. 45 (1) : 3 - 10.
- Murthy, Y.S. and S. Gupta, 1978. Morphological studies on the Meliaceae - III. A reinvestigation of floral anatomy of *Azadirachta* and *Melia*. J. Indian Bot. Soc. 57 : 195 - 204.

- Nag, T.N. and M.L. Harsh, 1982. Arid zone plants of Rajasthan : A source of steroidal sapogenins. *Acta Bot. Indica* 10 (1) : 8 - 11.
- Nagai, M., Kubo, M., Takashai, K., Fujita, M. and T. Inoue, 1983. Studies on constituents of Aceraceae plants 5. 2-Diaryl heptanoid glycosides and an aryl butanol apiosyl glucoside from *Acer nikoense*. *Chem. Pharm. Bull. (Tokyo)* 31 (6) : 1923 - 1928.
- Nagasampagi, B.A., Sriraman, M.C., Yankov, L. and Sukhdev, 1973. Siderin from *Cedrela toona*. *Phytochemistry* 14 : 1673.
- Naik, R.G., Kattige, S.L., Bhatt, S.V., Alreja, B., De Souza, N.J. and R.H. Rupp. 1988. An antiinflammatory cum immunomodulatory piperidiny benzopyranone from *Dysoxylum binectariferum*, isolation, structure and total synthesis. *Tetrahedron* 44 (7) : 2081-2086.
- Nair, A.G.R., Kotyal, J.P. and D.K. Bharadwaj. 1983. Myricetin - 7,4'-dimethyl ether and its 3-galactoside from *Rhus lancea*. *Phytochemistry* 22 : 318-319.
- Nair, A.G.R. and S.S. Subramanian, 1975. Isorhamnetin and quercetin glycosides from *Dodonea viscosa* and *Sapindus emarginatus*. *Indian J. Chem.* 13 : 639-640.
- Nair, A.G.R. and S.S. Subramanian, 1975. Quercetin glycosides from the leaves of *Soymda febrifuga* and *Melia azedarach*. *Indian J. Chem.* 13 : 527.

- Nair, A.G.R. and S.S. Subramanian, and K. Sriharan, 1963. Chemical examination of flowers of *Lannea coromandelica*. *Curr. Sci.* 32 (3) : 115-116.
- Nair, N.C. 1959. Floral morphology and embryology of *Naregamia alata* W. & A. *J. Indian Bot. Soc.* 38 : 353-366.
- Nair, N.C. 1963. Studies on Meliaceae VI. Morphology and anatomy of the flower of tribe Cedrelaceae and discussion on the floral anatomy of the family. *Ibid* 42 : 177-184.
- Nair N.C. 1968. Contribution of the floral morphology and embryology of two species of *Leea* with a discussion on the taxonomic position of the genus. *Ibid* 47 : 193 - 205.
- Nair, N.C. and I. Gupta, 1961. A contribution to the floral morphology and embryology of *Fagonia cretica* Linn. *Ibid* 40 : 635-640.
- Nair, N.C. and K. Kanta, 1961. Studies on Meliaceae IV : Floral morphology and embryology of *Azadirachta indica* Juss. A reinvestigation. *Ibid* 40 : 382-396.
- Nair N.C. and T. Joseph. 1960. Morphology and embryology of *Cardiospermum halicacabum* L. *Ibid.* 39 : 176 - 194.
- Nair, N.C. and K.V. Mani, 1960. Organography and floral anatomy of some species of the Vitaceae. *Phytomorphology* 10 : 138-144.
- Nair, N.C. and V. Parasuraman, 1954. Gametophyte of *Vitis pallida* W. & A. *Curr. Sci.* 23 : 163-164.

- Nair, N.C. and V. Parasuraman. 1962. Embryological studies on *Vitis pallida*. *Phyton* 18 : 157-164.
- Nair, N.C. and V.S. Sarma, 1961. Organography and floral anatomy of some members of the Rhamnaceae. *J. Indian Bot. Soc.* 40 : 47-55.
- Nair, N.C. and N.P. Sukumaran, 1960. Floral morphology and embryology of *Brucea amarissima*. *Bot. Gaz.* 121 : 175-185.
- Narayana, H.S. and P.K. Arora. 1963. Floral anatomy of *Monsonia senegalensis*. *Curr. Sci.* 32 : 185-187.
- Narayana, H.S. and C.G.P. Rao. 1963. Floral morphology and embryology of *Seetzenia orientalis* Decne. *Phytomorphology* 13 : 197-205.
- Narayana, L.L. 1959. Floral anatomy of Meliaceae-II. *J. Indian Bot. Soc.* 38 : 288-295.
- Narayana, L.L. 1965. Contribution to the embryology of Balsaminaceae-II. *J. Jap. Bot.* 40 : 104-116.
- Narayana, L.L. 1966. A contribution to floral anatomy of Oxalidaceae. *Ibid* 41 : 321-328.
- Narayana, L.L. and M. Sayeeduddin. 1958. Floral anatomy of Simaroubaceae-I. *J. Indian Bot. Soc.* 37 : 517-522.
- Narayana, L.L. and M. Sayeeduddin 1959. A study of gametophytes in *Impatiens leschenaultii* Wall. *Ibid.* 38 : 391-397.

- Narayanan, V., Seshadri, R. and T.R. Seshadri. 1973. Acergen-
in from the heartwood of *Acer caesium*. *Curr.*
Sci. 42 : 642-643.
- Nawwar, M.A.M., Ishak, M.S., Michael, H.N. and Buddrus, J.
1984. Leaf flavonoids of *Zizyphus spina-christi*.
Phytochemistry 23 : 2110-211.
- Nettleship, L. and M. Staylor. 1971. Ruine : A glucosidic
-carboline from *Peganum harmala*. *Ibid* 10
: 231-234.
- Nishizawa, M., Yamagishi, T., Gen-Ichiro nonaka and I. Nish-
loka. 1982. Tannins and related compounds
5. Isolation and characterisation of polygal-
loyl glucose from Chinese gallotannins. *J.*
Chem. Soc. Perkin Tras. 10 (12) 2963 - 2968.
- Nooteboom, H.P. 1962. Simaroubaceae. In *Flora Malaysiana*
6 : 194
- Nooteboom, H.P. 1966. Flavonols, leucoanthocyanins, cinnamic
acids and alkaloids in dried leaves of some
Asiatic and Malaysian Simaroubaceae. *Blumea*
14 (2) : 309 - 311
- Oohi, M., Kotsuki, H., Hirotsu, K. and To Tokoroyama. 1976.
Sendanin, a new limonoid from *Melia azedarach*
L. Var. *Japonica* Nakino. *Tetrahedron Lett.*
2877-2880
- Ohmoto, T. and K. Koike. 1983. The constituents of *Picrasma*
quassioides : 2. The alkaloidal constituents.
Chem. Pharm. Bull. (Tokyo) 31 (9) : 3198 -
3204.

- Okano, M., Lee, K. and I.H. Hall. 1981. Antitumor agents. 39. Bruceantinoside A & B. Novel antileukemic quassinoid glucosides from *Brucea antidysenterica*. *Lloydia* 44 (4) : 470 - 474.
- Osmarov, Z., Ibragimov, A.A. and S. Yu. Yunusov. 1982. A new alkaloid from above ground part of *Nitraria siberica*. *Khim. Prir. Soedin.* (Tashk.) 0 (2) : 225 - 227
- Overnon, L.S., Zaya, J. and E. Trousdale, 1986. Caftaric and coumaric acids in fruits of *Vitis*. *Phytochemistry* 25 (9) : 2127
- Paech, K. and M.V. Tracey, 1965. *Modern Methods of Plant Analysis II*. Springer - Verlag, Berlin.
- Palacios - Chavez, Rodolfo, M., Delaluz Arreguin-San, David L., Garcia, Q. and D. Ramos, 1987. Morphology of pollen grains of the genus *Bursera* (Burseraceae) from the valley of Mexico. *Phytologia* 62 (1) : 58 - 62
- Pandey, Y.N. 1969. Studies on the cuticular characters of some Meliaceae. *Bull. Bot. Surv. India*. 11 : 377 - 386
- Panishin, A.J. 1933 Comparative anatomy of the wood of the Meliaceae, subfamily Swietenioideae. *Amer. J. Bot.* 20 : 638 - 668
- Pardhasarathi, M. and G.S. Sindhu, 1972. Obtusifoliosin, syringetin and dihydrosyringetin from *Soymida febrifuga*. *Phytochemistry* 11 (4) : 1520 - 1522.

- Parvati, A. and L.L. Narayana. 1978. Systematic position of *Balanites* Delile. *Curr. Sci.* 47 (24) : 968 - 970.
- Payne, T.G. and P.R. Jefferies. 1973. Diterpene and flavonoid components of *Dodonea attenuata*. *Tetrahedron* 29 : 2575 - 2583
- Pax. F. 1885 Monographie der Gattung *Acer* : In *Engler's Bot. Jahrb.* 6 : 287 - 374
- Pennington, T.D. and B.T. Styles. 1975. A generic monograph of the Meliaceae. *Blumea* 22 : 419 - 540.
- Pernett, R. 1972. Phytochemie der Burseraceae. *Lloydia* 35 : 280 - 299.
- Polonsky, J., Varon, Z. and B.C. Das, 1976. Triterpenes tetracyclic der *Simarouba amara*. *Phytochemistry* 15 : 337 - 339
- Polonsky, Z., Varon, Z. and C. Moretti. 1980 The antineoplastic quassinoids of *Simaba cuspidata* and *Ailanthus grandis*. *Lloydia* 43 : 503 - 509
- Porter, D.M. 1971. Notes on the floral glands of *Tribulus* (Zygophyllaceae). *Ann. Missouri Bot. Gard.* 58 : 1 - 5.
- Pozzo - Balbi, T., Nobile. T., Scapini, G. and M. Cini. 1978. The triterpenoid acids of *Schinus molle*. *Phyto-*

chemistry 17 : 2107 - 2110

- Puri., V. 1951. Role of floral anatomy in the solution of morphological problems., Bot. Rev. 17 : 272 - 353.
- Puri, V. 1952. Placentation in Angiosperms, Ibid. 18 : 603 - 651
- Purushothaman, K.K., Duraiswamy, K. and J.C. Connolly. 1984. Tetranortriterpenoids from *Melia dubia*. Phytochemistry 23 : 135 - 137
- Purushothaman, K.K., Duraiswamy, K., Connolly, J.D. and D.S. Rycroft. 1985. Triterpenoids from *Walsura piscida*. Phytochemistry. 24 : 2349 - 2354
- Purushothaman, K.K., Sarada., A., Connolly, J.D. and A. Kinniyi. 1979. The structure of roxburghilin, a bis-amide of 2-aminopyrrolidine from leaves of *Aglaiia roxburghiana* (Meliaceae). J. Chem. Soc. Perkin Trans. 1 : 3171 - 3174.
- Purushothaman, K.K., Sarada, A., Sarawati, G. and J.D. Connolly 1977. 5, 7 - Dihydroxy - 6, 2', 4', 5' - tetra methoxy flavone from the leaves of *Chuckrasia tabularis*. Ibid. 16 : 398 - 399.
- Radlkofer, D. 1897. Sapindaceae. In Die Natürlichen Pflanzen Familien (A. Engler and K. Prantl ed) Engelman, Leipzig.
- Radlkofer, D. 1934. Sapindaceae In Das Pflanzenreich (A. Engler ed.) Leipzig 165.

- Radlkofer, D. 1956 Sapindaceae In Das Pflanzenreich 98 : 874 - 880.
- Ralph, I.A. and K.A. Leis. 1979. Pollen Morphology of the Anacardiaceae of northeastern North America. Bull. Torrey. Bot. Club. 106 (2) : 140 - 144
- Rao A.V.R., Bhide, K.S. and R.B. Majumdar. 1980. Phenolics from the bark of *Chloroxylon swietenia* Part-II. Isolation of swietenocoumarins G, H and I. Indian J. Chem. 19B : 1046 - 1048.
- Rao, B.K., Hanumaiah, T., Rao, J.V.M., Rao, K.V.J. and R.H. Thomson. 1984. Two anthraquinones from *Ventilago calyculata*. Phytochemistry 23 (9) : 2104 - 2105.
- Rao, K.V. 1962. Chemical examination of the leaves of *Dodonea viscosa*. J. Indian chem. Soc. 39 (8) : 561 - 562.
- Rao M.M., Gupta, P.S., Singh, P.P. and E.M. Krishna. 1979a. Structure of febrinin-A, a new tetranortriterpenoid from the heartwood of *Soyimida febrifuga*. Indian J. Chem. 17-B : 158 - 159.
- Rao, M.M., Gupta, P.S., Krishna, E.M. and P.P. Singh. 1979b. constituents of Heartwood of *Soyimida febrifuga* : Isolation of flavonoids. Ibid. 17 B : 178 - 180.
- Rao, N.S.P., Row, L.R. and R.T. Brown. 1973. Phenolic constituents of *Semecarpus anacardium*. Phytochemistry 12, 671 - 681.

- Reddy, T.S. and L.L. Narayana. 1986. Chemotaxonomy of Geraniaceae. *J. Indian Bot. Soc.* 65 : 428 - 435.
- Reinhold, L. and Y. Liwschitz. 1970. *Progress in Phytochemistry*. Interscience Publishers, London Vol. 2
- Remillard, S., Rebun, L.I., Howe, G.A. and S.M. Kupchan. 1975. Antimitotic activity of the potent antitumor inhibitor maytansine. *Science* 1989 : 1002 - 1004.
- Rendle, A.B. 1950. *The Classification of Flowering Plants* University Press, Cambridge. Vol. 2.
- Reppel, V.L. 1956. Die coumarine der rosskastanie (*Aesculus hippocastanum* L.). *Planta Med.* 4 : 199 - 203.
- Ridsdule, C.E. 1974. A revision of the family Leeaceae. *Blumea* 22 : 57 - 100.
- Robert, E., Woodson, Jr. Scheny, R.W. and collaborators 1975. Flora of Panama. *Ann. Missouri Bot. Gdn.* 62 (1) : 57.
- Roemer, A., Thomas, H., Kreules, B. and H. Budzikiewicz. 1981. Sesquiterpene ester of type B from *Euonymus europaeus* L. *Z. Naturforsch Teil B. An Org. Chem.* 36 (3) : 379 - 382.
- Row, T.R. and C. Rukmini. 1966a. Chemistry of saponins : Part-1. Saponins of *Sapindus mukorossi* Gaertn. *Indian J. Chem.* 4 : 36 - 38.
- Row, T.R. and C. Rukmini. 1966b. Saponins of *Sapindus emarginatus*. *Ibid* 4 : 149 - 150.

- Sachdev, K. and D.K. Kulshrestha 1983. Flavonoids from *Dodonea viscosa*. *Phytochemistry* 22 : 1253 - 1256.
- Sachdev, K. and D.K. Kulshrestha. 1986. Viscosol, a C-3 prenylated flavonoid from *Dodonea viscosa*. *Ibid* 25 : 1967-1969.
- Saifah, E. and V. Vaisiriraj. 1987. Constituents of the roots of *Cissus rheifolia* Planch. *Lloydia* 50 (2) : 328.
- Saifah, E., Jongbunprasert, V. and C.J. Kelley. 1988. Piriferin, a new pyrrolidine alkaloid from *Aglaia pirifera* leaves. *Ibid* 51 (1) : 80 - 82.
- Saifah, E., Kelley, C.J. and J.D. Leeary. 1983. Constituents of the leaves of *Cissus rheifolia* Planch. *Ibid* 46 (3) : 353 - 358
- Saleh, N.A.M. and M.N. El Hadidi. 1977. An approach to chemosystematics of the Zygophyllaceae. *Biochem. Syst. Ecol.* 5 : 121 - 128
- Saleh, N.A.M., El Hadidi and A.A. Ahmed. 1982. The chemosystematics of Tribulaceae. *Ibid* 10 (4) : 313 - 318.
- Sarma, M.R. 1954. Studies in the family Anacardiaceae. I. Vascular anatomy of the flower of *Mangifera indica* L. *Phytomorphology* 4 : 201 - 208.
- Sati., O.P. and V. Rana. 1987. A new molluscicidal triterpene glycoside from *Aesculus indica*. *Int. J. Crude Drug Res.* 25 (3) : 158 - 160.

- Saunders, E.R. 1939. **Floral Morphology**. 2. Heffer. Cambridge
- Seigler, D.S. and C.S. Butterfield. 1976. The origin of cyanolipids in *Koeleuteria paniculata*. *Phytochemistry* 15 (5) : 842 - 844.
- Seigler, D.S., Cortes, M. and J.M. Aguilera. 1987. Chemical components of guindilla seeds (*Valenzellia trinervis*). *Biochem. Syst. Ecol.* 15 (1) : 71 - 74.
- Seigler, D.A. and D. Kennard. 1977. Biosynthesis of cyanogenic lipid in *Cardiospermum grandiflorum* forma *hirsutum* seeds. *Phytochemistry* 16 : 1825 - 1826.
- Sen, S.P. 1964. Study of active constituents (Keto steroids) of *Cissus quadrangularis*. *Indian J. Pharm* 26 (9) : 247 - 248.
- Shah, A.H., Khan, R.M.A., Maurya, S.K. and V.P. Singh. 1989. Nummularine S : A cyclopeptide alkaloid from stem bark of *Zizyphus nummularia*. *Phytochemistry* 28 (1): 305 - 307.
- Sharma, S.C. and R. Kumar. 1983. Zizynummin - A dammarane saponin from *Zizyphus nummularia*. *Ibid* 22 : 1469 - 1472.
- Sherman, M.M., Borris, R.T., Ogura, M., Cordell, G. and N.R. Farnsworth. 1980. 3S,24S,25-trihydroxytirucall-7-ene from *Ailanthus excelsa*. *Ibid* 19 : 1499 - 1502.
- Shibata, S. 1977. Saponins with biological and pharmacological activity. In *New Natural Products and Plants with Pharmacological, Biological or therapeuti-*

- cal Activity (H. Wagner and P. Wolff ed) Springer-Verlag, Berlin, 191 - 192.
- Shiengthong, D. and A. Ungahakorn, 1979. Constituents medicinal plants. IV. New nitrogenous compounds-odorine and odorinol. *Tetrahedron Lett.* 24 : 2247 - 2250.
- Shimokariyana, M. and T.A. Geissman. 1962. Orientin, a component of *Oxalis cornu Thunb.* In *Chemistry of Natural and Synthetic Coloring Matters* (T. Gore, B.S. Joshi, S.V. Suntharkar and B.D. Tilak. ed.) Academic Press, London. 255 - 259.
- Shukla, R.D. 1955. On the morphology of two abnormal gynoecea in *Peganum harmala* L. *J. Indian Bot. Soc.* 34 : 382 - 387.
- Shukla, S.D., Modi, N.T. and B.S. Deshmankar. 1973. Some pharmacological actions of an alkaloidal fraction from *Cardiospermum halicacabum*. *Indian J. Pharm.* 35 (1) 40-41.
- Sierra, M.G., Mascaretti, O.A., Merkuza, V.M., Tosti. E.L., Ruveda, E.A. and C. Chang. 1974. Peptide alkaloids of *Scutia buxifolia*. *Phytochemistry* 13: 2865 - 2869.
- Singh, R.P. and R. Shiam. 1977. Microsporogenesis in *Litchi chinensis*. *Curr. Sci.* 46 (17) : 613 - 614.
- Singh, S., Garg, H.S. and N.M. Khanna 1976. Dysobinin, a new tetranortriterpene from *Dysoxylum binectariferum*. *Phytochemistry* 15 : 2001 - 2002.

- Singh, B., Agarwal, P.K. and R.S. Thakur. 1982. Aesculoside-A, a new triterpene glycoside from *Aesculus indica*. *Planta Med.* (5) : 409 - 410.
- Singh, B., Agarwal, P.K. and R.S. Thakur. 1987. Aesculoside-B, a new triterpene glycoside from *Aesculus indica*. *Lloydia* 50 (5) : 781 - 783.
- Simon, L. 1975. Le gynécée de *L'Impatiens balsamina* étude morphologique, ontogénique et tératologique. *Canadian J. Bot.* 53 : 2361 - 2384.
- Smith, T.A. 1980. Plant amines. In *Encyclopedia of Plant Physiology*. New Series (A. Pirson and M.H. Zimmerman ed) 8 433 - 460.
- Smith, A.C. and I.W. Bailey. 1941. *Brassiantha*, a new genus of Hippocrateaceae from New Guinea. *J. Arnold Arb.* 22 : 389 - 394.
- Souèges, R. 1953. Embryogénie des Peganaceae-Development de l'embryon chez le *Peganum harmala*. *C.R. Acad. Sci.* 236 : 2185 - 2188
- Souleles, C. and K. Vayas, 1986. Constituents of the leaves of *Aesculus hippocastanum*. *Fitoterapia* 57 (3) : 201-202.
- Srinivas, C.R. et al. 1987. Allergenic agent in contact dermatitis from *Holigarna ferruginea*. *Contact dermatitis* 17 (4) : 219 - 222.
- Srinivasan, S. et al. 1971. Effects of some flavonoids on the surface change characteristics of the vascular system and their antithrombogenic characteristics. *European Conf. Microcircula-*

tion 6 : 394.

- Srivastva, S.K. and V.K. Agnihotri. 1985. New glycoside from the stem bark of *Aphanamixis polystachya*. *Curr. Sci.* 54 (1) : 38 - 40.
- Srivastava, S.K. and J.S. Chauhan. 1977. Chemical examination of the whole plant of *Zizyphus nummularia*. *Planta Med.* 32 (4) : 384 - 387.
- Stankovii, S.K., Bastio, M.B. and J.A. Tarancio. 1985. Composition of the triterpene alcoholic fraction of the horse chestnut (*Aesculus hippocastanum*) seed. *Phytochemistry* 24 (1) : 119 - 122.
- Stephen, G.S. 1980. Cyanogenic compounds and angiosperm phylogeny In. *Encyclopedia of Plant Physiology*. New series. (A. Pirson and M.H. Zimmerman ed.) Vo. 8 : 80 - 116.
- Strappaghetti, G., Cotrsana, S., Craveiro, A. and G. Proiette. 1982. Constituents of essential oil of *Boswellia frereana*. *Phytochemistry* 21 : 2114
- Subash C., Sahai, M. and A.B. Ray. 1986. Chemical constituents of *Rhus Walllichii* Lloydia 49 (3) : 546-
- Subramanian, S.S. and A.G.R. Nair. 1972. Melicitrin - A New myricetin glycoside from the flower of *Melia azadirachta*. *Indian J. chem.* 10; 452
- Subramanian, S.S. and A.G.R. Nair. 1971. Polyphenols of *Lannea coromandelica*. *Phytochemistry* 10 : 1939 - 1940.

- Subramanian, S.S. Joseph, K.J. and A.G.R. Nair. 1968. Polyphenols of *Anacardium occidentale*. *Ibid.*: 8 : 673-674.
- Suradkar, S.S. and N.P. Vaikos, 1988. Nodal anatomy of *Filicium decipiens* Thw. (Sapindaceae). *Indian Bot. Repr.* 7 (1 & 2) : 112 - 113
- Swain, T. 1963. The distribution of alkaloids in the Rutales. In. *Chemical Plant Taxonomy*. Academic Press. London.
- Swain, T. Evolution of flavonoid compounds. In. *The Flavonoids* (J.B. Harborne ed.) Champmann and Hall. London.
- Swingle, W.T. 1938. *J.Wash. Acad. Sci.* 28: - 530.
- Takao, S. 1976. Embryogenic study on *Impatiens texortio*. *Phytomorphology* 26 (2) : 200 - 210.
- Tankashi., Sueyasu T., Nonaka, G. and I. Nishioko. 1984. Tannins and related compounds : 21. Isolation and characterisation of galloyl and -hydroxybenzoyl esters of benzophenone and xanthone (-glycosides from *Mangifera indica*. *Chem. Pharm.Bull.* (Tokyo) 32 (7) : 2676 - 2186.
- Takhtajan, A. 1980. Outline of the classification of flowering Plants (Magnoliophyta) *Bot. Rev.* 46 : 225 - 359.
- Takuo, O., Mori, K., Seno, K. and T. Natano. 1979. Constituents of *Geranium thunbergii* Seib. et Zucc. VII High-performance reversed - phase liquid

chromatography of hydrolysable tannins and related polyphenols. *J. Chromatogr.* 171 (1) : 313 - 320.

Takuo, O., Mori, K. and T. Hatano, 1980. The distribution of Geraniin and Mallotusinic acid in the order Geraniales. *Phytochemistry*. 19 : 547 - 552.

Taylor,, D.A.H. 1988. Biogenesis, distribution and systematic significance of limonoids in the Meliaceae, Cneoraceae and allied taxa. In *Chemistry and Chemical Taxonomy of Rutales*. (P.G. Waterman. and M.F. Grondon. ed.) Academic Press. New York.

Thorne, R.F. 1976. A phylogenetic classification of the Angiospermae. *Evol. Biol.* 9 : 35-106.

Thorne, R.F. 1981. Phytochemistry and angiosperm phylogeny. A summary statement. In *Phytochemistry and Angiosperm Phylogeny* (D.A. Young and S. Seigler ed.) Praeger Publishers, New York. 233 - 295.

Tillson, A.H. and R. Bamford. 1938. The floral anatomy of the Aurantioideae. *Amer. J. Bot.* 25 : 780 - 793.

Tomowa, M.P., Botschewa, D.M., Zaikin, W.G. and N.S.Wulfson. 1977. Steriodal saponins and sapogenins : V. Hecogenin. *Planta Med.* 32 (3) : 223 - 224.

- Tortosa, R.D. 1982. Organography and vascularization of flowers of *Discaria colletia* and *Condalia* (Rhamnaceae). *Kurtziana* 15 (0) : 19 - 40.
- Tripathi, V.D., Agarwal, S.K. and R.P. Rastogi, R.P. 1979. Chemical constituents of *Rhamnus nepalensis*. *Indian J. Chem.* 17B : 89 - 90
- Tschesche, R., Froberg, E. and H.W. Fehlhaber 1968. Alkaloids of Rhamnaceae - *Ceanothus integerrimus* *Tetrahedron Lett.* 11 : 1311 - 1315.
- Tucker, A.O. 1986. Frankincense and myrrh. *Econ. Bot.* 40 (4) : 425 - 433.
- Tulyaganov, T.S., Ibragimov, A.A. and S.Yu. Yunusov. 1981. *Nitraria komarovii* alkaloids : 4. Total synthesis of komarovine and komarovidine. *Khim. Priir. Soedin* (Tasheik) 0 (2) : 192 - 195.
- Turhan, B. 1977. Investigation on the anthracene derivatives in the barks of *Rhamnus oleoides* L. subsp. *gracilis* (Boiss et. Reuti.) Holmbi. *Istanbul Univ. Eozacilik Fak. Mecm.* 13 (1) 1 - 6.
- Tyman, J.H.P. and A.J. Matthews. 1982. Long chain phenols : 22. Compositional studies on Japanese lac (*Rhus Vernicifera*) by chromatography and Mass spectrometry. *J. Chromatogr.* 235 (1) : 149 - 164.
- Uvais, M. and S. Sultanbawa. 1981. Structures of 4 new oxygenated lupanes from *Pleurostylia opposita*. *J. Chem. Soc. Perkin Trans* 10 (10) : 2717 - 2723

- Vahl. J. 1974. *Syst. Bot.*: 3 54.
- Valenta, Z., Gray, A.H., Orr, D.E., Papadopoulos, S. and C. Podesva. 1962. The stereochemistry of quassin. *Tetrahedron* 18 : 1433 - 1441.
- Venkataramaiah, C. and K.N. Rao. 1986. Role of secondary constituents in the systematic position of *Filicium decipiens*. *Thw. Indian J. For.* 8 (4) : 298-300.
- Viswanathan, N.I, 1979., Salapermic acid, a new triterpene acid from *Salacia macrosperma* Wight. *J. chem. Soc. Perkin Trans. 1* (12) : 349 - 352.
- Volsteadt, F.D.R. and D.G. Roux. 1971. Zeyherin, a natural, 3, 8-coumaranoylflavanone from *Phyllogeiton zeyheri* Sond. *Tetrahedron Lett.* 20 : 11647 - 1650.
- Vrkoc, J. and P. Sedmera. 1972. Extractives of *Chloroxylon swietenia* DC. *Phytochemistry* 11 : 2647 - 2648.
- Wagh, A.D., Paknikar, S.K., and S.C. Bhattacharya. 1964. Terpenoids XLV. Structure and absolute configuration of canarone. *J. Org. Chem.* 29 : 2479.
- Wagner, H. and J. Burghat, 1977. Spermidine alkaloids and triterpenes from *Maytenus heterophylla* and *Pleurostylis africana*. *Planta Med.* 32 A : 9 - 14.

- Wagner, H. and J. Burghat. 1981. Macrocyelic spermidine alkaloids from *Pleurostylis africana* (Celastraceae). *Helv. Chim. Acta* 64 (1) : 283 - 296.
- Wang, O. 1983. The dent of the embryo and endosperm in *Zizyphus jujuba*. *Acta Bot. Sin.* 25 (6) : 526 - 531
- Watanabe, I. Saita, A., and K. Takagi . 1973. Pharmacological studies of *Zizyphus* seeds. *Japan J. Pharm* 23 : 563 - 565.
- Waterman P.G. 1983. Phylogenetic implications of the distribution of secondary metabolites within the Rutales. In *Chemistry and Chemical Taxonomy of the Rutales* (P.G. Waterman and M.F. Grondon eds.) A.P. London 377 - 400.
- Waterman, P.G. and M.F. Grondon. 1983. *Chemistry and Chemical Taxonomy of the Rutales*. AP. London.
- Webber, I.E. 1936. Systematic anatomy of the woods of the Simaroubaceae. *Amer. J. Bot.* 23 : 577 - 587.
- Webber, I.E. 1941. systematic anatomy of the woods of the Burseraceae, *Lilloa* 6 : 441 - 465
- Weberbauer, A. 1897. Rhamnaceae, In *Die Natürlichen Pflanzen Familien* (Engler. A. & K. Prantl. ed.)
- Weiffening, J.H. 1966. *Phytochemistry* 5 : 1055
- Wettstein, R. 1935. *Handbuch der Systematischen Botanik*, Frantz. Austria.

- Whitmore, T.C. 1972. *Tree Flora of Malaya*. Vol. Longman, London. 446 - 448.
- Wijeratne, Dharmassree, B.T., Vijay Kumar and M. Uvais 1981. 0-Oxalup-20 - (29) -en-30-ol, a new lupane from *Gymnosporia*. *J. Chem. Soc. Perkin Trans.* 10 (10) 2724 - 2730.
- Willd, H. 1965. Additional evidences from Africa-Madagascar-India Ceylon Land-bridge Theory with special references to the genera *Anisopappus* and *Commiphora*. *Webbia* 19 : 497 - 505.
- Wintersteiner, O., Krakower, G and M. Moore. 1965. Lupenone - and 18 -OC- oleanan- 19- α -01-3-one from *Samadera indica*. *J. Org. Chem.* 30 : 2847 - 2849.
- Wollen Weber, E. and K. Egger, 1970. Methylather des myricitins, quercetin, kaempferol in knospenol von *Aesculus hippocastanum*. *Tetrahedron Lett.* 19 : 1601 - 1604.
- Wolpert-Defilippes, M.K., Adamson, R.H. Cysyk, R.H. and D.G. Johns. 1975. Initial studies on the cytotoxic action of maytansine, a novel ansam-acrolide. *Biochem. Pharmacol.* - 24 : 751 - 754.
- Woo, W.S., Kang, S.S., Shim, S.H., Wagner, H., Chari, V.M., Seligmann, O. and G. Obermeier. 1979. The structure of spinosin (2'' -O- β - glucosylswer-tisin) from *Zizyphus vulgaris* var. *spinosus*. *Phytochemistry* 18 : 353 - 355.

- Woo, W.S., Kang, S.S., Wagner, H., Seligman, O and V.M. Chari 1980. Acylated flavone-C-glycosides from the seeds of *Zizyphus jujuba*. *Ibid* 19 : 2791 - 2793.
- Wüllf, G. and R. Tschesche. 1969. Über triterpenes-XXVI. Über die structure der Rosskastanien saponine (aescin) and die aglykone verwandter glykoside. *Tetrahedron* 25: 415 - 436.
- Yamada, K., Sugiura, K., Shizuri, Y., Wada, H. and Y. Hirata 1977. Isolations and structure of euonymine and neo-euonmine alkaloids from *Euonymus sieboldiana*. *Tetrahedron* 33 (14): 1725 - 1728.
- Yamada, K., Shizuri, Y. and Y. Hirata. 1978. Isolation and new structures of a new alkaloid alatamine and an insecticidal alkaloid wilfordine from *Euonymus alatus forma striatus* (Thunb.) Makino. *Ibid.* 34 (3) : 1915-1920.
- Young, D. 1979. Heartwood flavonoids and the infrageneric relationships of *Rhus* (Anacardiaceae). *Amer.J. Bot.* 66 (5) : 502 - 510.
- Young, A. and Sheilja, J. 1985. Leaf flavonoids of South African *Rhus*. *Ibid* 72 (6) : 916 - 917
- Zanno, P.R., Miura, I., Nakanishi, K. and D.L. Elder. 1975. structure of the insect phagorepellent azadirachtin. Application of PRFT/CWD carbon-13 NMR. *J. Amer. Chem. Soc.* 97 : 1975.

 *