

# LIST OF TABLES

---

- Table 1.** MS stock solutions and amount used for the preparation of 1litre medium (Murashige and Skoog 1962)
- Table 2a.** Macro salts of MS medium
- Table 2b.** Micro salts of MS medium
- Table 2c.** Iron stock solution of MS medium
- Table 2d.** Vitamin stock solution of MS medium
- Table 3.** Effect of individual cytokinins (BA/Kin) on leaf explants of *P oleracea*
- Table 4.** Effect of auxins (IAA/NAA/2,4D) on number of roots and shoots per explant of *P oleracea*
- Table 5.** Effect of BA and Kin on in vitro shoot formation from leaf explants of *P oleracea*
- Table 6.** Effect of BA and IAA on root and callus formation from leaf explant of *P oleracea*
- Table 7.** Effect of BA and NAA on callus, root and shoot induction from leaf explants of *P oleracea*
- Table 8.** Synergistic effect of Kin and IAA on leaf explant of *P oleracea*
- Table 9.** Synergistic effect of Kin and NAA on leaf explants of *P oleracea*
- Table 10.** Effect of IBA and M.S medium strength on number of roots of *P oleracea*
- Table 11.** Effect of individual cytokinin and auxins on leaf explant of *P grandiflora*
- Table 12.** Effect of synergistic concentrations of cytokinin (BA and Kin) in leaf explant of *P grandiflora*
- Table 13.** Effect of synergistic concentrations of BA and auxins on leaf explant of *P grandiflora*
- Table 14.** Effect of synergistic concentrations of Kin and auxins on leaf explant of *P grandiflora*
- Table 15.** Effect of cytokinins on nodal explants of *P grandiflora*
- Table 16.** Effect of Boron, CaCl<sub>2</sub> and Ca- gluconate on number of shoots per explants of *P grandiflora*
- Table 17.** Content of total fats (mg/g) in different samples of *P oleracea*
- Table 18.** Relative content of fatty acids present in different samples of *P oleracea*
- Table 19.** Effect of sodium ascorbate in different samples of *P grandiflora*

**Table 20.** Effect of different doses of EMS on in vitro leaf explants of *P oleracea*

**Table 21.** Relative content of fatty acids on EMS treated in vitro cultures of *P oleracea*

**Table 22.** Fatty acid profiling of variant in vitro shoot cultures at 0.4 and 0.6% EMS of *P oleracea*