

# LIST OF FIGURES

---

- Fig.1.** Percentage of global herbal market share
- Fig.2.** General habit of *P oleracea* Linn.
- Fig.3.** General habit of *P grandiflora* Hook.
- Fig.4.** Structural components of betalain
- Fig.5.** Effect of BA on leaf explants of *P oleracea*
- Fig.6.** Effect of Kin on leaf explants of *P oleracea*
- Fig.7.** Effect of auxins on leaf explant of *P oleracea*
- Fig.8.** Effect of 2.5  $\mu$ M BA with different concentrations of Kin on leaf explants of *P oleracea*
- Fig.9.** Effect of 5  $\mu$ M BA with different concentrations of Kin on leaf explants of *P oleracea*
- Fig.10.** Effect of 10  $\mu$ M BA with different concentrations of Kin on leaf explants of *P oleracea*
- Fig.11.** Effect of 15-20  $\mu$ M BA with different concentrations of Kin on leaf explants of *P oleracea*
- Fig.12.** Effect of BA and IAA on leaf explants of *P oleracea* after 8 weeks
- Fig.13.** Effect of BA and NAA on leaf explants of *P oleracea* after 8 weeks
- Fig.14.** Effect of BA and 2,4D on leaf explants of *P oleracea* after 8 weeks
- Fig.15.** Effect of BA and 2,4D on biomass of callus (gm) from leaf explants of *P oleracea*
- Fig.16.** Callus growth index (gm) on an optimized media of 2.5 $\mu$ M BA and 2.5  $\mu$ M 2,4D
- Fig.17.** Effect of Kin and IAA on leaf explants of *P oleracea* after 8 weeks
- Fig.18.** Effect of Kin and NAA on leaf explants of *P oleracea* after 8 weeks
- Fig.19.** Effect of Kin and 2,4D on leaf explants of *P oleracea* after 8 weeks
- Fig.20.** Effect of Kin and 2,4 D on biomass of callus (gm) from leaf explants of *P oleracea*
- Fig.21.** Effect of coconut water on in vitro shoot elongation of *P oleracea*
- Fig.22.** Effect of coconut water on shoot number in *P oleracea*
- Fig.23.** Histological changes during in vitro shoot regeneration of *P oleracea* from leaf explants

- Fig.24.** Regeneration of *P. oleracea* from in vitro leaf explant on optimized medium (5  $\mu$ M BA and 10  $\mu$ M Kin)
- Fig.25.** Number of shoots from in vitro leaf explants of *P. oleracea* after each subculture
- Fig.26.** Effect of media strength and IBA on rooting potential of *P. oleracea*
- Fig.27.** Effect of individual auxin on leaf explant of *P. grandiflora* after 8 weeks
- Fig.28.** Synergistic effect of BA and Kin on leaf explant of *P. grandiflora* after 8 weeks
- Fig.29.** Effect of BA and IAA on leaf explants of *P. grandiflora* after 8 weeks
- Fig.30.** In vitro shoot regeneration from young leaf explant in *P. grandiflora*
- Fig.31.** Effect of BA and NAA on leaf explants of *P. grandiflora* after 8 weeks
- Fig.32.** Effect of BA and 2,4D on leaf explants of *P. grandiflora* after 8 weeks
- Fig.33.** Effect of Kin and IAA on leaf explants of *P. grandiflora* after 8 weeks
- Fig.34.** Effect of Kin and NAA on leaf explants of *P. grandiflora* after 8 weeks
- Fig.35.** Effect of Kin and 2,4D on leaf explants of *P. grandiflora* after 8 weeks
- Fig.36.** Shoot buds formed at 2.5  $\mu$ M BA and 2.5  $\mu$ M NAA transferred to different media for bud break in *P. grandiflora* after 2 weeks
- Fig.37.** Shoot elongation at 5  $\mu$ M GA<sub>3</sub> in *P. grandiflora* after 12 weeks
- Fig.38.** Effect of GA<sub>3</sub> on the number of shoots in *P. grandiflora* after 2 weeks
- Fig.39.** Histological changes during in vitro shoot regeneration of *P. grandiflora* from leaf explants
- Fig.40.** Regeneration from in vitro leaf explants of *P. grandiflora*
- Fig.41.** Multiple shoot formation in *P. grandiflora* from nodal explants after 8 weeks
- Fig.42.** Shoot tip necrosis in shoot cultures of *P. grandiflora*
- Fig.43.**
- a. Effect of CaCl<sub>2</sub> and Ca-gluconate on % necrosis in shoot cultures of *P. grandiflora* after 4 weeks
  - b. Effect of Boron on % necrosis in shoot cultures of *P. grandiflora* after 4 weeks
- Fig.44.** Oil yielded from the whole plant of *P. oleracea* after soxhlet extraction
- Fig.45.** GC analysis of standard fatty acid mixture
- Fig.46.** GC analysis for the fatty acid profiling of in vivo shoot of *P. oleracea*
- Fig.47.** GC analysis for the fatty acid profiling of in vivo leaf of *P. oleracea*
- Fig.48.** GC analysis for the fatty acid profiling of stem of *P. oleracea*
- Fig.49.** GC analysis for the fatty acid profiling of in vitro shoots of *P. oleracea*
- Fig.50.** GC analysis for the fatty acid profiling of callus culture of *P. oleracea*
- Fig.51.** Total Ion Chromatogram in EI mode of in vivo shoots of *P. oleracea*

- Fig.52.** Total Ion Chromatogram in EI mode of in vivo leaves of *P oleracea*
- Fig.53.** Total Ion Chromatogram in EI mode of stem of *P oleracea*
- Fig.54.** Total Ion Chromatogram in EI mode of in vitro shoot of *P oleracea*
- Fig.55.** Total Ion Chromatogram in EI mode of callus cultures of *P oleracea*
- Fig.56.** Extraction of Betalains
- Fig.57.** Betalain content in whole plant of *P grandiflora* in presence of 50 mM sodium ascorbate
- Fig.58.** Betalain content in stem of *P grandiflora* with 50 mM sodium ascorbate
- Fig.59.** Spectrophotometric analysis of betalain content of stem of *P grandiflora* with and without Sodium ascorbate after 24 hours
- Fig.60.** Linear regression plot of the probit value against the log (dose) to determine LD50 dose in *P oleracea* leaf explant treated with EMS(0.1-1%)
- Fig.61.** Effect of different concentrations of EMS after 12 weeks
- Fig.62.** The variations expressed in different doses of in vitro leaf explants cultured on 5µM BA and 10µM Kin
- Fig.63.** Callus induction in EMS treated in vitro leaf explants
- Fig.64.** Effect of different doses of EMS on callus induction from in vitro leaf explants of *P oleracea* after 3 months
- Fig.65.** Fatty acid profiling of 0.1% EMS treated shoots of *P oleracea*
- Fig.66.** Fatty acid profiling of 0.2% EMS treated shoots of *P oleracea*
- Fig.67.** Fatty acid profiling of 0.3% EMS treated shoots of *P oleracea*
- Fig.68.** Fatty acid profiling of 0.1% EMS treated callus of *P oleracea*
- Fig.69.** Fatty acid profiling of 0.2% EMS treated callus of *P oleracea*
- Fig.70.** Fatty acid profiling of 0.3% EMS treated callus of *P oleracea*
- Fig.71.** Fatty acid profiling of 0.4% EMS treated shoot cultures of *P oleracea*
- Fig.72.** Fatty acid profiling of 0.6% EMS treated shoot cultures of *P oleracea*
- Fig.73.** Total Ion Chromatogram of fatty acids present in 0.1% EMS treated shoots of *P oleracea*
- Fig.74.** Total Ion Chromatogram of fatty acids present in 0.2% EMS treated shoots of *P oleracea*.
- Fig.75.** Total Ion Chromatogram of fatty acids present in 0.3% EMS treated shoots of *P oleracea*.
- Fig.76.** Total Ion Chromatogram of fatty acids present in 0.1% EMS treated callus of *P oleracea*.

- Fig.77.** Total Ion Chromatogram of fatty acids present in 0.2% EMS treated callus of *P oleracea*
- Fig.78.** Total Ion Chromatogram of fatty acids present in 0.3% EMS treated callus of *P oleracea*
- Fig.79.** Total Ion Chromatogram of fatty acids present in 0.4% EMS treated shoots of *P oleracea*
- Fig.80.** Total Ion Chromatogram of fatty acids present in 0.6% EMS treated shoots of *P oleracea*
- Fig.81.** Mass Spectra of Linoleic Acid (9,12 Octadecadienoic Acid, methyl ester) in 0.1% EMS shoot cultures of *P oleracea*
- Fig.82.** Mass Spectra of Linolenic Acid (9,12,15 Octadecatrienoic Acid, methyl ester) in 0.1% EMS shoot cultures of *P oleracea*
- Fig.83.** Effect of different concentrations of EMS on leaf explants of *P oleracea*