

LIST OF FIGURES

Figure No.	Title	Page No.
1	Distribution of medicinal plants by habits (GOI 2000)	14
2	The proportion of different parts of medicinal plants used (GOI 2000)	14
3	Constituents of Dashmoola	24
4	<i>Oroxylum indicum</i>	28
5	<i>Stereospermum suaveolens</i>	31
6	<i>O.indicum</i> and <i>S.suaveolens</i> plants	56
7	Explants of <i>O.indicum</i> and <i>S.suaveolens</i>	66
8	Methodology for regeneration studies in <i>O.indicum</i> and <i>S.suaveolens</i>	67
9	Germination of <i>O.indicum</i> seeds in different substrates	79
10	Seed germination in <i>O.indicum</i>	80
11	Effect of different substrates on length of <i>O.indicum</i> seedlings	81
12	Effect of different substrates on collar diameter of <i>O.indicum</i> seedlings	82
13	Effect of different substrates on biomass of <i>O. indicum</i> seedlings	83
14	Morphogenic response in different explants of <i>O.indicum</i> in presence of individual cytokinins after 4 weeks	86
15	Effect of individual cytokinins in inducing multiples from <i>in vitro</i> nodes of <i>O.indicum</i> after 8 weeks	89
16	Shoot induction from cotyledonary node explants of <i>O.indicum</i> in presence of two cytokinins after 4 weeks	92
17	Shoot induction from nodal explants of <i>O.indicum</i> in presence of individual cytokinins after 4 weeks	100
18	Shoot induction from nodal explants of <i>O.indicum</i> in presence of two cytokinins after 4 weeks	104
19	Effect of two cytokinins in inducing multiples from <i>in vitro</i> nodes of <i>O.indicum</i> after 8 weeks	106
20	Multiple shoots developed from <i>in vitro</i> nodes of <i>O.indicum</i> in presence of two cytokinins	107
21	Number of shoots from <i>in vitro</i> nodes of <i>O.indicum</i> in MS and WPM medium after 12 and 16 weeks	108
22	Average number of shoot and shoot length of <i>O.indicum</i> in MS+ BAP(8 μ M) + TDZ(0.25 μ M) combination after 24 and 32 weeks	109
23	Shoot induction from nodal explants of <i>O.indicum</i> in presence of cytokinins and auxins after 4 weeks	111
24	Multiples developed from <i>in vitro</i> nodes of <i>O.indicum</i> in presence of cytokinins and auxins	115
25	Effect of cytokinins and auxins in inducing multiples from <i>in vitro</i> nodes of <i>O.indicum</i> after 8 weeks	116
26	Effect of MS+BAP (20 μ M)+IBA(0.5 μ M) and WPM+BAP(16 μ M) +IAA(1 μ M)on inducing multiples in <i>O.indicum</i> after 12 and 16 weeks	117
27	Average shoot number and shoot length of <i>O.indicum</i> in MS+BAP(20 μ M)+IBA(0.5 μ M) combination	117
28	Shoot differentiation from morphogenic basal callus in <i>O.indicum</i>	120

29	Multiple shoots formation from morphogenic callus in MS and WPM medium fortified with different PGRs after 12, 16, 24 weeks in <i>O.indicum</i>	121
30	Average number of shoots from morphogenic callus in MS+BAP (20µM)+NAA(1µM) after 8,12,16,24 weeks in <i>O.indicum</i>	122
31	Shoot differentiation in <i>O.indicum</i> from morphogenic callus in MS medium fortified with BAP (20µM) + NAA (1µM)	123
32	Indirect organogenesis from nodal explants of <i>O.indicum</i> in MS medium in following combinations	125
33	<i>In vitro</i> root induction in <i>O.indicum</i> after 4 weeks	129
34	Hardening of <i>O.indicum</i> plants in cocopeat substrate under lab conditions	132
35	Hardening of <i>O.indicum</i> plants in Cocopeat substrate	134
36	Percent survival in different substrates after four weeks of transfer of <i>O.indicum</i> plants to greenhouse	135
37	Leaf number and leaf length of <i>O.indicum</i> plants in different substrates after 4 weeks	136
38	Initial and final shoot, root length and plant height in different substrates in <i>O.indicum</i>	136
39	Hardening of <i>O.indicum</i> plants in Sand substrate	137
40	Growth in <i>O.indicum</i> plants in Sand, Soil and Sand:Soil substrate after 8 weeks	138
41	Plant height and girth in <i>O.indicum</i> (four) plants after transfer to garden	139
42	Tissue cultures plants of <i>O.indicum</i> grown in botanical garden	140
43	Hardening of <i>O.indicum</i> plants in Soil substrate	142
44	Hardening of <i>O.indicum</i> plants in Cocopeat:Sand substrate	143
45	Hardening of <i>O.indicum</i> plants in Sand:Soil substrate	144
46	Hardening of <i>O.indicum</i> plants in Cocopeat:Sand:Soil substrate	146
47	Effect of sodium alginate and calcium chloride on synseed formation	149
48	Germination of <i>in vitro</i> nodes encapsulated in half and full WPM matrix in different regenerative medium in <i>O.indicum</i> after 4 weeks	152
49	Germination of <i>in vitro</i> nodes encapsulated in half and full WPM+BAP (16µM) matrix in different regenerative medium in <i>O.indicum</i> after 4 weeks	157
50	Germination of synseeds encapsulated in half and full strength WPM+BAP(16µM)+IBA(0.1µM) matrix in different regenerative media in <i>O.indicum</i> after 4 weeks	159
51	Effect of agar substrate in ½WPM+BAP(16µM)+IBA(0.1µM) matrix and ½ WPM+NAA (5 µM) regenerative medium in <i>O.indicum</i> after 4 weeks	161
52	Germination of <i>S.suaveolens</i> seeds in different substrates	162
53	Seed germination in <i>S.suaveolens</i>	164
54	Germination rate (GR) of <i>S.suaveolens</i> seeds in different planting substrates	165
55	Mean daily germination (MDG) of <i>S.suaveolens</i> seeds in different planting substrates	165
56	Germination index (GI) of <i>S.suaveolens</i> seeds in different planting substrates	166
57	Morphogenic response of cotyledonary leaf and hypocotyl explants of <i>S.suaveolens</i> in presence of individual cytokinins after 4 weeks	169
58	Shoot formation from cotyledonary node explants of <i>S.suaveolens</i> in presence of individual cytokinins after 4 weeks	172

59	Effect of individual cytokinins in inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i> after 8 weeks	173
60	Development of shoots from <i>in vitro</i> nodes of <i>S.suaveolens</i> in MS and WPM medium	174
61	Effect of individual cytokinins in inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i> after 12 weeks	176
62	Shoot formation from cotyledonary node explants of <i>S.suaveolens</i> in presence of two cytokinins after 4 weeks	179
63	Effect of two cytokinins from <i>in vitro</i> nodes of <i>S.suaveolens</i> in inducing multiples after 8 weeks	183
64	Shoot induction from cotyledonary node explants of <i>S.suaveolens</i> in presence of cytokinins and auxins after 4 weeks	185
65	Shoot induction from nodal explants of <i>S.suaveolens</i> in presence of individual cytokinins after 4 weeks	188
66	Effect of individual cytokinins in inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i> after 8 weeks	191
67	Effect of individual cytokinins on inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i>	192
68	Effect of individual cytokinins in inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i> after 12 weeks	193
69	Effect of individual cytokinins from <i>in vitro</i> nodes of <i>S.suaveolens</i> on shoot number and length after 16, 24 and 32week	194
70	Effect of MS+TDZ (0.1 μ M), MS+TDZ (0.2 μ M) and WPM+Kn(8 μ M) on shoot number and shoot length of <i>S.suaveolens</i> in different passages	195
71	Effect of additives on shoot number and length of <i>S.suaveolens</i> after 32 weeks	196
72	Development of healthy shoots of <i>S.suaveolens</i> in MS and WPM medium supplemented with additives	197
73	Shoot induction from nodal explants of <i>S.suaveolens</i> in MS and WPM medium fortified with two cyokinins after 4 weeks	200
74	Synergistic effect of cytokinins on inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i> after 8 weeks	203
75	Development of healthy shoot from <i>in vitro</i> nodes of <i>S.suaveolens</i> after 8 weeks	204
76	Effect of two cytokinins on inducing multiples from <i>in vitro</i> nodes of <i>S.suaveolens</i> at different passages	205
77	Enhancement of <i>S.suaveolens</i> shoots from <i>in vitro</i> nodes of in combinations of two cytokinins : MS+ BAP(4 μ M) +Kn(8 μ M), MS+ BAP(8 μ M) +TDZ(0.1 μ M)	206
78	Multiple shoot formation from <i>in vitro</i> nodes of <i>S.suaveolens</i> in combinations of two cytokinins : MS+BAP(8 μ M)+TDZ(0.2 μ M), MS+Kn(8 μ M) +TDZ(0.2 μ M)	207
79	Effect of MS medium fortified with Kn (8 μ M) with TDZ(0.2 μ M) on shoot length of <i>S.suaveolens</i> at each passage	208
80	Indirect shoot formation in WPM medium fortified with individual cytokinin in <i>S.suaveolens</i>	212
81	Effect of WPM medium fortified with coconut water (10%) on shoot number and length of <i>S.suaveolens</i> at three passages	213

82	Indirect organogenesis in WPM medium fortified with Kn (8 μ M) with IAA (0.1 μ M, 1 μ M), Kn (8 μ M) with NAA (0.1 μ M) after 12 weeks in <i>S.suaveolens</i>	214
83	Indirect organogenesis in presence of cytokinin and auxin in <i>S.suaveolens</i>	215
84	Root induction of <i>S.suaveolens</i> shoots in MS and WPM medium after 4 weeks	218
85	Hardening of <i>S.suaveolens</i> plants in cocopeat under culture room conditions	221
86	Hardening of <i>S.suaveolens</i> plants in Cocopeat substrate	223
87	Percent survival of <i>S.suaveolens</i> after four weeks in different substrates under greenhouse conditions	224
88	Leaf number and leaf length of <i>S.suaveolens</i> in different substrates after 4 weeks	224
89	Initial and final shoot, root length and plant height of <i>S.suaveolens</i> in different substrates	225
90	Hardening of <i>S.suaveolens</i> plants in Cocopeat: Sand substrate	226
91	Hardening of <i>S.suaveolens</i> plants in Sand:Soil substrate	227
92	Hardening of <i>S.suaveolens</i> plants in Cocopeat:Soil substrate	229
93	Hardening of <i>S.suaveolens</i> plants in Cocopeat: Sand: Soil substrate	230
94	Effect of MS basal medium matrix on <i>S.suaveolens</i> synseed germination after 4 weeks	234
95	Effect of ½ MS+Kn(8 μ M) matrix on <i>S.suaveolens</i> synseed germination after 4 weeks	240
96	Effect of full strength MS+TDZ (0.2 μ M) matrix on <i>S.suaveolens</i> synseed germination after 4 weeks	245
97	Effect of agar substrate in MS + Kn (8 μ M) matrix and MS+NAA (2 μ M) regenerative medium in <i>S.suaveolens</i>	249
98	Effect of agar substrate on <i>S.suaveolens</i> synseed germination after 4 weeks	250