



List of Figures

| Figure no. | Page no. |
|---|----------|
| 1.1 A schematic representation of identified current branches during the Southwest monsoon..... | 5 |
| 1.2 Schematic representation of surface circulation in the Indian Ocean during the NE monsoon..... | 6 |
| 2.1 Locations of the sediment cores employed in this study along with the Arabian Sea bottom topography..... | 20 |
| 2.2 Schematic diagram of CO ₂ preparation using CAPS and the Mass Spectrometer inlet system..... | 38 |
| 3.1 Locations of the core SK145-9 and other cores with which it has been compared..... | 46 |
| 3.2 September Salinity contours in the eastern Arabian Sea..... | 47 |
| 3.2 Calibrated radiocarbon ages and sedimentation rates (cm/ka) for various intervals in the core SK145-9..... | 48 |
| 3.4 Downcore variations of $\delta^{18}\text{O}$ of three species of foraminifera in the core SK 145-9 shown along with % <i>G.bulloides</i> data from the ODP Site 723A..... | 51 |
| 3.5 Downcore variation of CaCO ₃ , C _{org} and C/N ratio in the core SK145-9 covering the last ~2800 years..... | 55 |
| 3.6 Downcore variation of $\delta^{13}\text{C}$ of the three species of foraminifera in the core SK 145-9..... | 58 |
| 3.7 Downcore variation of $\delta^{15}\text{N}$ of sedimentary organic matter in the core SK145-9..... | 60 |
| 3.8 Comparison of the TSI data and precipitation proxies for the past 1200 years..... | 63 |
| 3.9 Power spectra of oxygen isotope timeseries in 3 species of foraminifera..... | 64 |

| Figure no. | | Page no. |
|------------|---|----------|
| 4.1 | Locations of the core SS3827 G and other cores with which it has been compared..... | 69 |
| 4.2 | The inflow of low salinity water from the Bay of Bengal into the Arabian Sea along the NMC in the month of December.. | 71 |
| 4.3 | Calibrated Radiocarbon ages and sedimentation rates for the core SS 3827G..... | 72 |
| 4.4 | Temporal variations in the oxygen isotope ratios ($\delta^{18}\text{O}$) of three species of foraminifera in the core SS3827 G..... | 75 |
| 4.5 | Comparison of the $\delta^{18}\text{O}$ record of the core SS3827 G with the GISP2 $\delta^{18}\text{O}$ record..... | 78 |
| 4.6 | Downcore variation of CaCO_3 (%) in the core SS3827 G for the past 35 ka..... | 84 |
| 4.7 | Comparative study of the core SS3827 G with other cores from eastern Arabian Sea viz. SS3104 G and SS3101 G..... | 87 |
| 4.8 | Downcore variations of C_{org} plotted along with CaCO_3 (%) and C/N ratio..... | 89 |
| 4.9 | Downcore variation of $\delta^{15}\text{N}$ in the core SS3827 G for the past 35 ka..... | 90 |
| 4.10 | $\delta^{15}\text{N}$ vs. Nitrogen content for all the three cores viz. SS3827 G, SS4018 G and SK 145-9 analyzed in this study..... | 91 |
| 4.11 | Downcore variation of $\delta^{13}\text{C}$ values of the three species of planktonic foraminifera in the core SS3827 G..... | 92 |
| 4.12 | Power spectra for various paleoclimatic proxies of the core SS3827 G..... | 94 |
| 5.1 | Locations of core SS 4018G and other cores with which it has been compared..... | 100 |
| 5.2 | SST contours (in $^{\circ}\text{C}$) during August..... | 102 |
| 5.3 | Calibrated radiocarbon ages and sedimentation rates for SS4018G.... | 104 |

| Figure no. | | Page no. |
|------------|--|----------|
| 5.4 | Downcore variations of productivity proxies in the core SS 4018 G..... | 106 |
| 5.5 | Downcore variation of $\delta^{13}\text{C}$ of three species of foraminifera..... | 109 |
| 5.6 | Plot showing the downcore variation of $\delta^{15}\text{N}$ of the sedimentary organic matter in the core SS4018 G..... | 110 |
| 5.7 | Downcore variation in $\delta^{18}\text{O}$ of three species of foraminifera in the core SS4018 G..... | 114 |
| 5.8 | Power spectra for various paleoclimatic proxies of the core SS4018 G..... | 119 |