

List of Figures

Figure 1.1: Maize field in the region of Chhani (North of Vadodara). The most preferred crop of <i>Spodoptera frugiperda</i>	3
Figure 1.2: Maize field in the region of Savli (North of Vadodara).....	3
Figure 1.3: Castor field in the region of Padra (Southwest of Vadodara).....	4
Figure 1.4: Cotton field in the region of Dabhoi (South east of Vadodara).....	4
Figure 1.5: Cauliflower field in the region of Sherkhi (West of Vadodara).....	5
Figure 1.6: Banana field in the region of Waghodia (East of Vadodara).....	5
Figure 1.7: FAW in News.....	16
Figure 1.8: FAW in News (2).....	17
Figure 1.9: Diamide family (Chlorantraniliprole)	23
Figure 1.10: Avermectin family (Emamectin benzoate).....	23
Figure 2.1: Tree depicting the relationship between the GST classes present in insects.....	38
Figure 2.2: Venn diagram depicting the fat bodies and midguts differentially expressed genes in the of <i>S. litura</i> against treatment with tomatine.....	39
Figure 3.1: Surveying in the fields of Savli and Waghodia.....	44
Figure 3.2: Cauliflower and Cotton fields in Waghodia.....	45
Figure 3.3: Maize field in the Channi and Sherkhi regions of Vadodara.....	46
Figure 3.4: Collection of Fall Armyworm from Agricultural Fields.....	47
Figure 3.5: Maize-producing states in India (ICAR).....	49
Figure 3.6: Larvae Reared on Natural and Artificial Diets.....	52
Figure 3.7: Insecticides found in an agricultural field (Cypermethrin + Chlorpyrifos).....	58
Figure 3.8: Commercial grade Emamectin benzoate.....	60
Figure 3.9: Technical grade product and structural formula of Chlorantraniliprole.....	62
Figure 3.10: Technical details of Emamectin benzoate.....	64

Figure 3.11: Stock solutions and preparing different concentrations of insecticide solutions.....	66
Figure 3.12 A-F: Diet preparation process (weighing ingredients, boiling Part A, mixing parts A and B, filling in bottles, pouring in trays, releasing insects).....	67
Figure 3.13: Flow of histological studies.....	68
Figure 3.14: Flow of molecular work.....	69
Figure 3.15: Flow of transcriptome analysis.....	73
Figure 4.1: <i>S. frugiperda</i> infestation in maize fields (creating windows), Damage by FAW in Maize (complete holes), Excreta deposition and FAW on maize.....	77
Figure 4.2: Life cycle of Fall Armyworm (<i>Spodoptera frugiperda</i>).....	78
Figure 4.3: Egg mass with scales of <i>Spodoptera frugiperda</i> S. (4X).....	79
Figure 4.4: Egg mass of <i>Spodoptera frugiperda</i> S (4X).....	80
Figure 4.5: Magnified neonate/first instar (4X) and fully grown caterpillar (normal view....	81
Figure 4.6: Pupa and Adult of <i>S. frugiperda</i> (male).....	82
Figure 4.7: Histological slides.....	106