LIST OF TABLES

Table	No.	Page	No
1.	Nutritional demands of the growing fetus and the nursing infant in relation to maternal diet (Rajalakshmi, 1979).	4	
2.	Intakes of critical nutrients as compared to the nutritional requirements of pregnancy and lactation.	13	
3.	Classes of factors in behavioral deve- lopment.	28	,
4.	Critical phases in the developing brain	31	`
5.	The nutritional stress of reproduction in rat and man.	32	
6.	Variability in conversion ratios of Carotene to Vitamin A.	67	
7.	Composition of the stock diet (Expt. 1).	76	
8.	Food consumption pattern in low and high income groups in Baroda as compared to values reported for the British	77 0	
•	(Experiment 2).	79	
9.	Pattern of vegetable consumption in different groups (Experiment 2).	81	
10.	Vitamin content of diets consumed by different groups (Experiment 2).	84	
11.	Composition of diets containing different levels of protein (Experiment 3).	86	
12.	Composition of wheat and wheat + lysine diets (Experiment 4).	88	

lable No.		Page No
13.	Composition of diets providing different levels of calcium (Experiment 5).	90
14.	Composition of selt mixtures providing different amounts of calcium (Expt. 5).	91
15.	Composition of diet providing vitamin A or carotene (Experiment 6).	93
16.	Summary of various experiments performed.	94
17.	Composition of vitamin mixture.	98
18.	Composition of salt mixture.	99
19.	The sources of the chemicals used.	100
20.	Details of reagents and standards used.	113
21 <u>.</u> ę	Growth rate and efficiency of food utilization in male rats subjected to different degrees of postweaning food restriction in relation to preweaning nutritional status.	122
22 _B	Growth and efficiency of food utilization in female rats subjected to different degrees of postweaning food restriction in relation to preweaning nutritional status \hat{V} .	126
23.	Growth rate and efficiency of food utili- zation in response to dietary rehabilita- tion in male rats subjected to different degrees of preweaning and/or postweaning food restriction.	130•
24•	Growth rate and efficiency of food utilization in response to dietary rehabilitation, in female rats subjected to different degrees of preweaning and/or postweaning food restriction.	134

-

Table No.		Page No.
25.	Adult weights of rats subjected to different degrees of preweaning and/or postweaning food restruction and subsequent rehabilitation.	142
26.	Activity in the open field after rehabi- litation of male rats subjected to differ- ent degrees of preweaning and/or post- weaning food restriction.	145
27.	Voluntary activity of undernourished animals as reported by various investigators.	147
28.	Behavioral measures in an activity wheel of male rats subjected to different degrees of preweaning and/or postweaning food restriction.	150
29•	Reproductive performance after rehabilita- tion of female rats subjected to different degrees of preweating and/or postweating food restriction	155
30.	Maternal behaviour of rats subjected to different degrees of preweaning and/or postweaning food restriction.	168
31.	Reproductive performance of rats fed diets consumed by different population groups.	177
32.	Changes in blood-hemoglobin during gesta- tion and lactation in rats fed diets	•
<i>3</i> 3•	consumed by different population groups. Changes in serum protein during gestation and lactation in rats fed diets consumed by	181
	different population groups.	183

Table No.		Page No
34.	Longitudinal data on blood-hemoglobin and serum protein during gestation and lactation in females fed LIG, HIG and W diets.	186
35	Food intake and utilization during gesta- tion and lactation in rats fed diets consumed by different population groups.	191
36	Nitrogen retention during gestation and lactation in rats fed diets consumed by	
37	Apparent digestibility, per cent absorbed N retained, weight gain per g N and estimate of per cent protein in tissue gained during gestation and lactation in rats fed diets consumed by different population groups.	193 195
38	Values during gestation and lactation in rats fed diets consumed by different population groups expressed as per cent of pregestation (initial) values.	197
39	Nitrogen balance at different stages of the reproductive cycle of females fed LIG, HIG and W diets.	198
40	Nitrogen balance data for non-pregnant, non-lactating animals of generation 1 fed diets consumed by different population groups at points corresponding to the stages specified.	297
41	Comparison of a nitrogen balance in generations 0 and 1 during gestation and lactation of rats fed diets consumed by different	
	population groups.	209

Table	No.	Page No.
42.		216
43.	Growth in phase II (11-20 weeks) of animals fed different levels of protein.	217
₹ \ \$	Growth in Phase II(20-23 weeks) of animals fed different levels of protein.	218
45,	Growth in phase IV (23-31 weeks) of animals feed different levels of protein.	219•
46,	Effect of feeding different levels of protein on blood-hemoglobin and serum protei	n 225
47.	Longitudinal data on blood-hemoglobin and serum protein changes of animals fed different levels of protein.	226,
48.	Food, fecal and urine nitrogen of animals fed different levels of protein.	230
49,	Apparent digestibility, mg N retained and per cent absorbed N retained in animals fed different levels of protein.	231
50.	reference values for animals fed different	
51	levels of protein. Incorporation of C ¹⁴ -DL-leucine into serum protein in rats fed different levels of	232
52₀	protein. I. at the end of Phase-I. Incorporation of C ¹⁴ -DL-leucine into serum protein in rats fed different levels of protein-values expressed as percent 12 hr.	234
	values. I. At the end of phase-I.	235

able No	•	Page No
53 .	Incorporation of C ¹⁴ -DL-Leucine into serum protein in rats fed different levels of protein- phase I and phase II values compared.	236
54.	Incorporation of C ¹⁴ -DL-leucine into serum- protein in rats fed different levels of protein-values expressed as per cent control values.	237 ,
55•	Growth of rats fed wheat diet with and without addition of lysine at different ages.	252
56.	Nutritional status as judged by blood- hemoglobin, serum protein, liver/weight and liver protein of rats fed a wheat diet with and without addition of lysine.	256
<i>5</i> 7.	Values for lysine supplemented group as per cent of wheat group - values for data presented in Table 52 and 53.	257
58.	Incorporation of C ¹⁴ -DL-leucine into serum protein in rats fed on wheat diets with or without addition of lysine at different ages.	259
5 9•	Body weight of rats fed different amounts of dietary calcium at different ages.	267
60.	Body weight gains of rats fed different levels of calcium at different ages.	268
61.	Values for low calcium and high calcium diets as per cent of the group fed 440 mg Ca per 100 g diet for data presented in Tables 59, 60 and 62.	2 ′69ે

•

Table	No.	Page No
. 62,	Blood hemoglobin and serum protein of rats fed differentlevels of calcium at different ages.	272
63 •	Composition of the femur of rats fed different levels of calcium at different ages.	273
64.	Composition of femur at different ages in rats fed a standard diet providing 440 g Ca throughout.	1 279
65.	Body weight gain during period of depletion and repletion in rats fed vitamin A in carotene prior to depletion.	2 90
66.	Serum vitamin A during period of depletion and repletion in rats fed vitamin A or carotene prior to depletion.	292