

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE NO.
3.1	Picture of (a) <i>Aconitum ferox</i> plant, (b)Dried <i>Aconitum</i> root.	30
3.2	Structure of Aconitine.	36
3.3	Pictures of (a) Plant parts, (b) Plant (c) Roots of <i>Solanum indicum</i>.	41
3.4	Structures of glycoalkaloids of <i>Solanum indicum</i>.	46
3.5	Chemical Structure of Piperine.	51
3.6	Pictures of (a) <i>Piper longum</i> climber and (b) <i>Piper longum</i> Fruit.	56
3.7	Pictures of <i>Piper nigrum</i> (a) plant and (b) dried fruits	58
3.8	Picture of Cinnabar	60
3.9	Pictures showing Purification of Cinnabar using lemon juice.	65
3.10	Picture of Crude Sulphur.	66
3.11	Picture depicting the purification of sulphur.	69
5.1	Transverse section of <i>Aconitum ferox</i> roots.	130
5.2	Diagnostic features for the powder microscopy of the roots of <i>Aconitum ferox</i>. (1) cork cells; (2) vessels with alternate pitting; (3) vessels with helical thickening; (4) Fibres.	130
5.3	Transverse section of <i>Solanum indicum</i> roots.	131
5.4	Diagnostic features for the powder microscopy of the roots of <i>Solanum indicum</i> (1) vessels with simple pitting; (2) tracheids; (3) xylem parenchyma (4) Fibres.	131
5.5	HPTLC Chromatograms of <i>Aconitum</i> alkaloids (a) Image at 235 nm, (b) Image at 254 nm and (c) Image at 366 nm.	142
5.6	HPTLC Chromatograms of Alkaloid fractions A-1(a), A-2 (b), A-3 (c), A-4 (d), A-5 (e), A-6 (f).	143-144

5.7	IR Spectra of Aconitum Alkaloids	147
5.8	X-ray diffraction spectra of S-1(a), S-2 (b), S-3 (c), S-4 (d), S-5 (e).	150-151
5.9	Differential thermograms of Fig. X-ray diffraction pattern of S-1(a), S-2 (b), S-3 (c), S-4 (d), S-5 (e).	152-154
5.10	X-ray diffraction spectra of H-1 (a), H-2 (b), H-3 (c).	157
5.11	XRD spectra of B-1 (a), B-2 (b), B-3 (c), B-4 (d).	159-160
5.12	Differential thermograms of B-1 (a), B-2 (b), B-3 (c).	161-162
5.13	UV spectra of Aconitine (200µg/ml).	169
5.14	Representative HPLC chromatogram of standard Aconitine.	170
5.15	Effect of solvent used for extraction on the absorbance of the alkaloid fraction	171
5.16	Representative Chromatograms of Aconitine (10-100 µg/ml).	171
5.17	Calibration Curve of Aconitine.	172
5.18	Representative chromatogram of FORM-1 (F-1), FORM-2 (F-2) and FORM-3 (F-3).	176
5.19	Image of TLC plate before derivatization at 254nm.	180
5.20	Image of the TLC plate after Derivatization with Dragendroff's reagent at 500 nm.	180
5.21	Representative HPTLC chromatograms of (a)Aconitine Standard and (b)Form-1.	181
5.22	Excitation of Aconitine at 270 nm and emission at 505 nm.	182
5.24	Representative HPLC chromatohram of Aconitine.	186
5.25	Representative chromatogram aconitine in presence of 0.1 N HCl at different intervals of time.	186
5.26 (a)	Chromatographic separation of ACN (50 µg/mL) in stressed sample of the standard subjected to acid hydrolysis.	187
5.26 (b)	Chromatographic separation of ACN (50 µg/mL) in	187

	stressed sample of the standard subjected to alkali hydrolysis.	
5.26 (c)	Chromatographic separation of ACN (50 µg/mL) in stressed sample of the standard subjected to water hydrolysis.	188
5.26 (d)	Chromatographic separation of ACN (50 µg/mL) in stressed sample of the standard subjected to oxidation.	188
5.26 (e)	Chromatographic separation of ACN (50 µg/mL) in stressed sample of the standard subjected to thermal degradation.	189
5.26 (f)	Chromatographic separation of ACN (50 µg/mL) in alkaline stressed sample of the standard subjected to photolytic degradation.	189
5.26 (g)	Chromatographic separation of ACN (50 µg/mL) in stressed sample of the standard subjected to 75% RH at 30°C.	190
5.27	Chromatograms showing separation of standard ACONITINE (10-100 µg/mL).	190
5.28	Chromatogram showing separation of ACONITINE in degraded formulation.	191
5.29	Representative peak purity curve of ACN in peak purity analysis. The peak has a positive value of minimum peak purity index confirming the purity of the peak and specificity of the method developed in the presence of degradation products and other components.	193
5.30	UV spectra of Solanine.	1961
5.31	Representative HPLC Chromatograms of Solanine standard.	198
5.32	Calibration curve of solanine.	199
5.33	Representative chromatograms of solanine (10-100 µg/ml).	199
5.34	Representative chromatogram of FORM-1 (F-1), FORM-2 (F-2) and FORM-3 (F-3).	203

5.35	Image of the TLC plate before derivatization at 254nm	207
5.36	Image of the TLC plate after derivatization with Dragendroff at 500 nm.	207
5.37	Representative HPTLC Chromatograms of (a)Solanine standard and (b)Form-1.	208
5.38	Image of TLC plate before derivatization at 254nm.	213
5.39	Image of TLC plate at 366nm after derivatization.	213
5.40	Representative HPTLC Chromatograms of (a) Piperine standard and (b) Form-1.	213-214
5.41(a)	Chromatographic separation of PIN (75 µg/mL) in stressed sample of the standard subjected to acid hydrolysis.	215
5.41 (b)	Chromatographic separation of PIN (75 µg/mL) in stressed sample of the standard subjected to alkali hydrolysis.	215
5.41 (c)	Chromatographic separation of PIN (75 µg/mL) in stressed sample of the standard subjected to water hydrolysis.	216
5.41 (d)	Chromatographic separation of PIN (75 µg/mL) in stressed sample of the standard subjected to oxidation.	216
5.41 (e)	Chromatographic separation of PIN (75 µg/mL) in stressed sample of the standard subjected to thermal degradation.	217
5.41 (f)	Chromatographic separation of PIN (75 µg/mL)in alkaline stressed sample of the standard subjected to photolytic degradation.	217
5.41 (g)	Chromatographic separation of PIN (75 µg/mL) in stressed sample of the standard subjected to 75% RH at 30°C.	218
5.42	Representative chromatograms of Linearity study of Piperine.	219
5.43	Representative peak purity curve of PIN in peak purity analysis. The peak has a positive value of minimum peak	221

	purity index confirming the purity of the peak and specificity of the method developed in the presence of degradation products and other components.	
5.44	Representative chromatogram showing separation of Piperine in degraded formulation.	223
5.45	UV spectra of a-ACN (λ_{max}-223nm), b-SON (λ_{max}-218nm) and c-PIN (λ_{max}-343nm).	226
5.46	Representative Chromatogram of SON, PIN and ACONITINE.	226
5.47	Representative chromtograms of Ternary mixture for linearity study.	227
5.48	Representative chromatogram of (a)FORM-1 , (b)FORM-2 and (c)FORM-3 (F-3).	232-233
5.49	UV spectra of Aconitine (λ_{max}-223nm), Solanine (λ_{max}-218nm).	236
5.50	Calibration curve of Solanine.	237
5.51	Calibration curve of Aconitine	237
5.52	Representative Chromatograms of Solanine and Aconitine (100μg/ml)	238
5.53	Representative chromatogram of Solanine and Aconitine.	239
5.54	Representative Chromatograms of Form-1(F-1), Form-2 (F-2), Form (F-3).	242
5.55	Representative chromatograms of HPLC in linearity range.	245
5.56	Calibration curve of Piperine.	246
5.57	Representative Chromatograms of Solanine and Piperine Standards	248
5.58	Representative chromatogram of Form-1(F-1), Form-2 (F-2)and Form-3 (F-3).	250
5.59	Overlain spectra for Sulphur.	252
5.60	Effect of Hydrazine hydrate concentration.	252
5.61	Effect of Sodium hydroxide Concentration on absorbance	253

	of hydrazine sulphide.	
5.62	Continuous variation plot for Hydrazine sulphide.	254
5.63	Effect of Sonication time on the extractive value of Sulphur.	254
5.64	Effect of Toluene volume on the extractive value of sulphur.	255
5.65	Calibration plot of Sulphur.	256
5.66	Effect of pH on the fluorescence intensity (%).	260
5.67	Effect of reagent concentration on the fluorescence intensity (%).	260
5.68	Effect of temperature on the fluorescence intensity (%).	261
5.69	Excitation of Boron at 470 nm and emission at 570 nm.	262
5.70	Calibration curve of Boron using spectrofluorimetry.	263
5.71	Differential voltammogram of Alizarin red S.	266
5.72	Differential voltammograms of Boron-Alizarin red S complex.	266
5.73	Effect of Accumulation time on the peak current (Na).	267
5.74	Effect of Accumulation potential on the peak current (nA).	268
5.75	Effect of pH on the peak current (nA).	269
5.76	Calibration curve of Boron-alizarin red S using Voltammetry.	271
5.77	Calibration curve of Boron.	274
5.78	Calibration curve of Mercury.	278
5.79	Microscopic images of mice heart, kidney and liver illustrating effect of single dose administration of FORM1, FORM2 and FORM3	279
5.80	Effect of FORM1 on percentage viability of H9c2 cells at different time interval	281
5.81	Effect of FORM2 on percentage viability of H9c2 cells at different time interval	282
5.82	Effect of FORM3 on percentage viability of H9c2 cells at	283

	different time interval	
5.83	Effect of Formulations on (a) CK-MB (b) LDH (c) SGOT (d) ALKP (e) Uric acid.	287
5.84	Effect of Formulations on the Heart weight to body weight ratio.	288
5.85	Fig.5.85. Microscopic images of rat heart illustrating effect of pretreatment if FORM1, FORM2 and FORM3 in ISO induced and MI rats	289-290