

LIST OF PLATES

<u>Plate</u>	<u>Title</u>	<u>Page</u>
I.1	View of the Almora hill	xi
III.1	Porphyroblastic gneiss (Loc. Chaunsali)	45
III.2	Folded quartz veins (Loc. Chitai)	48
III.3	Relict F_1 fold in quartzose mica-schists (Loc. near Dyolidanda hill)	52
III.4	Mesoscopic F_1 reclined fold in flaggy quartzite (Loc. Khatiari)	55
IV.1	Crenulation cleavage (S_2) due to micro- folding (gneiss from Dyolidanda hill)	66
IV.2	Crinkling related to F_1 , in schists (Dyolidanda fold core)	71
IV.3	F_1 reclined fold in flaggy quartzite, micaceous layers showing axial-plane schistosity (Loc. south of Chaunsali)	73
IV.4	L_1 lineation of striation type in flaggy quartzite (Loc. Khatiari)	75
IV.5	L_1 ribbons folded by a small F_2 fold in quartzite (Loc. Phalsimi)	78
IV.6	L_1 lineation of ribbon type folded on F_2 (Loc. south of Kasardevi)	117
V.1	Texture of coarse-grained garnet mica-schist type (i)	132
V.2	Mica flakes showing tight folding of earlier cleavage	132
V.3	Texture of medium-grained garnet mica-schist type (ii)	134

V.4	Texture of garnet mica-schist type (iii)	134
V.5	Garnet grain in mica-schist showing rotational growth	138
V.6	Stauroilite crystal in garnet mica-schist	138
V.7	Crinkled garnet mica-schist with porphyroblasts of chlorite	141
V.8	Crinkled garnet mica-schist with static garnet	144
V.9	Texture of quartzose mica-schist	146
V.10	Texture of graphitic schist showing relicts of earlier schistosity	148
V.11	Plagioclase in augen gneiss showing a porphyroblastic tendency	152
V.12	Basal section of tourmaline with inclusions of quartz and plagioclase	154
V.13	Texture of augen gneiss with augen of plagioclase (Chaunsali band)	157
V.14	Porphyroblast of potash felspar in porphyroblastic gneiss. Relict carlsbad twin is seen (Chaunsali band)	160
V.15	Augen of potash-felspar from augen gneiss of Dyolidanda band showing superimposition of cross hatching over carlsbad twinning	164
V.16	Porphyroblast of Potash-felspar in the porphyroblastic gneiss of Dyolidanda hill	166
V.17	Texture of gneissic granite	168
V.18	Rims of quartz grains along borders of felspar porphyroblast in gneissic granite	170
V.19	Perthitic veins in orthoclase of gneissic granite	170

V.20	Porphyroblastic muscovite after sericite	173
V.21	Texture of quartzite type (i)	176
V.22	Texture of quartzite type (ii)	176
V.23	Porphyroblasts of actinolite and garnet with inclusions of quartz and showing a late growth	178
VII.1	Relict quartzite layer within gneisses (Loc. Dyolidanda hill)	201

LIST OF TABLES

<u>Table</u>		<u>Page</u>
V.1	2Vx values of K-felspars in gneissic granites of Dyolidanda hill	171
VI.1	Chemical analyses of mica-schists	187
VII.1	Chemical analyses of the gneissic rocks of Dyolidanda hill band	205
VII.2	Chemical analyses of the gneissic rocks of Chaunsali band	208
VII.3	Chemical analyses of the gneissic rocks of Dyolidanda hill band (for Von Wolf's Q-L-M diagram)	211
VII.4	Chemical analyses of the gneissic rocks of Dyolidanda hill band (for Hejtman's diagram)	214
VII.5	Chemical analyses of the gneissic rocks of Dyolidanda hill band (for ACF and AKF diagrams)	228

-xi-

PLATE I.1

View of the Almora hill

