

REFERENCES



REFERENCES

- Bhan, S.K., (1972). Geological Mapping in parts of Banaskantha, Mehsana and Sabarkantha Districts, Gujarat. Unpublished G.S.I. Report.
- Binns, R.A., (1962). Metamorphic pyroxenes from the Broken Hill District, New South Wales, Mineral. Mag. Vol. 33, pp. 320-338.
- Bowes, D.R., Wright, A.E. and Park, R.G., (1964). Layered intrusive rocks in the Lewisian of the Northwest Highlands of Scotland; Quart.J.Geol.Soc. London., Vol.120, pp 153-192.
- Carey, S.W., (1954). The Rheid Concept in geotectonics, Jour. Geol. Soc. Australia., Vol.1 (for 1953), pp. 67-117.
- Cooray, P.G., (1960). Khondalites and charnockites of the Laggala Pallegma Area, Ceylon. Bull.Mys. Geol.Assn. Vol.18, pp. 3-54.
- , (1962). Charnockites and their associated gneisses in the Pre-Cambrian of Sri Lanka, Quart. Jour. Geol. Soc. London, Vol.18, pp.239-273.
- , (1965). Geology of the country around Alutgama, Ceylon, Geol.Surv.Dept., Mem.3, pp.1-111.
- , (1972). Hornblende garnet granulites from Hatton, Ceylon, Geol.Mag. Vol.109, No. 1, pp.37-44.
- Crawford, A.R., (1970). The Pre-Cambrian geochronology of Rajasthan and Bundelkhand, Northern India, Can. Jour.Earth Sci., Vol.7, No.1, pp.91-110.
- Davis, B.F., (1974). A layered basic complex in the Lewisian, South of Loch Laxford, Sutherland, U.K., Jour. Geol. Soc. London, Vol.130, pp. 279-284.
- Deb, M., (1979). Polymetamorphism of ores in Pre-Cambrian stratiform massive sulfide deposits at Ambaji-Deri, Western India, Mineral.Deposita (Berlin) Vol.14, pp.21-31.
- Dempster, T.J., Harrison, T.N., Brown P.E. and Hutton, D.H.W., (1991). Low pressure granulites from the Ketilidian mobile belt of Southern Greenland., Jour. Petrol., Vol. 32, No.5, pp. 979-1004.

- Desai, S.J., (1974). Petrological studies of High Grade Metamorphic rocks from an area between Balaram and Kanpura, Banaskantha district, Gujarat, India. Unpublished thesis for DIC, Imp. College of Science and Technology, London, p. 1-115.
- , Patel M.P. and Merh, S.S., (1978). Polymetamorphites of Balaram - Abu Road area north Gujarat and southwest Rajasthan Jour. Geol. Soc. India, Vol. 19, No.9, pp. 383-394.
- , and Patel M.P., (1986). A tectonic model of the polymetamorphic rocks of Balaram - Bhatana - Kapasia area southwest Rajasthan, north Gujarat Abst. Publ., Seminar on Evolution of Pre-Cambrian crust in Aravalli Mountain belt, Udaipur
- , and Patel, M.P., (1987). Occurrence of inliers of granulite facies rocks in amphibolite facies terrains of north Gujarat. Abst., Publ. in IV Gujarat Science Congress, Baroda. p.9.
- , and Patel, M.P., (1992). Granulitic rocks of the Banaskantha district of north Gujarat, India, in High Grade Metamorphics, Theophrastus Publ. S.A., pp. 103-121.
- Deshpande, G.G., Powar, K.B., Peshwa, V.V. and Sukhtankar, R.K., (1970). Occurrence of hornfels granulite transitional facies in the metasediments of Danta area, Gujarat, Jour. Indian Geol. Assoc., Vol.12, pp.72-82.
- Devaraju, T.C., Sadashivaiah, M.S., (1975). Structural geology of a granulite area in southern Karnataka. Jour. Karn. Uni., Vol. 20, pp.99-123.
- Dewaard, D., (1964). Mineral assemblages and metamorphic subfacies in the granulite facies terrane of the little Moose mountain syncline, south central Adirondack Highlands, Koninkl. Nederlandse Akad. Wetensch. Proc., Amsterdam, B, V.67, pp.344-362.
- , (1965). A proposed subdivision of the granulite facies, Am. Jour. Sci., Vol.263, pp.455-461.
- Drury, S.A., (1990). Spot Image data as an aid to structural mapping in the southern Aravalli hills of Rajasthan, India. Geol. Mag. Vol.,127, No.3, pp. 195-207.

- Eskola, P., (1939). Die metamorphen Gesteine, in Barth, T.F.W., Correns, C.W., and Eskola Pentti, Die Entstehung der Gesteine : Berlin, Julius Springer, pp. 263-407.
- , (1952). On the granulites of Lapland, Am. Jour. Sci. Bowen Vol. , pp. 135-171.
- Fareeduddin, Sharma, A.K. and Bose, V., (1991). Tectonic slices of high grade rocks in the Delhi fold belt of north central Rajasthan. Jour. Geol. Soc. Ind. 38, pp. 427-436.
- Fyfe, W.S., Turner, F.J. and Verhoogen, J., (1958). Metamorphic reactions and metamorphic facies, Geol. Soc. Am. Mem. 73.
- Gable, D.J. and Sims, P.K., (1969). Geology and metamorphism of some high grade cordierite gneisses, Front Range, Colorado. Geol. Soc. Am. spec. paper 128, p.87.
- Gandhi, S.M., Paliwal, H.V. and Bhatnagar, S.M., (1984). Geology and ore reserve estimates of Rampura - Agucha zinc lead deposits dist. Bhilwara Rajasthan, Jour. Geol. Soc. India. V.25, pp.689-705.
- Green, D.H. and Ringwood, A.E., (1967). An Experimental investigation of the gabbro to eclogite transformation and its petrological applications. Geochem. et. Cosmochimica Acta., Vol.31, pp.767-833.
- Griggs, D.T. and Handin, J., (1960). Observations on fracture and a hypothesis of earth quakes, in Rock Deformation : Geol. Soc. Am. Mem. 79, pp. 347-364.
- Groves, A.W., (1935). The charnockite series of Uganda, British East Africa, Quar. Jour. Geol. Soc. London. Vol. 91, pp. 150 - 207.
- Hapuarachchi, D.J.A., (1967). Hornblende granulite subfacies mineral assemblages from areas in Ceylon. Geol. Mag., Vol. 104, No.1, pp. 29-34.
- Harris, N.B.W., Holt, R.W. and Drury, S.A., (1982). Geobarometry, geothermometry and late Archean geotherms from the granulite facies terrains of south India, Jour. Geol., V.90, pp. 509-527.
- Heron, A.M. and Ghosh, P.K., (1938). The geology of Palanpur, Danta and part of Idar states. Rec. Geol. Surv. India, 79, pp.367-412.

- Heron, A.M., (1953). The geology of central Rajputana, Mem. Geol. Surv. India, 72, No.4, pp.367-411.
- Hietanen, A., (1943). Uber das Grundergebirge des Kanfantegebirges in Sudwestlichen Finland Acad. Fennicae; Serie A., III,6; Com.Geol. Bull. 130.
- , (1947). Archean geology of the Turku dist. in southwestern Finland, Bull. Geol.Soc.Am.Vol.58, pp. 1019-1084.
- Kohler, A. and Raaz, F., (1951). Uber eine neue Berechnung und graphische Darstellung Von Geseinsanalysen : N. Jb. Miner. Mn., pp.247-263.
- Kranck, E.H., (1931). Petrologische Übersicht des Kustengebietes E.Von Helsingfors. Bull. Com. Geol. Finlande, Vol.89,p.67.
- Krishna Rao , J.S.R., (1992). Petrochemistry and metamorphism of metabasites from Kanjamalai, Salem dist. Tamilnadu, India, High Grade Metamorphics, Theophrastus Publications S.A., pp.141 - 156.
- Krupicka, J., (1973). Geology of Devon Island, Canadian Shield.
- Miyashiro, A., (1973). Metamorphism and metamorphic belts, George Allen and Unwin Ltd., London, p.492.
- Myers, J.S., (1973). Igneous structures and textures in the Major gap gava outcrop of the Fiskenaeset anorthosite complex, Rapp. Gronlands, Geol. Unders. No. 51, pp.47-53.
- Naha, K. and Halyburton, R.V., (1974b). Late stress system for conjugate folds and kink bands in the main Raialo syncline, Udaipur dist., Rajasthan, India, Geol. Soc. Am. Bull. 85, pp. 251-256.
- Nambiar, G.C., Bhaskar Rao, B., Parthasarathy R. and Fedkin, V.V., (1992). Geochemistry and genesis of charnockites and associated gneisses from northern Kerala, India, High Grade Metamorphics, Theophrastus Publications S.A., pp. 187-216.
- Parras, K., (1946). On the coarse grained garnet cordierite gneisses of south and southwest Finland Comm. Geol. Finland. Bull. 138, pp.1-7.

- , (1958). On the charnockites in the light of a highly metamorphic complex in south western Finland, Comm. Geol. Finland, Bull. 181, pp.
- Patel, M.P., Desai, S.J. and Merh, S.S., (1985). Structural and metamorphic studies of the Pre Cambrian rocks of Balaram-Swarup Ganj area of north Gujarat and SW Rajasthan, Proc. of Fifth session Ind. Geol. Cong., pp. 65-75.
- Powar, K.B. and Patwardhan, A.M., (1984). Tectonic evolution and base metal mineralisation in the Aravalli Delhi belt, India. PreCam. Res. Vol.25, pp.309-323.
- Prider, R.T., (1945). Charnockite and related cordierite bearing rocks from Dangin, Western Australia, Geol. Mag. Vol.82, pp. 145-172.
- Radhakrishna, B.P., (1954). On the nature of certain cordierite bearing granulites bordering Closepet granite of Mysore, Rec. Mys. Geol. Dept. Bull.21.
- Raja Rao, C.S., (1967). On the age of Pre-Cambrian groups of Rajasthan, Jour. Miner. Met. Fuels, Vol. 15, No.9, pp. 306-309.
- Rama Rao, B. and Krishnachar, T.P., (1937). Cordierite hypersthene granulites and their associated schistose rocks from Bidaloti, Mysore state, Rec. Mys. Geol. Dept. Vol. 36, pp. 40-71.
- Ramsay, J.G., (1967). Folding and Fracturing of Rocks, McGraw Hill Inc. P. 562.
- Reinhardt, E.W. and Skippen, G.B., (1970). Geol. Surv. Can. Rept. Activities Paper 70.1 Pt.B, pp. 48-54.
- Sadashiviah, M.S., Hanagodimath, R.S., (1973). Cordierite - hypersthene - anthophyllite granulite from Mysore State, Jour. Kam. Uni. Vol. XVIII, pp.58-69.
- Scheumann, K.H. and Huckenholz, G., (1961). Die cordieritgneise und grantgneise des Sächsischen Granulitgebirges : Neues Jahrb. Mineralogie Abh., Vol. 96, pp. 131-161.
- Schreurs, J. and Westra, L., (1986). The thermotectonic evolution of a Proterozoic, low pressure granulite dome, west Uusimaa, SW Finland, Contr. Miner. Petrol. Vol.93, pp. 236-250.

- Sen, S.,(1959). Mineralogenetic trends in the evolution of metamorphic rocks and origin of granulites of east Manbhum, India. Proc. Nat. Acad. Sci.Vol.25A, pp. 118-138.
- Sharma, N.L. and Purkayastha, S., (1935). The heavy minerals of the Erinpura granite and microgranite of Danta State (N.Gujarat) Proc. Ind. Acad. Sci. Vol.2, No.4 , pp.369 - 376.
- , Nandy, N.C., (1936). A note on the petrological classification of the basic intrusives of Danta State (N.Gujarat) Proc. Indian. Acad. Sci. Vol. 3 (B), pp. 366-376.
- Sharma, N.L., (1953). Problems in correlation of the Pre-Vindhyan igneous rocks of Rajasthan, Presid. Add. Ind. Sci. Cong. pt. IX:30.
- Sharma, R.S. and Narayan, V., (1975). Petrology of polymetamorphic schists from an Archean complex terrain southwest of Beawar, Rajasthan, India, Neues Jahrb. Mineralogie Abh., Vol. 124, pp.190-222.
- Sharma, R.S.,(1988). Patterns of metamorphism in the Pre-Cambrian rocks of the Aravalli mountain belt. Mem. Geol. Soc. India, No.7, pp.33-75.
- Shaw, D.M., (1954). Trace elements in pelitic rocks, pt. II, Bull. Geol. Soc. Am. Vol.65, pp.1167-1182.
- Sherbakov, I.B.,(1992).Granulites and charnockites of the Ukrainian Shield, High Grade Metamorphics, Theophrastus Publications S.A., pp. 157-186.
- Sims, P.K. and Gable, D.J., (1967). Petrology and structure of Pre-Cambrian rocks, Central City Quadrangle, Colorado, U.S. Geol. Surv., Prof. Paper 554-E, p. 56.
- Sutton, J. and Watson, J.,(1951). The Pre-Torridonian metamorphic history of the Loch Torridon and Scourie areas in the NW highlands and its bearing on the chronological classification of the Lewisian. Quar. J. Geol. Soc. London, Vol. 106, pp. 241-308.
- Sychanthavong, S.P.H. and Merh, S.S.,(1984). Proto Plate Tectonics : the energetic model for the structural, metamorphic and igneous evolution of the Pre-Cambrian rocks northwest Peninsular India. Spec. Publ. Geol. Surv. Ind. Vol.12, pp.419-457.

- Thompson, A.B., (1976b). Mineral reactions in pelitic rocks; II, calculation of sum P-T-X (Fe-Mg) phase reactions, Am. J. Sci. Vol.276, pp. 425-454.
- Turner, F.J., and Verhoogen, J., (1960). Igneous and Metamorphic Petrology, McGraw Hill Publ. p.694.
- Turner, F.J. (1968). Metamorphic Petrology Mineralogical and Field Aspects, McGraw Hill Publ., pp.403.
- Wager, R., (1938). studies in gneisgebirge des schwarzaltes IX, Akad. Wiss. Math. Nat. K1, 4. Abh.
- Waldmann, L., (1929). Ein Cordieritreicher kinzigit Vom Rieserferner Tonalit in Ostirol, Mitt. Wiener. Geol. Ges. Bd. 22.
- Watson, J. (1949). The Geology of The Lewisian of the Scourie Area, Unpublished Ph.D. Thesis, University of London.
- , (1973). Effects of reworking on high grade gneiss complexes. Phil. Trans. Roy.Soc.London,A. 273, pp. 443-455.
- Wilson, A.F., (1952). The charnockite problem in Australia, Sir Douglas Mawson Anniv. University of Adelaide, pp. 205-224.
- Windley, B.F., (1973). Crustal development in the Pre-Cambrian. Phil. Trans.Roy.Soc. London. A. 273, pp. 321-342.
- Winkler, H.G.F., (1965). Petrogenesis of Metamorphic Rocks, Springer Verlag, NewYork, Inc., p.344.
- Wynne-Edwards, H.R., (1957). Structure of the westport concordant pluton in the Greenville, Ontario, Jour. Geol. Vol. 65, pp.639-649.
- , (1963). flow Folding, Am.J.Sci. Vol.261, pp. 793-814.
- and Hay, P.W., (1963). Coexisting cordierite and garnet in regionally metamorphosed rocks from the west port area, Ontario, Can. Mineralogist, Vol. 7, pp.453-478.
- Wynne-Edwards, H.R., (1967). Westport map area, Ontario with special emphasis on the Pre-Cambrian rocks, Geol. Surv. Can. Mem. 346 p. 1-142.

Young, E.D., Anderson, J.L., Clarke, S.H. and Thomas, W.M. (1989). Petrology of biotite - cordierite - garnet gneiss of the McCullough Range, Nevada. 1. evidence for proterozoic low pressure fluid absent granulite grade metamorphism in the southern Cordillera., J. Petrol. Vol.30, pp. 39-60.

Zen, E-An., (1960). Am. Mineral. Vol.45, pp. 129-175.