ACKNOWLEDGEMENT

It gives me great pleasure to express my deep sense of gratitude towards Prof. K.N. TRIVEDI, Professor of Chemistry, Department of Chemistry, M.S. University of Baroda, for suggesting the problem and valuable guidance with encouragement throughout the work.

I wish to express my special thanks to Prof. P.K. Bhatta-charya, Head, Chemistry Department, M.S. University of Baroda for his keen interest during the course of this work.

Special thanks are due to Shri S.S. Madhava Rao, Chemistry Department, M.S. University of Baroda for PMR spectra and carrying out microanalysis, Shri K.R. Chokshi, Morton-Thiokol Inc., Woodstock, Illinois, U.S.A. for PMR and Mass Spectra and Smt. Sushila Amin for IR Spectra.

Thanks are also due to Manager, Sarabhai Research Centre, Wadiwadi, Baroda and Director, Malti-Chem Research Centre, Nandesari, Baroda for their kind permission to allow me to use the facilities at their laboratories.

I am indebted to my father, Shri Padmakar Chandratre, who has been the source of inspiration and energy, throughout this work, without whose encouragement, this work would have

been an uphill task. I also express my gratitude towards my father-in-law Shri P.L. Soman for his timely help in looking after my son, Saurabh, during the writing of this work.

I also thank all my friends and colleagues for their co-operation.

Finally, I must mention the name of my husband, SANJAY SOMAN, but for fhis patience, interest, encouragement and co-operation in all respects, it would have been an impossible task to complete this thesis. To him I owe the most gratitude of all.

S P. Chandrathe. -SHUBHANGI P. CHANDRATRE