ACKNOWLEDGEMENT

It is with a profound sense of indebtness that I wish to express my sense of gratitude to PROF. K.N. TRIVEDI, who not only suggested the problem but also provided meticulous and ever inspiring guidance at every stage of this endevour.

Thanks are due to Prof. P.K. Bhattacharya, Head, Chemistry Department, M.S. University of Baroda, for providing necessary facilities.

Thanks are due to Dr. F.M. Dean, Robert Robinson Laboratories, University of Liverpool, Livepool, England for C¹³ nmr and pmr spectra of some compounds.

Thanks are due to Dr. R.T. Pardasani, Department of Chemistry, University of Manchester for Mass spectra of some compounds.

I am greatful to Shri S.S. Madhava Rao for carrying out the micro-analysis of the compounds, recording of pmr spectra and for taking deep interest in my research work.

I am also thankful to Mr. Nerugaonker and Mr. V.K. Kulkarni, Analytical division, Malti-Chem Research Centre for pmr spectra.

I am very much thankful to CSIR, New Delhi for financial support.

I am thankful to Mr. Dinesh Bhavsar for carrying out the typing of the thesis.

I thank all my colleagues at Chemistry Department and Malti-Chem Research Centre for their helpful comments and encouragements.

Finally without the patience, encouragement, 'forbearence

ŝ