

ACKNOWLEDGEMENTS

It is with a profound sense of indebtedness that I wish to record my sense of gratitude to Dr.Sukh Dev, D.Sc., FNA, Director, Malti-Chem Research Centre, Nandesari, Baroda, who not only advanced the problem but provided an ever inspiring guidance at every stage during its progress. I am indeed fortunate in getting an opportunity to be in his research group and am very grateful for all his kindness. I shall always cherish the memories of my association with him.

Grateful thanks are also due to Dr.N.P.Damodaran for the benefit of many valuable suggestions and encouragements.

My warmest thanks are due to Dr.H.P.S.Chawla for stimulating discussions in the course of these studies.

I thank Dr.C.L.Bhagchandani and his staff for MASS, PMR, UV and IR spectra; Mr.V.K.Kulkarni for GLC data and Prof.K.N.Trivedi, Head, Department of Chemistry, M.S. University, Baroda for microanalysis.

I am thankful to Mr.T.R.Viswanath for an excellent typing of this thesis and Mr.Arun Dindorkar for his help in compiling the same. I thank all my colleagues for their whole-hearted co-operation.

Financial assistance by Malti-Chem Research Centre, Nandesari, Baroda is gratefully acknowledged.

Malti-Chem Research Centre
Nandesari 391 340
Baroda.

January 1979

R C Desai

R.C.Desai