

# C O N T E N T S

	<u>Pages</u>
Summary	1- 12
 <u>Part I</u>	
Formation of malonmono arylamides on partial hydrolysis of cyanacet arylamides using (a) polyphosphoric acid (b) 75 % sulphuric acid ; and preparation of malonmono arylacids.	
Theoretical	13- 35
Experimental	36- 47
 <u>Part II</u>	
Synthesis of 2,4-dihydroxyquinolines by cyclisation of malon arylacids and malonmono arylamides using polyphosphoric acid as the cyclising agent.	
Theoretical	48- 70
Experimental	71- 82
 <u>Part III</u>	
Formation of methylene bis-derivatives of acetoacet arylamides by means of sodium hydroxy methane sulphonate.	
Theoretical	83- 104
Experimental	105- 114
 <u>Part IV</u>	
Synthesis of bis-hydroxyquinoline methanes by cyclisation of methylene bis-(acetoacet arylamides) using acetic anhydride and concentrated sulphuric acid as the cyclising agent.	
Theoretical	115- 134
Experimental	135- 141

Part V

Studies in the absorption spectra of  
acetoacet arylamides, hydroxyquinolines  
and their methylene bis-derivatives.

Theoretical	...	...	...	142- 163
Experimental	...	...	...	164- 176

Part VI

Halogenation of 2-hydroxy-4-methyl-  
quinolines and 2,4-dihydroxy-  
quinolines.

Theoretical	...	...	...	177- 198
Experimental	...	...	...	199- 226

Part V

Absorption Plates ( 1 to 13 )	...	227- 232
-------------------------------	-----	----------