

# List of Figures

<b>1.1:</b> The nuclear phenomenon takes place based on the impact factor of the incoming projectile. Schematic representation of involved processes based on impact factor.....	3
<b>1.2:</b> Different stages of nuclear interaction when the projectile is bombarded on the target nucleus. ....	4
<b>1.3:</b> Schematic presentation of a transfer pick-up reaction. ....	6
<b>1.4:</b> Schematic presentation of a transfer stripping reaction.....	6
<b>1.5:</b> Schematic presentation of a transfer knock-out reaction.....	6
<b>1.6:</b> Schematic representation of the formation and decay of Compound Nucleus (CN).....	8
<b>1.7:</b> Energy spectra for charge particles and neutrons. Figure is taken from J. S. Lilley, Nuclear Physics – Principles and applications (John Wiley and Sons Ltd.....	9
<b>1.8:</b> Differential cross section for an outgoing particle for direct reaction and decay of compound nucleus.....	10
<b>1.9:</b> Cluster structure in some weakly bound stable and unstable nuclei.....	11
<b>1.10:</b> The detailing of the process when weakly bound projectile nuclei interact with target nuclei. A typical example taken over here is WBP ${}^6\text{Li}$ ( $\alpha$ +d). ....	13
<b>1.11:</b> Part of Segre chart containing weakly bound category of nuclei.....	13
<b>1.12:</b> Typical example of TA behavior for system ${}^{16}\text{O} + {}^{208}\text{Pb}$ .....	17
<b>1.13:</b> Typical example of BTA behavior for system ${}^6\text{Li} + {}^{208}\text{Pb}$ .....	17
<b>2.1:</b> Schematic block diagram of a typical pelletron accelerator tank .....	28
<b>2.2:</b> Deposition of ${}^{92}\text{Mo}$ during the run and internal view of the chamber after deposition respectively.....	32
<b>2.3:</b> Floating off the thin film and taking on SS target frame for target fabrication.....	32

<b>2.4:</b> Schematic illustration of $\Delta E$ -E telescope detector setup. $\Delta E$ is the thinner solid state/gas detector .....	34
<b>2.5:</b> Schematic block diagram of electronic setup corresponding to the experimental arrangement in the thesis.....	36
<b>2.6:</b> Illustrative example of the form of Wood-Saxon potential.....	38
<b>2.7:</b> There is integration over two densities and therefore it is called the double folding model.....	42
<b>2.8:</b> The linear segments of $W(r, E)$ . The figure is taken from reference .....	44
<b>2.9:</b> Represent the coordinates of arrangement and now the three body becomes two body.....	48
<b>3.1:</b> Photograph of mounting arrangement for detector telescopes (T1, T2, T3, and T4), Monitors (M1, M2), target ladder, and Faraday cup (outside the chamber).....	55
<b>3.2:</b> Particle $\Delta E$ -( $E+\Delta E$ ) spectrum at $E_{\text{lab}}=25.7\text{MeV}$ , $\theta_{\text{lab}}=40^\circ$ , for the system ${}^6\text{Li}+{}^{51}\text{V}$ .....	56
<b>3.3:</b> Elastic scattering angular distribution data fitting through Woods –Saxon potential .....	58
<b>3.4:</b> The sensitivity radius deduced from optical model analysis for the imaginary part for the system ${}^6\text{Li}+{}^{51}\text{V}$ at energies a) 14 MeV b) 12 MeV.....	59
<b>3.5:</b> Elastic scattering angular distribution data fitting through the microscopic double folding model.....	60
<b>3.6:</b> Left panel: Real and imaginary potentials at sensitivity radius $R=9.4\text{ fm}$ using optical model. dynamic polarization potential (DPP) ( $\Delta V$ , $\Delta W$ ) and bare + DPP, these calculations are obtained from CDCC calculations. Right Panel: Real ( $N_R$ ) and imaginary ( $N_I$ ) normalization factors from the microscopic double folding model. ....	63

<b>3.7:</b> Result of CDCC calculation for real ( $\Delta V$ ) and imaginary ( $\Delta W$ ) polarization potential around sensitivity radius (9.4 fm). .....	65
<b>3.8:</b> Real and Imaginary normalization factors for systems .....	67
<b>3.9:</b> (a) Reduced reaction cross section for targets from different mass regions with ${}^6\text{Li}$ . (b) The ratio of systematic radius $R_{\text{sys}}^i = 1.12A_i^{1/3} - 0.96 A_i^{-1/3} + 3.75$ fm to spherical radius $R_{\text{sp}}^i = 1.3 A_i^{1/3}$ fm is given in the inset. ....	68
<b>3.10:</b> A systematic reaction cross section for a variety of projectile categories viz. strongly bound, Weakly bound, and rare ion beam with target ${}^{51}\text{V}$ . ....	69
<b>4.1:</b> Particle $\Delta E$ -( $E+\Delta E$ ) spectrum at $E_{\text{lab}}=19.7\text{MeV}$ , $\theta_{\text{lab}}=30^\circ$ , for the system ${}^6\text{Li}+{}^{51}\text{V}$ . ....	77
<b>4.4:</b> Energy integrated angular distribution of $\alpha$ -particles at various energies (a) 25.7 MeV, (b) 22.7 MeV, (c) 19.7 MeV, and (d) 13.6 MeV at ${}^6\text{Li}+{}^{51}\text{V}$ . ....	81
<b>4.5:</b> Schematic diagram of coupling states included for Coupled Reaction Channel Calculations (CRC) for 1-n transfer.....	85
<b>4.6:</b> Schematic diagram of coupling states included for Coupled Reaction Channel Calculations (CRC) for 1-p transfer.....	86
<b>4.7:</b> The contribution of several possible reaction channels to direct $\alpha$ production viz. non-capture breakup, 1 p transfer , 1 n transfer , 1 d transfer , transfer + Non-capture breakup (NCBU) , and Incomplete fusion +transfer +NCBU .....	88
<b>4.8:</b> Experimentally deduced direct $\alpha$ - particle cross section for ${}^6\text{Li}$ with various targets. ....	90
<b>4.9:</b> A ratio showing relative production of $\alpha$ - production cross section to deuteron production cross section for various targets around the Coulomb barrier. ....	91

<b>5.1:</b> Schematic diagram of the GPSC chamber of the experimental setup for the experiment....	99
<b>5.2:</b> Schematic diagram of the inner view of GPSC chamber of the experimental setup for the experiment.....	100
<b>5.3:</b> An energy calibrated experimental spectra <b>(a)</b> for the system ${}^7\text{Li}+{}^{92}\text{Mo}$ at energy $E_{\text{lab}} = 35 \text{ MeV}$ and $\theta_{\text{lab}} = 27^\circ$ ; <b>(b)</b> the projection of the drawn box reveals the first few inelastic states of target ${}^{92}\text{Mo}$ .....	101
<b>5.4:</b> Elastic scattering angular distribution data fitting through Woods –Saxon potential (WSP) and São-Paulo Potential (SPP for system ${}^7\text{Li}+{}^{92}\text{Mo}$ and diamonds for system ${}^7\text{Li}+{}^{100}\text{Mo}$ .....	104
<b>5.5:</b> The sensitivity radius deduced from optical model analysis for the real and imaginary parts for the system ${}^7\text{Li}+{}^{100}\text{Mo}$ system at energies 21.5 MeV and 30 MeV respectively.....	105
<b>5.6:</b> Left panel: Real and imaginary potentials at sensitivity radius $R=9.58 \text{ fm}$ using optical model WSP . Right panel: Real and imaginary potentials at sensitivity radius using optical model SPP for system ${}^7\text{Li}+{}^{92}\text{Mo}$ .....	109
<b>5.7:</b> Left panel: Real and imaginary potentials at sensitivity radius $R=10.28 \text{ fm}$ using optical model WSP. Right panel: Real and imaginary potentials at sensitivity radius using optical model SPP. for system ${}^7\text{Li}+{}^{100}\text{Mo}$ .....	110
<b>5.8:</b> The CDCC calculations are demonstrated with and without including breakup coupling effects for the system ${}^7\text{Li}+{}^{92}\text{Mo}$ .....	112
<b>5.9:</b> The CDCC calculations are demonstrated with and without including breakup coupling effects for the system ${}^7\text{Li}+{}^{100}\text{Mo}$ .....	113

<b>5.10:</b> A Comparative study of the experimental cross section with CDCC calculated reaction cross section and breakup cross section for ${}^7\text{Li}+{}^{92,100}\text{Mo}$ systems. ....	114
<b>5.11:</b> The comparative plot for total fusion cross sections ( $\sigma_{\text{fus}}$ ) calculated by CCFULL and the total reaction cross sections ( $\sigma_{\text{R}}$ ) for the systems ${}^7\text{Li}+{}^{92}\text{Mo}$ , ${}^7\text{Li}+{}^{100}\text{Mo}$ , and ${}^6\text{Li}+{}^{100}\text{Mo}$ .....	117
<b>5.12:</b> Reduced reaction cross section for targets from lighter to heavy mass regions with ${}^7\text{Li}$ ..	118
<b>6.1:</b> Schematic diagram and inner view of GPSC chamber of the experimental setup for the experiment.....	127
<b>6.2:</b> Elastic scattering angular distribution data fitting through Woods –Saxon potential for system ${}^7\text{Li}+{}^{100}\text{Mo}$ . ....	130
<b>6.3:</b> The sensitivity radius deduced from optical model analysis for the real and imaginary parts for the system ${}^6\text{Li}+{}^{100}\text{Mo}$ system at energy 23 MeV.....	132
<b>6.3:</b> Real and imaginary potentials at sensitivity radius $R=10.97$ fm using optical model WSP (blue solid circle).....	134

