

LIST OF TABLES

| <u>TABLE NO.</u> | <u>TITLE</u> | <u>PAGE</u> |
|------------------|--|-------------|
| 4-1 | : Important physico-chemical results | 4-3 |
| 2(a) | : Changes in the soluble organic constituents | 8 |
| 2(b) | : Chlorophyll content | 8 |
| 3 | : Biological changes | 11 |
| 4 | : Changes in coliform group density and total colony counts. | 14 |
| 5 | : Morphological characteristics of bacteria isolated. | 17 |
| 6 | : Biochemical characteristics of Bacteria isolated. | 22 |
| 7 | : Types of bacteria isolated. | 26 |
| 8 | : Comparison between Bacterial phase I and algal phase II | 32 |
| 5 - 1 | : Cultural characteristics of the bacteria isolated from raw sewage and the viscous scum. | 5 - 9 |
| 2 | : Bio-chemical reactions of the bacteria isolated from raw sewage and the viscous scum. | 11 |
| 3 | : Bacterial types in raw sewage and viscous scum | 12 |
| 4 | : Zoogloea formation seen in different synthetic liquid media. | 15 |
| 5 | : Morphological and bio-chemical characteristics of the isolate classified as zoogloea spp. | 16 |
| 6 | : Comparative characteristics of the isolates from activated sludge, trickling filter and stored raw sewage. | 18 |
| 7 | : Sub-grouping of the zoogloea spp on the basis of nitrate reduction and ammonia oxidation. | 21 |
| 8 | : Important biochemical characteristics of the zoogloea spp. | 21 |
| 9 | : Comparative study of the zoogloea strains isolated from the viscous scum and those obtained from Dr.Freedman and Dr.Crabtree of U.S.A. | 23 |
| 10 | : Taxonomical characteristics of <u>zoogloea ramigera</u> . | 29 |

List of Tables (Cohtd.)

| <u>Table No.</u> | <u>Title</u> | <u>Page</u> |
|------------------|--|-------------|
| 5 -11 | Distribution of certain morphological and cultural types among the dominant activated sludge bacteria from Dias (1964) compared with those found in the viscous scum. | 5 - 37 |
| 12 | Distribution of some physiological characteristics among the dominant activated sludge bacteria compared with those found in the viscous scum formed during bacterial phase I | 38 |
| 13 | A comparison between the bacterial types in the seven different sludge samples (Dias 1964) and in the viscous scum formed during bacterial phase I. | 40 |
| 6 -1 | Microorganisms accompanying the zoogloeas. | 6 - 5 |
| 2 | Physico-chemical analysis of sewage samples drawn on 7-4-1966. | 7 |
| 3 | A comparative study of the viscous scums formed in raw sewage samples taken on 7-10-66 and 13-2-67 from the Indian Institute of Science, Bangalore; and Wadi Sewage Disposal Works, Baroda on 13-2-67. | 10 |
| 4 | Zoogloea colonies and protozoans formed in stored industrial wastes from Alembic Chemical Works, Baroda. | 11 |
| 5 | Zoogloea colonies and protozoans formed in stored sewage samples taken from Atladra Sewage Disposal Works, Baroda. | 13 |
| 6 | Samples drawn from Okhla Sewage Disposal Works, New Delhi on 14-1-1968. | 14 |
| 7 -1 | Microscopic examination of slimy growths collected from Okhla Sewage Disposal Works at New Delhi on 14-1-1968. | 7 -10 |
| 8 - 1 | Turbidity changes noted with different strains and mixture of bacteria. | 8 - 7 |
| 2 | pH changes noted with different strains and mixture of bacteria. | 8 |
| 3 | Changes in Relative stability noted with different strains and mixture of bacteria. | 9 |
| 4 | Changes in acid KMnO_4 (4 hrs.) value noted with different strains and mixture of bacteria. | 10 |
| 5 | Changes in orthophosphate content noted with different strains and mixture of bacteria. | 11 |
| 6 | Changes in Am-N content noted with different strains and mixture of bacteria. | 12 |

List of Tables (Contd.)

| <u>Table No.</u> | <u>Title</u> | <u>Page</u> |
|------------------|---|-------------|
| 8 - 7 | : Turbidity changes noted with different fast-growing and mixed algae with and without bacteria. | 8 - 21 |
| 8 | : pH changes noted with different fast growing and mixed algae with and without bacteria. | 22 |
| 9 | : Changes in Relative stability noted with different fast-growing and mixed algae with and without bacteria. | 23 |
| 10 | : Changes in phenolphthalein alkalinity value noted with different fast-growing and mixed algae with and without bacteria. | 24 |
| 11 | : Changes in acid KMnO_4 (4 hrs.) value noted with different fast-growing and mixed algae with and without bacteria. | 25 |
| 12 | : Changes in orthophosphate content noted with different fast-growing and mixed algae with and without bacteria | 26 |
| 13 | : Changes in Am-N content noted with different fast-growing and mixed algae with and without bacteria. | 27 |
| 14 | : Turbidity changes noted with different slow-growing and mixed algae with and without bacteria. | 34 |
| 15 | : pH changes noted with different slow-growing and mixed algae with and without bacteria. | 35 |
| 16 | : Changes in Relative stability noted with different slow-growing and mixed algae with and without bacteria. | 36 |
| 17 | : Changes in Phenolphthalein alkalinity value noted with different slow-growing and mixed algae with and without bacteria. | 37 |
| 18 | : Changes in acid KMnO_4 (4 hrs.) value noted with different slow-growing and mixed algae with and without bacteria. | 38 |
| 19 | : Changes in Orthophosphate content noted with different slow growing and mixed algae with and without bacteria. | 39 |
| 20 | : Changes in Am-N content noted with different slow-growing and mixed algae with and without bacteria. | 40 |
| 9 - 1 | : Effluents from the laboratory models of activated sludge and oxidation pond compared. | 9 -27 |