

ACKNOWLEDGEMENT

I am in duty bound to record my deep sense of gratitude to Dr.J.C.George, Professor of Zoology, M.S.Uni - versity of Baroda, for the suggestion of the problem and also for the constant guidance and encouragement, but for which this work would not have been completed. I am indebted to Dr.Suresh Sethna, Mr.A.N.Godbole and Mr.S.S.Lele of the Department of Chemistry, to Dr.C.V.Ramakrishnan, Mr.T.N.Sekhara Varma and Mr.W.M.Deshpande of the Department of Biochemistry and to Mr.Y.K.Sane of the Department of Botany for the generous help extended to me during the course of this work. It is a pleasure to thank Dr.K.K.Nair, Professor of Zoology, Wilson College, Bombay and all my colleagues in this department for helpful criticism and cooperation and Mr.V.K.Chari, Curator, Bombay Natural History Museum, for kindly identifying the bat used in this work. Finally I must acknowledge the generosity of the M.S.University of Baroda, in the award of a special grant for the purchase of chemicals and of E.I. du Pont de Nemours & Co.(Inc.), U.S.A., Atlas Powder Co., U.S.A., Carbide and Carbon Chemicals Co., U.S.A. and Sigma Chemical Company, U.S.A for the free gift of some of the chemicals used in this study.