

## Contents

S. No.	Title	Page No.
<b><u>Chapter I:</u></b>		
<b>Introduction</b>		
1.	Introduction	2
1.1.	Organic Pollutants	3
1.2	Inorganic Pollutants	3
1.2.1	Lead	4
1.3.	Dyes	5
1.3.1.	Methylene Blue	5
1.3.2	Malachite Green	5
1.3.3.	Methyl Orange	6
1.3.4.	Rhodamine 6G	6
1.3.5	Rhodamine B	6
1.4	Adsorption	7
1.5	Adsorbent	7
1.5.1	Magnetic Nanoadsorbents	8
1.5.1.1	Cyclodextrins	8
1.5.2.	Hydrogel	9
1.5.2.1.	Dextran	9
1.5.3.	Bioadsorbent	10
1.6	Sensing	10
	References	12-17
<b><u>Chapter II:</u></b>		
<b>Synthesis, characterization and application of <math>\beta</math>-cyclodextrin based magnetic nanoadsorbent for simultaneous adsorption of hydrophilic and hydrophobic dyes</b>		
2.1	Introduction	19
2.2.	Materials and Methods	21

2.2.1.	Materials	21
2.2.2	Synthesis of Super paramagnetic Nano-Adsorbent (SPNA)	22
2.2.3.	Characterization of SPNA	22
2.2.4.	Adsorption of Dyes	23
2.2.5.	Simultaneous Adsorption Studies of Hydrophilic and Hydrophobic Dyes	24
2.2.6.	Desorption Experiment and Reusability	24
2.3.	Results and Discussion	24
2.3.1	Synthesis of SPNA	24
2.3.2.	NMR Spectra Analysis	25
2.3.3.	FTIR Spectra Analysis	27
2.3.4.	X-Ray Diffractogram Analysis	28
2.3.5.	HR-TEM Analysis	29
2.3.6.	Dynamic Light Scattering Analysis	30
2.3.7.	VSM Analysis	31
2.3.8.	TGA Analysis	32
2.3.9.	BET Analysis	33
2.3.10.	Selectivity of Adsorption (Effect of Cavity of Cyclodextrin)	34
2.3.11.	Effect of SPNA dosage	35
2.3.12.	Effect of contact time	36
2.3.13.	Effect of pH	36
2.3.14.	Effect of Initial Concentration	38
2.3.15.	Effect of Temperature	38
2.3.16.	Adsorption isotherms	39
2.3.17.	Adsorption Kinetics	42
2.3.18.	Simultaneous Adsorption Studies of Hydrophilic and Hydrophobic Dyes	43
2.3.19.	Desorption and Recyclability	44
2.3.20.	Adsorption Mechanism	45
2.3.21.	Comparison with Other Adsorbents	47
2.4.	Conclusion	48

References	49-56
Published Paper	57

### **Chapter III:**

#### **The application of dextran based hydrogel for elimination of organic dyes and reduction of nitrophenols**

3.1	Introduction	59
3.2	Materials and Methods	61
3.2.1	Materials	61
3.2.2.	Synthesis of Hydrogel (DEX-HMDI)	61
3.2.3.	Characterization of DEX-HMDI	62
3.2.4.	Measurements of Swelling Characteristics	63
3.2.4.1.	Swelling kinetics	63
3.2.4.2.	pH and Salt sensitivity	63
3.2.4.3.	Water Retention Capacity	63
3.2.5.	Adsorption of Dyes	64
3.2.6.	Desorption Experiment and Reusability	65
3.2.7.	Preparation of Ag@Hydrogel	65
3.2.8.	Reduction of 4-NP Catalyzed by Ag@Hydrogel	65
3.3.	Result and Discussion	66
3.3.1	Characterization Techniques	66
3.3.1.1.	Solid State NMR Spectra Analysis	66
3.3.1.2.	FTIR Spectra Analysis	67
3.3.1.3.	XRD Analysis	67
3.3.1.4.	EDS Analysis	68
3.3.1.5.	Thermogravimetric Analysis	69
3.3.2.	Swelling behavior of the hydrogel	70
3.3.2.1.	Deionized water	70
3.3.2.2.	Different pH values	70
3.3.2.3	Different concentration of saline solutions	70
3.3.3.	Water retention profile of the hydrogel	71

3.3.4.	Effect of various parameters on adsorption of dyes	71
3.3.4.1.	Effect of pH	71
3.3.4.2.	Effect of the Hydrogel Dosage	72
3.3.4.3.	Effect of Initial Concentration	72
3.3.4.4.	Effect of Temperature	72
3.3.4.5.	Effect of Contact Time	74
3.3.5.	Adsorption Isotherm	74
3.3.6.	Adsorption Thermodynamics	76
3.3.7.	Adsorption Kinetics	79
3.3.8.	Adsorption Studies in Binary Mixture of Dyes	81
3.3.9.	Desorption and Recyclability	82
3.3.10.	Adsorption Mechanism	84
3.3.11.	Comparison Table of other Hydrogels	86
3.3.12.	Valorization of Waste Expensive Metal Ions entrapped in the hydrogel matrix to Valuable Metal Nanoparticles for removal of Nitroaromatics in Water	88
3.3.13.	Reduction of Nitroaromatics and Sunlight Mediated Photocatalytic Capability of Ag@Hydrogel	90
3.4	Future perspectives for commercialization of hydrogel and remediation on a large scale	93
3.5.	Conclusion	96
	References	97-104
	Published Paper	105

#### **Chapter IV:**

#### **Development of crosslinked $\beta$ -cyclodextrin polymer functionalized gold nanosensor for detection of sulphur based amino acid and agrochemicals**

4.1.	Introduction	107
4.2.	Materials and Methods	109
4.2.1	Materials	109

4.2.2.	Characterization	109
4.2.3.	Synthesis of Au and Ag derived nano-sensors	109
4.2.3.1.	CDPA polymer synthesis	109
4.2.3.2.	Synthesis of Au derived functional nano-sensor AuNS@CDPA	110
4.2.3.3.	Synthesis of Ag derived functional nano-sensor AgNS@CDPA	110
4.2.4.	Preparation of Samples	110
4.2.5.	The colourimetric sensing ability of AuNS@CDPA and AgNS@CDPA towards sulfur-based compounds	111
4.3.	Result and Discussion	111
4.3.1.	Characterization of nano-sensors	112
4.3.1.1.	NMR Analysis	112
4.3.1.2.	FTIR Analysis	114
4.3.1.3.	UV-vis spectrophotometric determination	114
4.3.1.4.	HR-TEM Analysis	116
4.3.1.5.	DLS and Zeta potential	118
4.3.1.6.	FESEM-EDX Analysis	118
4.3.2.	Colourimetric Detection of sulfur-based compounds	118
4.3.2.1.	Sensing of Amino Acids	118
4.3.2.2.	Selective sensing of Cysteine in a mixture of amino acids	121
4.3.2.3.	Sensing of Agrochemicals	121
4.3.2.4.	Selective sensing of sulfur-based Agrochemicals in a mixture	124
4.3.3.	Effect of pH and NaCl concentration	124
4.3.4.	Natural and real sample analysis	127
4.3.5.	Selectivity and Sensitivity of AuNS@CDPA and AgNS@CDPA as a visible colourimetric detection system in a mixture of analytes	128
4.3.6.	Analytical Performance	129
4.3.7.	Mechanism	130
4.4.	Conclusion	135
	References	136-142
	Published Paper	143

---

**Chapter V (A):****Assessment of seaweed bioadsorbent (*Fucus vesiculosus*) for removal of methylene blue and rhodamine B dyes**

<b>5.1.1.</b>	Introduction	<b>145</b>
<b>5.1.2.</b>	Materials and Methods	<b>147</b>
<b>5.1.2.1.</b>	Materials	<b>147</b>
<b>5.1.2.2.</b>	Preparation of <i>Fucus vesiculosus</i> Seaweed Bioadsorbent (FVSB)	<b>147</b>
<b>5.1.2.3.</b>	Characterization of FVSB	<b>147</b>
<b>5.1.2.4.</b>	Adsorption of Organic Dyes	<b>148</b>
<b>5.1.2.5.</b>	Simultaneous Adsorption of MB and RB	<b>149</b>
<b>5.1.2.6.</b>	Desorption Experiment and Reusability	<b>149</b>
<b>5.1.3.</b>	Results and Discussion	<b>149</b>
<b>5.1.3.1.</b>	Characterization of Bioadsorbent	<b>149</b>
<b>5.1.3.1.1.</b>	FTIR Spectra Analysis	<b>149</b>
<b>5.1.3.1.2.</b>	SEM-EDX Analysis	<b>150</b>
<b>5.1.3.1.3.</b>	TGA Analysis	<b>152</b>
<b>5.1.3.2.</b>	Effect of FVSB Dosage	<b>153</b>
<b>5.1.3.3.</b>	Effect of Initial Concentration	<b>153</b>
<b>5.1.3.4.</b>	Effect of pH	<b>154</b>
<b>5.1.3.5.</b>	Effect of Contact Time	<b>156</b>
<b>5.1.3.6.</b>	Effect of Temperature	<b>156</b>
<b>5.1.3.7.</b>	Adsorption Isotherms	<b>157</b>
<b>5.1.3.8.</b>	Adsorption Kinetics	<b>160</b>
<b>5.1.3.9.</b>	Adsorption Thermodynamics	<b>161</b>
<b>5.1.3.10.</b>	Simultaneous Adsorption Study of MB and RB	<b>163</b>
<b>5.1.3.11.</b>	Desorption and Regeneration	<b>164</b>
<b>5.1.3.12.</b>	Adsorption Mechanism	<b>166</b>
<b>5.1.3.13.</b>	Adsorption of real sample using FVSB	<b>167</b>
<b>5.1.3.14.</b>	Comparison with Other Adsorbents	<b>168</b>
<b>5.1.4.</b>	Conclusion	<b>169</b>
	References	<b>170-177</b>

---

## **Chapter V (B):**

### ***Citrus limetta* derived eco-friendly bioadsorbent for efficient elimination of organic dyes and heavy metal ions**

<b>5.2.1.</b>	<b>Introduction</b>	<b>179</b>
<b>5.2.2.</b>	<b>Materials and Methods</b>	<b>181</b>
<b>5.2.2.1.</b>	<b>Materials</b>	<b>181</b>
<b>5.2.2.2.</b>	<b>Synthesis of Modified <i>Citrus limetta</i> (MCL)</b>	<b>181</b>
<b>5.2.2.3.</b>	<b>Characterization of MCL</b>	<b>182</b>
<b>5.2.2.4.</b>	<b>Adsorption of organic and inorganic pollutants</b>	<b>183</b>
<b>5.2.2.5.</b>	<b>Simultaneous Adsorption of MB and Pb<sup>2+</sup></b>	<b>184</b>
<b>5.2.2.6.</b>	<b>Desorption Experiment and Reusability</b>	<b>184</b>
<b>5.2.3.</b>	<b>Result and Discussion</b>	<b>184</b>
<b>5.2.3.1.</b>	<b>Characterization of Bioadsorbent</b>	<b>184</b>
<b>5.2.3.1.1</b>	<b>FTIR Spectra Analysis</b>	<b>184</b>
<b>5.2.3.1.2.</b>	<b>SEM-EDX Analysis</b>	<b>186</b>
<b>5.2.3.1.3.</b>	<b>BET Analysis</b>	<b>187</b>
<b>5.2.3.1.4.</b>	<b>TGA Analysis</b>	<b>188</b>
<b>5.2.3.1.5.</b>	<b>DLS Analysis</b>	<b>189</b>
<b>5.2.3.1.6.</b>	<b>XRD Analysis</b>	<b>190</b>
<b>5.2.3.2.</b>	<b>Effect of MCL Dosage</b>	<b>191</b>
<b>5.2.3.3.</b>	<b>Effect of Initial Concentration</b>	<b>191</b>
<b>5.2.3.4.</b>	<b>Effect of pH</b>	<b>193</b>
<b>5.2.3.5.</b>	<b>Effect of Contact Time</b>	<b>194</b>
<b>5.2.3.6.</b>	<b>Effect of Temperature</b>	<b>195</b>
<b>5.2.3.7.</b>	<b>Adsorption Isotherms</b>	<b>196</b>
<b>5.2.3.8.</b>	<b>Adsorption Kinetics</b>	<b>199</b>

5.2.3.9.	Adsorption Thermodynamics	200
5.2.3.10.	Simultaneous Adsorption Study of MB and Pb <sup>2+</sup>	203
5.2.3.11.	Desorption and Regeneration of MCL	204
5.2.3.12.	Adsorption Mechanism	205
5.2.3.13.	Adsorption of environmental sample using MCL bioadsorbent	206
5.2.3.14.	Comparison with Other Bioadsorbents	207
5.2.4.	Conclusion	208
	References	210-218
	Published papers, book chapters, conferences and seminars	219-232

### List of Figures

Figure No.	Title	Page Number
2.1	Schematic for synthesis of SPNA	24
2.2	NMR spectra for CD-MA	25
2.3	<sup>1</sup> H NMR spectra of Maleic Anhydride	26
2.4	<sup>1</sup> H NMR spectra of β- cyclodextrin polymer	26
2.5	FTIR spectra of MNPs, β-CD, MA, and SPNA	27
2.6	XRD pattern for fresh SPNA and recycled SPNA	29
2.7	HR-TEM images for (A) Pure MNPs and (B) & (C) SPNA	30
2.8	DLS profile for SPNA	31
2.9	Room temperature magnetization curves of SPNA (Inset: Photograph showing separation of SPNA using an external magnet)	31
2.10.	Thermo Gravimetric Analysis curve of SPNA	32
2.11	Nitrogen adsorption–desorption isotherm of SPNA	33
2.12	Nitrogen adsorption-desorption isotherm of pristine Iron Oxide	34

Nanoparticles		
<b>2.13</b>	Effect of SPNAs dosage on the percentage removal of 30mgL <sup>-1</sup> of MG, 20 mgL <sup>-1</sup> of MB and R6G at 25 °C and pH-7.0	<b>35</b>
<b>2.14</b>	Effect of contact time on the percentage removal of 30mgL <sup>-1</sup> of MG, 20 mgL <sup>-1</sup> of MB and R6G at 25 °C and pH-7.0	<b>36</b>
<b>2.15</b>	Effect of pH on the percentage removal of 30mgL <sup>-1</sup> of MG, 20 mgL <sup>-1</sup> of MB and R6G at 25 °C	<b>37</b>
<b>2.16</b>	Effect of initial concentration on adsorption of dyes on SPNA	<b>38</b>
<b>2.17</b>	Effect of temperature on dye removal efficiency at 35 °C and 45 °C	<b>39</b>
<b>2.18</b>	Langmuir adsorption isotherm for the removal of 30mgL <sup>-1</sup> of MG, 20 mgL <sup>-1</sup> of MB and R6G at 25 °C pH 7.0	<b>40</b>
<b>2.19</b>	Freundlich adsorption isotherm for the removal of 30mgL <sup>-1</sup> of MG, 20 mgL <sup>-1</sup> of MB and R6G at 25 °C pH 7.0	<b>41</b>
<b>2.20</b>	(A) Zero order (B) First Order (C) Second Order (D) Third Order (E) Pseudo first order (F) Pseudo Second Order kinetic curves for MB, MG and R6G	<b>42</b>
<b>2.21</b>	Percentage removal of dyes after successive desorption/adsorption cycles	<b>44</b>
<b>2.22</b>	FTIR spectra of the recycled adsorbent	<b>45</b>
<b>2.23</b>	Interaction of the dyes MB, MG and R6G with SPNA	<b>46</b>
<b>3.1</b>	Schematic description of process for synthesis of dextran derived Hydrogel	<b>62</b>
<b>3.2</b>	Solid state <sup>13</sup> C NMR of Hydrogel	<b>66</b>
<b>3.3</b>	FTIR spectra of Dextran and Hydrogel	<b>67</b>
<b>3.4</b>	X-ray diffractograms of Dextran and hydrogel	<b>68</b>
<b>3.5</b>	SEM images and elemental mapping via EDX in (i) dextran and (ii) hydrogel	<b>69</b>
<b>3.6</b>	Thermogravimetric Curves for Hydrogel	<b>69</b>
<b>3.7</b>	Swelling characteristics of hydrogel in (A) distilled water (B) buffer solutions of pH (2-10) (C) KCl solutions (D) water retention profile	<b>71</b>
<b>3.8</b>	Effect on removal efficiency of MB and MO by varying parameters (A) pH; Inset: Photographs of vial before and after adsorption, pH 12 for MB	<b>73</b>

	and pH 2 for MO (B) adsorbent dosage (C) initial concentration (D) at 35°C, 45°C and 55°C temperature (E) contact time (For all the experiments 100 mgL <sup>-1</sup> dye solution was and 0.1 g of hydrogel was used)	
<b>3.9</b>	Adsorption isotherms for MB and MO uptake; (A & B) Langmuir and (C) Freundlich model fitting	<b>75</b>
<b>3.10</b>	Thermodynamic plot for removal of (A) MB and (B) MO using hydrogel	<b>77</b>
<b>3.11</b>	Thermodynamic surface plots for (A) MB and (B) MO uptake by hydrogel	<b>78</b>
<b>3.12</b>	Graphs of (A) zero order (B) first order (C) second order (D) third order (E) pseudo first order (F) pseudo second order kinetic models for adsorption of MB and MO on hydrogel (conditions: initial concentration 100 mgL <sup>-1</sup> , adsorbent dosage 0.1 g at room temperature)	<b>80</b>
<b>3.13</b>	Assessment of hydrogel's adsorption behaviour in a binary mixture of dyes (A) UV-vis spectra recorded before and after adsorption at different time intervals digital (Inset: images depicting color changes prior to mixing, after mixing of dyes and post adsorption.) (B) %R <sub>e</sub> for MB and MO individually from mixture (Conditions: initial concentration of binary solution 25 mgL <sup>-1</sup> , adsorbent dosage 0.1 g at room temperature)	<b>81</b>
<b>3.14</b>	(A) Studies for desorption of dyes in different eluents (B) assessment of recyclability of the hydrogel	<b>83</b>
<b>3.15</b>	Characterizations of recycled hydrogel (A) (i) X-ray diffractograms and (ii) FTIR spectra (B) EDX spectra of adsorbent samples with dyes adsorbed on its surface (i) MB and (ii) MO	<b>84</b>
<b>3.16</b>	Probable mechanism for adsorption of cationic and anionic Dyes onto hydrogel depicting some possible interactions	<b>85</b>
<b>3.17</b>	Schematic for preparation of valorized metallopolymer Ag@Hydrogel	<b>88</b>
<b>3.18</b>	Characterizations of valorized metallopolymer (A) SEM image and EDS spectrum showing presence of Ag (B) appearance of SPR band due to formation of Ag nanoparticles observed by UV-vis spectrophotometric determination (C) XPS and (D) DLS spectrum	<b>89</b>
<b>3.19</b>	General scheme for the reduction of nitroaromatics and UV-vis	<b>91</b>

	spectrophotometric determinations for reduction of varying concentrations of 4-NP using Ag@Hydrogel	
<b>3.20</b>	Probable mechanism for reduction of 4-NP to 4-NA over valorized catalyst Ag@Hydrogel	<b>92</b>
<b>3.21</b>	(A) FT-IR overlay of fresh and recycled metallopolymer (B) Recycling studies	<b>93</b>
<b>3.22</b>	Process flow diagram for commercial production of dextran derived Hydrogels	<b>94</b>
<b>3.23</b>	Process flow diagram for practical application of Ag@Hydrogel for nitroaromatics reduction	<b>95</b>
<b>4.1</b>	Schematic for synthesis of cyclodextrin phthalate ester (CDPA) polymer and subsequent entrapment of Au/Ag nanoparticles to form the nanosensor	<b>112</b>
<b>4.2</b>	(A) <sup>1</sup> H NMR spectrum of CDPA polymer (B) overlay of FTIR spectra (i) CDPA polymer, (ii) AuNS@CDPA and (iii) AgNS@CDPA	<b>113</b>
<b>4.3</b>	UV-vis determinations demonstrating spectral properties of synthesized nanoparticles (A) AuNS@CDPA and (B) AgNS@CDPA (inset images of cuvettes containing sensor solutions)	<b>115</b>
<b>4.4</b>	Characterization via HRTEM imaging for (A)AuNS@CDPA (B) AgNS@CDPA; comparison of the hydrodynamic size of both sensors via (C) DLS measurements (D) zeta-potential graph and (E) elemental mapping of the sensors	<b>117</b>
<b>4.5</b>	(A) Photographs and (B) Absorption spectra of AuNS@CDPA in absence and presence of various amino acids and their mixture	<b>119</b>
<b>4.6</b>	(A) Photographs and (B) Absorption spectra of AgNS@CDPA in absence and presence of amino acids (Cysteine, Proline, Glycine, Aspartic Acid, Tyrosine and Mixture)	<b>120</b>
<b>4.7</b>	(A) Photographs of AuNS@CDPA solution in presence of different agrochemical compounds. The colour change is observed only in the case of sulfur-containing compounds (B-D) Comparative UV-visible spectra of AuNS@CDPA in a mixture of sulfur and non-sulfur agrochemicals	<b>122</b>

<b>4.8</b>	(A) Photographs and (B) Absorption spectra of AgNS@CDPA in absence and presence of agrochemicals (PDO, SDO, DST, DAHO, SHC, DPA, ATC, SMB, TGA and SDDC)	<b>123</b>
<b>4.9</b>	UV-visible spectra demonstrating effect of (A) NaCl concentration and (B) pH values in presence/absence of Cysteine. The graph of absorption ratio ( $A_{670}/A_{524}$ ) versus NaCl concentration/pH shows the optimum sensing conditions	<b>125</b>
<b>4.10</b>	UV-visible spectra demonstrating effect of (A) NaCl concentration and (B) pH values in presence/absence of SDDC. The graph of absorption ratio ( $A_{670}/A_{524}$ ) versus NaCl concentration/pH shows the optimum sensing conditions	<b>126</b>
<b>4.11</b>	(A) UV-vis spectra and (B) Absorption ratio ( $A_{670}/A_{524}$ ) of AuNS@CDPA in presence of allicin containing onion and garlic extract. (C)Photographs of AuNS@CDPA and AgNS@CDPA in presence of sulfur based pesticide samples and their (D&E) Absorption spectra. Linear relationship of (F) Cysteine and (G) SDDC with absorbance at 670 nm for quantification and determination of LOD (Sensing samples were prepared by adding 700 $\mu$ L of AuNS@CDPA solution with 300 $\mu$ L of different concentration of Cysteine and SDDC, the error bars represent standard deviation obtained from three independent measurements)	<b>127</b>
<b>4.12</b>	Schematic showing a probable mechanism for detection of (A) Cysteine and (B) SDDC using AuNS@CDPA	<b>132</b>
<b>4.13</b>	FTIR overlay spectra for AuNS@CDPA in presence of (i) Cysteine and (ii) SDDC	<b>133</b>
<b>4.14</b>	Various analytical characterization of sensor in presence of analytes; HRTEM image of (A) Cysteine added AuNS@CDPA (B) SDDC added AuNS@CDPA (C) Cysteine containing AgNS@CDPA and (D) SDDC containing AgNS@CDPA; (E) & (F) DLS spectra of Cysteine and SDDC containing sensor solutions; elemental mapping of (G) Cysteine and (H) SDDC added AuNS@CDPA	<b>134</b>
<b>5.1.1</b>	Preparation of bioadsorbent FVSB	<b>147</b>

<b>5.1.2</b>	FTIR spectra of FVSB, FVSB+MB and FVSB+RB	<b>150</b>
<b>5.1.3</b>	(I) SEM images of (A) FVSB, (B) FVSB+MB and (C) FVSB+RB; (II) EDS curves of (A) FVSB, (B) FVSB+MB and (C) FVSB+RB	<b>151</b>
<b>5.1.4</b>	Thermogravimetric curves for FVSB	<b>152</b>
<b>5.1.5</b>	Effect of FVSB dosage on the percentage removal of 100mg/L of MB and RB at room temperature	<b>153</b>
<b>5.1.6</b>	Effect of initial concentration of MB and RB on adsorption capacity and removal efficiency using FVSB at room temperature	<b>154</b>
<b>5.1.7</b>	Effect of pH on removal efficiency of MB and RB on FVSB at room temperature	<b>155</b>
<b>5.1.8</b>	Effect of contact time on the percentage removal of 100mgL <sup>-1</sup> of MB and RB at pH-8.0 and pH- 4.0, respectively at room temperature	<b>156</b>
<b>5.1.9</b>	Effect of temperature on removal efficiency of MB and RB with increasing temperature from 25 °C to 45 °C	<b>157</b>
<b>5.1.10</b>	(A&B) Langmuir and (C) Freundlich adsorption isotherm for the removal of 100mgL <sup>-1</sup> of MB and RB at room temperature	<b>159</b>
<b>5.1.11</b>	Graphs of (A) Zero order (B) First Order (C) Second Order (D) Third Order (E) Pseudo first order (F) Pseudo Second Order kinetic curve for MB and RB adsorption on FVSB	<b>160</b>
<b>5.1.12</b>	ln (Q <sub>e</sub> /C <sub>e</sub> ) versus 1/T graph for estimation of thermodynamic parameters for adsorption of MB and RB on FVSB	<b>162</b>
<b>5.1.13</b>	(A) Photographic images of binary solution before and after adsorption (B) UV-vis spectra of MB and RB in binary solution with time	<b>163</b>
<b>5.1.14</b>	(A) Percentage removal of MB and RB after successive desorption/adsorption cycles using 0.1M HCl as eluent; (B) FTIR spectra of regenerated FVSB	<b>165</b>
<b>5.1.15</b>	Probable mechanism of adsorption of MB and RB on FVSB	<b>166</b>
<b>5.1.16</b>	Photographic images and UV visible spectra of environmental sample adsorption using FVSB	<b>167</b>
<b>5.2.1</b>	Synthetic scheme of MCL	<b>181</b>
<b>5.2.2</b>	FTIR spectra for MCL, MCL+MB and MCL+Pb <sup>2+</sup>	<b>185</b>

<b>5.2.3</b>	(I) SEM images of (A) MCL, (B)MCL+MB and (C) MCL+Pb <sup>2+</sup> ; (II) EDS curves of (A) MCL, (B) MCL+MB and (C) MCL+Pb <sup>2+</sup>	<b>187</b>
<b>5.2.4</b>	Nitrogen adsorption-desorption isotherm of MCL bioadsorbent	<b>188</b>
<b>5.2.5</b>	Thermogravimetric curves for MCL bioadsorbent	<b>189</b>
<b>5.2.6</b>	DLS size distribution diagram of MCL bioadsorbent	<b>189</b>
<b>5.2.7</b>	XRD spectra of MCL, MCL+Pb <sup>2+</sup> and MCL+MB	<b>190</b>
<b>5.2.8</b>	Effect of MCL dosage on the percentage removal of 100mg/L of MB and Pb <sup>2+</sup> at 35°C	<b>191</b>
<b>5.2.9</b>	Effect of initial concentration of MB and Pb <sup>2+</sup> on adsorption capacity and removal efficiency using MCL at 35°C	<b>192</b>
<b>5.2.10</b>	(A) point zero charge (pH pzc) estimation, (B) Effect of pH on removal efficiency of MB and Pb <sup>2+</sup> on MCL at 35°C	<b>193</b>
<b>5.2.11</b>	Effect of contact time on the percentage removal of 100mgL <sup>-1</sup> of MB and Pb <sup>2+</sup> at pH-6.0 and pH- 8.0, respectively at 35°C	<b>195</b>
<b>5.2.12</b>	Effect of temperature on dye removal efficiency with increasing temperature from 35 °C to 55 °C	<b>196</b>
<b>5.2.13</b>	(A&B) Langmuir and (C) Freundlich adsorption isotherm for the removal of 100mgL <sup>-1</sup> of MB and Pb <sup>2+</sup> at 35°C	<b>198</b>
<b>5.2.14</b>	Graphs of (A) Zero order (B) First Order (C) Second Order (D) Third Order (E) Pseudo first order (F) Pseudo Second Order kinetic curves for MB and Pb <sup>2+</sup> adsorption on MCL	<b>199</b>
<b>5.2.15</b>	(A) In (Q <sub>e</sub> /C <sub>e</sub> ) versus 1/T graph for estimation of thermodynamic parameters for adsorption of MB and Pb <sup>2+</sup> , (B&C) Thermodynamic surface plots of MB and Pb <sup>2+</sup> , respectively	<b>202</b>
<b>5.2.16</b>	(A) UV-vis spectra of MB in binary solution with time and (B) Removal Efficiency of MB and Pb <sup>2+</sup> with time in binary solution	<b>203</b>
<b>5.2.17</b>	(A) Desorption study of MB and Pb <sup>2+</sup> with different eluents; (B) Percentage removal of MB and Pb <sup>2+</sup> after successive desorption/adsorption cycles using HCl as eluent; (C) FTIR spectra of regenerated MCL after adsorption of MB and Pb <sup>2+</sup>	<b>205</b>

<b>5.2.18</b>	Probable mechanism of adsorption of MB and Pb <sup>2+</sup>	<b>206</b>
<b>5.2.19</b>	UV visible spectra of environmental sample adsorption using MCL bioadsorbent	<b>207</b>

### List of Tables

<b>Table No.</b>	<b>Title</b>	<b>Page Number</b>
<b>2.1</b>	BET Parameters of SPNA	<b>33</b>
<b>2.2</b>	Isotherm constants for the adsorption of 30mgL <sup>-1</sup> of MG, 20 mgL <sup>-1</sup> of MB and R6G at 25 °C pH 7.0	<b>41</b>
<b>2.3</b>	Elemental composition of adsorbent determined before and after adsorption following 4 adsorption/desorption cycles	<b>47</b>
<b>2.4</b>	Comparison of maximum adsorption capacities of SPNA with other adsorbents	<b>47</b>
<b>3.1</b>	Isotherm Constants for the Adsorption of 100mgL <sup>-1</sup> of Methylene Blue and 100 mgL <sup>-1</sup> of Methyl Orange	<b>75</b>
<b>3.2</b>	Thermodynamic Parameters for adsorption of dyes onto the surface of hydrogel at three different temperatures	<b>79</b>
<b>3.3</b>	Elemental Composition of Recycled Hydrogel following Adsorption/Desorption Cycle	<b>86</b>
<b>3.4</b>	Comparison of Removal Efficiencies of Hydrogel with Literature Reports	<b>86</b>
<b>3.5</b>	Elemental Composition of Ag@Hydrogel	<b>90</b>
<b>3.6</b>	Reduction of 4-NP at Different Concentrations using Ag@Hydrogel as Catalyst	<b>92</b>
<b>4.1</b>	Optimization for preparation of AuNS@CDPA	<b>114</b>
<b>4.2</b>	Optimization for preparation of AgNS@CDPA	<b>115</b>
<b>4.3</b>	A comparison table of various Au/Ag derived nanosensors employed for colourimetric detection of cysteine/pesticides	<b>129</b>

<b>5.1.1</b>	Elemental composition of FVSB before and after adsorption of MB and RB	<b>152</b>
<b>5.1.2</b>	Adsorption isotherm constants for 100 mgL <sup>-1</sup> of MB and RB at room temperature	<b>158</b>
<b>5.1.3</b>	Kinetic parameters for adsorption of MB and RB on FVSB	<b>161</b>
<b>5.1.4</b>	Thermodynamic parameters for the adsorption of MB and RB on FVSB	<b>163</b>
<b>5.1.5</b>	Comparison of maximum adsorption capacities of FVSB with other adsorbents	<b>168</b>
<b>5.2.1</b>	Elemental composition of MCL bioadsorbent before and after adsorption of MB and Pb <sup>2+</sup>	<b>186</b>
<b>5.2.2</b>	Isotherm constants for the adsorption of 100mgL <sup>-1</sup> of MB and Pb <sup>2+</sup> at 35 °C	<b>197</b>
<b>5.2.3</b>	Kinetics parameters for adsorption of MB and Pb <sup>2+</sup> on MCL	<b>200</b>
<b>5.2.4</b>	Thermodynamic parameters for the adsorption of MB and Pb <sup>2+</sup> on MCL	<b>201</b>
<b>5.2.5</b>	Comparison of maximum adsorption capacities of MCL with other adsorbents	<b>206</b>